


Petrous Bone Cholesteatoma: Facial and Hearing Preservation

Ear, Nose & Throat Journal
2021, Vol. 0(0) 1–8
© The Author(s) 2021
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/01455613211056554
journals.sagepub.com/home/ear


Annalisa Pace, MD¹ , Irene Claudia Visconti, MD², Giannicola Iannella, MD²,
Alessandro Milani, MD², Valeria Rossetti, MD², Salvatore Cocuzza, MD³,
Antonino Maniaci, MD³ , Daniela Messineo, MD⁴ and Giuseppe Magliulo, MD² 

Abstract

Objectives: Petrous bone cholesteatoma (PBC) is a rare condition of the petrous portion of the temporal bone. Treatment of choice consists of radical surgical removal, paying attention to protect the facial nerve and inner ear as far as possible. The aim of the present study was to evaluate the efficacy of modified translabyrinthine techniques in preserving hearing function and the use of the adjuvant endoscopic techniques in a group of PBC patients. **Methods:** This study comprised 16 cases of PBCs surgically treated in our Department. Pre- and post-operative hearing status was assessed with pure tone audiometry and speech discrimination and graded according to the Gardner- Robertson classification system. Facial function was based on the House Brackman (HB) classification. PBCs were grouped using Sanna's classification. The choice of surgical technique was based on the above findings together with preoperative evidence. Post-operative follow-up ranged from 1 to 10 years and also included Computed Tomography and Magnetic Resonance Imaging assessment. **Results:** PBCs were classified as follows: 37.5% infralabyrinthine; 43.75% supralabyrinthine; and 18.75% massive. Preservation of the bone conduction threshold was feasible in 62.5% of patients. For supralabyrinthine PBCs a subtotal petrosetomy was performed in all cases and hearing preservation was possible in 57% of them: an adjuvant endoscopic approach was performed in 43%. Infralabyrinthine PBCs were treated using a modified translabyrinthine approach with preservation of bone conduction in 83% of patients; an adjuvant endoscopic approach was performed in 50% cases. One patient with a massive cholesteatoma was treated by modified translabyrinthine approach, preserving a serviceable level of hearing. In all massive cases, an adjuvant endoscopic approach was performed. In 2 patients with preoperative palsy, facial nerve function showed an improvement. The follow-up period revealed evidence of limited recurrence at CT imaging in 2 patients. **Conclusions:** The introduction of modified surgical approaches, able to preserve the anatomical-functional structures, have shown an improvement of post-operative hearing outcomes.

Keywords

petrous bone cholesteatoma, facial nerve, hearing preservation, modified translabyrinthine approach, endoscopic surgery

Introduction

Petrous temporal bone cholesteatoma (PBC) is a rare disease developing in the petrous portion of the temporal bone with an incidence ranging from 4 to 9% of all petrous pyramid lesions.¹ Since it has a slow and silent growth, it may produce severe functional damage, reducing patients' quality of life.

Radical surgery of PBCs is performed by means of different techniques based on the location and extension of the lesion, respecting the facial nerve and inner ear.² The main complications of PBC surgery are facial nerve palsy and severe hearing loss, depending on their proximity to the disease.³ Since PBC

¹Scienze Chirurgiche Department, Sapienza University of Rome, Italy

²Organi di Senso Department, Sapienza University of Rome, Italy

³Otorinolaringoiatria Department, University of Catania, Italy

⁴Scienze Radiologiche, Oncologiche e Anatomo-Patologiche Department, Sapienza University of Rome, Italy

Corresponding Author:

Giuseppe Magliulo, MD, Organi di Senso Department, Sapienza University of Rome, Via Gregorio VII n.80, Rome 00165, Italy.

Email: giuseppe.magliulo@uniroma1.it



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons

Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE

and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

involves the otic capsule, it comes into contact with the membranous labyrinth structures: cochlea (anterior labyrinth) and the vestibular system (posterior labyrinth).⁴

Over the last years, advances in neuroradiology and modern surgical techniques have made it possible to remove PBCs more safely, with minimal recurrence, and perioperative morbidity.

Sanna et al classified PBCs into 5 subtypes (supralabyrinthine, infralabyrinthine, infralabyrinthine-apical, apical, and massive) depending on their localization and defined possible surgical approaches.⁵

According to many authors, otic capsule and hearing preservation are impossible to achieve in most supralabyrinthine or infralabyrinthine PBCs, due to the high risk of recurrence.^{6,7}

In cases of supralabyrinthine PBCs,⁵ the middle cranial fossa approach is preferred to the transotic one in order to maintain the otic capsule and the preoperative level of serviceable hearing, even though this implies a higher incidence of recurrence.

Infralabyrinthine PBCs are usually treated by translabyrinthine and transotic approaches with resulting anacusia. Hence, Fish proposed the infratemporal fossa approach as a hearing preservation technique.⁸

In the recent literature, few studies have reported good hearing preservation outcomes after PBC surgery.^{9–13} On the contrary, good hearing outcomes have been reported after vestibular Schwannoma surgery performed using modified translabyrinthine techniques.^{14,15}

Adoption of intraoperative endoscopic procedures is generally accepted, but their use in PBC surgery is mentioned in only a few papers.

The aims of this study were to evaluate the efficacy of modified translabyrinthine techniques in preserving hearing function and the use of adjuvant endoscopic techniques in a group of patients affected by PBCs. The outcomes and the possible disadvantages of these methods are discussed.

Patients and methods

This retrospective study described a series of 16 patients with petrous bone cholesteatoma who were referred to our Department for surgery between 2008 and 2020. This research was performed in accordance with the principle of the Declaration of Helsinki and approved by the local Ethics Committee of the University “Sapienza,” Rome.

Patients' characteristics

The database was analyzed for presenting signs and symptoms, surgical details, and post-operative complications. A cohort of 23 consecutive patients, surgically treated for PBCs, was reported in the database. The inclusion criterion was related to hearing status that was assessed with pure tone audiometry (PTA) and speech discrimination. The degrees of hearing preservation and loss were defined according to the Gardner–Robertson classification system.¹⁶ Sixteen patients belonging to Class I–III of this

classification (serviceable in class I and II and with a potential improvement with hearing aid in class III) were enrolled.

Facial function was based on the House Brackman (HB) classification.¹⁷

Preoperative Computed Tomography (CT) showed the extension of the disease. Topographically, the lesions were grouped using Sanna's classification.^{2,5}

When a PBC was observed, a temporal bone magnetic resonance was performed to differentiate it from other lesions such as cholesterol granuloma, mucocele and temporal bone tumors.

Surgical management

The choice of surgical technique was based on the database and the preoperative conditions of each patient; all definitive surgical procedures were performed by the senior author (GM).

In cases of supralabyrinthine cholesteatoma with serviceable hearing, subtotal petrosectomy with blind closure of the EAC was performed, while in those with non-serviceable hearing a transotic approach was preferred (Table 1).

In cases of infralabyrinthine cholesteatoma with serviceable hearing, a modified translabyrinthine approach was chosen, while in other cases subtotal petrosectomy with blind closure of EAC was performed (Table 1).

Modified translabyrinthine approach

The modified translabyrinthine technique consisted of a cutaneous incision similar to that used in the traditional one.¹⁵ After drawing the periosteal flap back and exposing the cortical mastoid, total mastoidectomy was performed. All of the mastoid cells were removed, exposing the medial cranial fossa tegmen, the sinodural angle, and the dura of the posterior cranial fossa at the front and back of the sigmoid sinus, which was skeletonized with exposure of the jugular bulb. The three semicircular canals were identified from the horizontal canal as the starting point. The bone covering of the semicircular canal is removed using a diamond burr. This passage is performed keeping the dissection area as wet as possible and avoiding suction of the fluid around the labyrinth, which may violate the membranous labyrinth. When the posterior bony labyrinth has been completely removed, the area of the posterior, horizontal, and upper semicircular canals is occluded with bone wax.

Particular care must be taken to close off completely the access to the vestibule of the semicircular canals. Maintaining the vestibule anatomically intact and isolating it completely is fundamental for successful preservation of hearing function. If the cholesteatoma reaches the internal auditory channel, the bone around the vestibule must be scrupulously removed until the “blue line” of the internal auditory canal can be identified. The bone between the jugular bulb and the resected semicircular canals must be removed.

Table 1. Comparison between hearing preservation techniques for PBCs.

Surgical Technique	Indications	Advantages	Disadvantages
Subtotal petrosectomy	Supralabyrinthine and infralabyrinthine PBCs with/without serviceable hearing	✓Preservation of the facial nerve and inner ear function	✓Conductive hearing loss
Middle fossa approaches	Supralabyrinthine PBCs with serviceable hearing	✓Preservation of the facial nerve and inner ear function	✓Invasive: Craniotomy; temporal lobe retraction ✓Not applicable in infralabyrinthine PBCs
Modified translabyrinthine approaches	Infralabyrinthine and some massive PBCs with serviceable hearing	✓Preservation of the facial nerve and inner ear function ✓Poorly invasive: not needed craniotomy or mandibular involvement	✓Needed of the intact membranous labyrinth
Infratemporal fossa approach A-B-C	Infralabyrinthine PBCs with serviceable hearing	✓Preservation of the facial nerve and inner ear function ✓Appropriate control of the vertical and horizontal intrapetrous internal carotid artery	✓Conductive hearing loss. ✓Invasive: Sacrifice of the mandibular nerve; disarticulation of the mandibular condyle. ✓High risk of recurrence

In the event of a semicircular canal fistula being identified, the cholesteatoma matrix was isolated and gently raised, exposing the membranous labyrinth. Firstly, the membranous labyrinth with the matrix was excised using a sickle knife and then the opening in the labyrinth was rapidly closed with bone wax. For the time it remained open, the membranous labyrinth was kept wet with Ringer's solution and no suction was applied close to the openings of the semicircular canals to protect the anatomy and function of the membranous labyrinth of the vestibule. Using a cottonoid, the closure of the fistula was then reinforced with bone paste and temporalis fascia. This allowed preservation of the vestibule, sealing the openings of the semicircular canals.¹⁸

Subtotal petrosectomy

Subtotal petrosectomy is defined as a complete exenteration of all pneumatic cell tracts of the temporal bone with closure of the isthmus of the Eustachian tube. The first step is retroarticular temporal incision followed by a blind sac closure of the EAC. The skin of the auditory canal is totally removed from the lateral bony canal. Mastoidectomy is performed exposing the medial cranial fossa tegmen, the sinodural angle and the dura of the posterior cranial fossa at the front and back of the sigmoid sinus, which was skeletonized with exposure of the jugular bulb. The facial nerve is identified and preserved, removing the posterior canal wall and following the passage of the Fallopian canal from the oval window to the stylomastoid foramen.

Then temporal pneumatic cells are removed (retrosigmoid, retrofacial retrolabyrinthine, supralabyrinthine, infralabyrinthine, pericarotid, and supratubal). Finally, the middle ear is obliterated with abdominal fat and fibrin glue.

Endoscopic adjuvant approach

For endoscopic adjuvant surgery, a rigid endoscope with an angulation of 45°, a length of 14-cm, and an outer diameter of 3 mm and 4 mm (Storz, Germany) was used. It was connected to a camera head (Storz, Germany) and a high-definition monitor positioned in front of the surgeon. It allows better visualization and removal of the matrix of the cholesteatoma from the dura of the middle and posterior cranial fossa and near the posterior labyrinth.

Follow-up

Post-operative follow-up ranged from 1 to 10 years (average of 3.9 years) and included CT and Magnetic Resonance Imaging (MRI) to estimate cholesteatoma recurrence. The Gardner–Robertson classification system and facial nerve function were evaluated at 3–6–12 months; after the first year of follow-up, they were repeated once a year.

Results


Sixteen patients (9 men and 7 women, with a median age of 49.6 years and a range of 13–83 years) presented an extensive petrous bone cholesteatoma in one of their ears. 10 patients had left-sided disease, while in 6 it involved the right side (Table 2).

All the cholesteatomas were acquired: 11 patients underwent primary surgery, while 5 had previously undergone otologic surgery in other structures.

Otomicroscopic exams revealed 5 epitympanic perforations, 4 subtotal or total perforations; 3 normal tympanic membranes associated with swelling in their inferior portion; 2 canal wall up tympanoplasty, 1 canal wall down tympanoplasty, and 1 radical tympanoplasty.

Table 2. Patients characteristics.

Pt n	Age	Sex	Side	Symptoms	Route	Gardner-Robertson Hearing Scale		Facial Nerve Function	
						Preop	Postop	Preop	Postop
1	83	M	L	Tinnitus, otorrhea, hearing impairment	SUPRAL.	II	IV	I HB	I HB
2	13	F	R	Otorrhea, hearing impairment	MASSIVE	II	II	I HB	I HB
3	48	F	L	Tinnitus	SUPRAL.	I	I	I HB	I HB
4	46	M	R	Hearing impairment, headache	INFRAL.	II	III	I HB	I HB
5	62	M	L	Hearing impairment	SUPRAL.	II	II	I HB	I HB
6	16	M	L	Hearing impairment, otorrhea, aural fullness	INFRAL.	II	II	I HB	I HB
7	57	F	R	Hearing impairment, tinnitus	SUPRAL.	II	IV	I HB	I HB
8	61	M	L	Hearing impairment, otorrhea, otalgia	INFRAL.	II	II	I HB	I HB
9	50	M	R	Hearing impairment, aural fullness	SUPRAL.	III	III	I HB	I HB
10	39	F	L	Hearing impairment	INFRAL.	II	II	I HB	I HB
11	29	F	L	Hearing impairment, vertigo	SUPRAL.	II	II	I HB	I HB
12	77	F	R	Hearing impairment, otorrhea, tinnitus	SUPRAL.	III	V	V HB	III HB
13	41	M	L	Hearing impairment	INFRAL.	III	IV	I HB	I HB
14	74	F	L	Hearing impairment, otorrhea, aural fullness	MASSIVE	III	V	III HB	II HB
15	53	M	R	Hearing impairment, headache	MASSIVE	III	V	I HB	I HB
16	44	M	L	Hearing impairment, otorrhea, vertigo	INFRAL.	II	II	I HB	I HB

**Figure 1.** (a) CT scan examination of massive petrous bone cholesteatoma in contiguity with the jugular bulb and involving the dura mater; and (b) CT scan examination of massive petrous bone cholesteatoma in contiguity with the jugular bulb and the dura mater.

Hearing impairment was the most common symptom, present in 94% of patients (15/16) (Table 2). Preoperative Gardner–Robertson hearing evaluation evidenced 6.3% Class I (1/16); 62.5% Class II (10/16); and 31.25% Class III (5/16) (Table 2).

Facial palsy was present in 12.5% (2/16): one patient had a grade III HB for 4 months and the other had grade V HB for 7 months (Table 2).

PBCs were categorized according to Sanna's classification: 6/16 (37.5%) were infralabyrinthine cholesteatomas; 7/16 (43.75%) were supralabyrinthine cholesteatomas, and 3/16 (18.75%) massive ones (Figure 1).

Preservation of the bone conduction threshold was possible in 62.5% of patients (10/16) (Table 2).

In our series of supralabyrinthine PBCs subtotal petrosectomy was performed in all cases and hearing preservation was possible in 4 patients (57%) (Table 2). The endoscopic adjuvant approach was performed in 3 cases (43%).

Infralabyrinthine PBCs are usually removed by subtotal petrosectomy with labyrinthectomy or translabyrinthine or transotic approaches, sacrificing hearing. In our series of patients, they were surgically treated with the modified translabyrinthine approach and showed preservation of bone conduction in 83% of patients (5/6). The adjuvant endoscopic adjuvant approach was adopted in 3 cases (50%).

In massive PBCs hearing preservation is usually sacrificed to achieve radicality and to avoid recurrence. In our series, one patient presented a massive cholesteatoma that mainly




Figure 2. Massive petrous bone cholesteatoma: (1) jugular foramen; (2) jugular bulb; and (3) removal of the cholesteatoma from the dura mater.




Figure 3. Massive petrous bone cholesteatoma: (1) facial nerve; (2) sigmoid sinus; and (3) jugular bulb after removal of infralabyrinthine petrous bone cholesteatoma.

involved the posterior labyrinth and the internal auditory canal: hence a modified translabyrinthine approach was chosen and a serviceable level of hearing was preserved (Figures 2 and 3). In all massive cases, an adjuvant endoscopic approach was performed.

Intraoperative findings evidenced dehiscence of the facial nerve in 10/16 patients (62.5%): in 2 of these cases, the nerve was compressed but not infiltrated by the cholesteatoma (Table 3).

Partial erosion of the ossicular chain was observed in 3 patients, while 5 did not present any ossicular chain.

A fistula of the labyrinth was present in 62.5% of patients: in 19% (4/16) it was located in the anterior portion (cochlea), while in 37.5% (6/16) it was on the posterior side.

In one case of massive cholesteatoma, the lesion involved the internal auditory canal.

In the two cases with compression of the facial nerve, the cholesteatoma matrix was lifted off the nerve with a blunt dissector without requiring any graft.

Facial nerve function in two patients with preoperative palsy showed an improvement: 1 patient went to grade III HB from grade V HB; the other went to grade II HB from grade III HB (Table 2).

During surgery, no cochlear implant was positioned in anacusic patients. The same choice was made for bone conduction implants in patients with hearing preservation.

The follow-up period showed that 2 patients (12.5%) had a diagnosis of limited recurrence at CT imaging. Patients who constantly maintained their hearing presented a post-operative T2-weighted vestibule and cochlea signal (Figure 4). Two patients with hearing preservation underwent positioning of a Sophono device about two years later while three patients opted for a non-implanted bone conduction aid. None of the patients with anacusia opted for the cochlear implant.

Discussion

Petrous bone cholesteatoma (PBC) is an epidermoid cyst of the petrous portion of the temporal bone with a congenital or acquired etiopathogenesis. Its growth is slow and asymptomatic in the deep portion of the temporal bone and it is commonly misdiagnosed for years. It has an erosive expansion that may involve several important structures (inner ear, facial nerve, internal carotid artery, and dura) even reaching as far as the cerebellopontine angle.⁵

CT-scan imaging is the mainstay method since it displays isotropic sections and three-dimensional (3D) reconstructions of PBCs and their location in the temporal bone, thus providing fundamental information for surgical planning.¹⁹

Magnetic resonance (MR) imaging provides preoperative information for differential diagnosis with cholesterol granuloma, mucocele, and tumors. It detects small lesions by DWI thinner sections imaging and is also essential for identifying signs of recurrence during follow-up.¹⁹

Radical surgery of PBCs employs a variety of techniques whose target is to achieve removal of the entire cholesteatoma matrix with maximal preservation of nearby structures.² Surgical goals of all the procedures are radical excision of the disease; preservation of facial nerve function; prevention of CFS leakage; prevention of meningitis; preservation of hearing when feasible.²

In our series, 87.5% of patients did not have any long-term recurrence. This incidence is in line with the literature, despite the different approaches performed.⁶ Therefore, eradication of the disease may also be possible using less invasive techniques.

The facial nerve is a structure located in the temporal bone that may be involved by PBCs. Facial palsy is a common complication depending on compression and/or infiltration of the disease or may be a consequence of surgical removal procedures.

Table 3. Surgical management and Intraoperative findings.

Pt n	Route	Approach	Involved Inner Ear	Dura	FN	Recurrence
1	SUPRAL.	RSP	Fistula of Anterior Labyrinth	MF		None (1 year)
2	MASSIVE	MTL+ ENDOSCOPIC GUIDANCE	Fistula of Posterior Labyrinth; IAC	PF	Dehiscent tymp	None (2 years)
3	SUPRAL.	SP + ENDOSCOPIC GUIDANCE			Dehiscent GG	None (10 years)
4	INFRAL.	MTL	Fistula of Posterior Labyrinth		Dehiscent tymp	None (7 years)
5	SUPRAL.	RSP + ENDOSCOPIC GUIDANCE	Fistula of Anterior Labyrinth	MF	Dehiscent GG and tymp	None (4 years)
6	INFRAL.	MTL		MF		None (4 years)
7	SUPRAL.	RSP	Fistula of Anterior Labyrinth	MF	Dehiscent GG	None (1 year)
8	INFRAL.	MTL + ENDOSCOPIC GUIDANCE			Dehiscent tymp	None (1 year)
9	SUPRAL.	SP + ENDOSCOPIC GUIDANCE	Fistula of Posterior Labyrinth			None (3 years)
10	INFRAL.	MTL + ENDOSCOPIC GUIDANCE		PF	Dehiscent tymp	None (5 years)
11	SUPRAL.	RSP	Fistula of Anterior Labyrinth	MF		Limited (7 years)
12	SUPRAL.	SP			Compressed GG	Limited (2 years)
13	INFRAL.	MTL		PF		None (3 years)
14	MASSIVE	SP + ENDOSCOPIC GUIDANCE	Fistula of Posterior Labyrinth		Compressed GG and tymp	None (1 year)
15	MASSIVE	TOA + ENDOSCOPIC GUIDANCE	Fistula of Posterior Labyrinth	PF	Dehiscent GG and tymp	None (6 years)
16	INFRAL.	MTL + ENDOSCOPIC GUIDANCE	Fistula of Posterior Labyrinth	MF		None (6 years)




Figure 4. (a) CT scan examination of massive petrous bone cholesteatoma in contiguity with the dura mater; (b) CT scan evidenced the same massive PBC involving of internal auditory canal; (c) post-operative MRI axial images that evidenced of the persistence of intralabyrinthine fluid; (d) post-operative MRI: coronal images that evidenced of the persistence of intralabyrinthine fluid; and (e) MRI 3D reconstruction that showed the presence of fluid in post-operative labyrinthine.

The incidence of preoperative facial palsy^{6,20,21} reported in the literature averages 51%. However, in our experience, its incidence is decreasing. In 2007, Magliulo³ reported an incidence of 54% in patients with PBCs, treated at Sapienza University of Rome. From 2008 to 2020, in the same Department, preoperative facial palsy was observed in only 12.5% of PBC patients. This reduction may be ascribed to our protocol based on an early MRI diagnosis in patients with a suspect lesion.

Radical matrix removal is undoubtedly the aim of PBC surgery. Therefore, preservation of the otic capsule and hearing are secondary issues and depend on the extension of the disease and the possibility of preserving homeostasis of the anterior (cochlea) and posterior (vestibular) labyrinth (Table 1).

According to Sanna et al,⁵ selected patients with supralabyrinthine PBC and serviceable hearing, but without any

evidence of fistula, should be treated by combined middle fossa and trans-mastoid approach. However, this strategy is possible only in cases without any otic capsule involvement.

Prasad et al⁶ evidenced post-surgical anacusia in 74.4% of supralabyrinthine PBCs treated with these surgical approaches. In our series, only Gardner–Robertson Class I–III patients were enrolled, namely those likely to maintain a serviceable level of hearing with/without aid after surgery. Supralabyrinthine PBCs were removed by subtotal petrosectomy, and in 43% this approach was combined with an adjuvant endoscopic approach. None of the patients were treated via the middle cranial fossa or transotic approach. Hearing outcomes showed preservation of the bone threshold in 68.75% of patients.

Surgical strategies for infralabyrinthine PBCs are subtotal petrosectomy with labyrinthectomy, translabyrinthine,

and transotic approaches. Since they all sacrifice the otic capsule, in 1977 Fisch⁸ proposed the infratemporal fossa approach as a hearing preservation technique. The translabyrinthine approach, and its related anacusia, is preferred for PBCs with labyrinthine fistula or internal auditory canal involvement.

Several surgical approaches for hearing preservation have been reported in the field of vestibular Schwannoma surgery.

In 1991, McEleven et al¹⁴ proposed a modified translabyrinthine approach based on the destruction of all the semicircular canals and a concomitant sealing of the vestibule with bone wax. Sharma et al used this technique in 1 patient with PBC and reported a good long-term hearing outcome.

In 2004, Magliulo et al¹⁵ described a modified translabyrinthine technique to preserve hearing during vestibular schwannoma surgery, involving the internal auditory canal. It was a less invasive technique than McEleven's one because it spared the posterior labyrinth. In the recent literature, there are no reports of this technique being used for PBCs, although it was firstly performed to treat infralabyrinthine PBCs, associated with the endoscope in 50% of cases. Outcomes showed total hearing preservation in 5/6 patients (83%) including the case with internal auditory canal involvement. Several advantages of this approach emerged during follow-up: cholesteatoma was totally removed in all patients who underwent this method and none has shown signs of persistence or recurrence of the disease on radiological imaging.

The preservation of endolabyrinthine fluids in the vestibule is the basis of hearing preservation showed by a well-depicted post-operative T2-weighted labyrinthine signal (Figure 4). To date, no cases of recurrence have been reported.

Our data are in contrast with those reported by Moffat et al in 2008.²¹ In their report, 43 patients were categorized according to the Moffat–Smith classification, showing a prevalence of massive (massive labyrinthine and massive labyrinthine apical), and supralabyrinthine types. Preservation of the otic capsule was performed in only 14% of patients, even though the series included patients with normal hearing and patients with serviceable hearing. The authors preferred to sacrifice hearing to gain better access for removing the cholesteatoma.

Similar findings were reported by Zaabi et al,⁷ who described preservation of hearing after traditional PBC surgery in only 1 out of 6 patients. A limitation of their study was the lack of a PBC classification.

On the contrary, our results are in line with Senn et al,²² who reported 82% of hearing preservation in patients treated with traditional approaches. However, the percentage of infralabyrinthine PBCs was low (10%), despite the supralabyrinthine ones (67%). Moreover, they reported hearing preservation techniques with a high rate of recurrence (19%), while in our cases none of the patients treated with a modified translabyrinthine approach presented evidence of recurrence during follow-up.

The endoscopic surgical technique is a recent source of debate due to its increased use in association with

microsurgery. According to Orhan et al,²³ the association with an angled endoscope avoids invasive techniques (es: middle fossa) better visualizing the hidden area of the temporal bone and allowing a more radical removal of the cholesteatoma matrix, while reducing the incidence of disease recurrence.

Moreover, the improved view of anatomic recesses helps inner ear preservation and hearing function. The disadvantages of this technique are its two-dimensional vision and the need to perform one-handed surgery.²⁴

Gao et al,²⁵ using the combined microscopic and endoscopic technique, reported cochlea preservation in 76.9% of patients with supralabyrinthine PBC, followed by 66.7% of infralabyrinthine and 66.7% of infralabyrinthine-apical ones.

In our series, the endoscope was mainly used in revision surgery and in cases of more extensive PBCs, providing a better view of the potential dehiscence of the facial nerve, dura mater, and labyrinthine fistula.

The main limitation of this paper is the small sample reported, due to the rarity of the disease. Therefore, further multicenter studies with larger samples are under way to confirm these results.

Conclusions

PBC is a rare disease that requires surgical treatment. Its management is based on radical excision evaluating preoperative hearing function, facial function, and the risk of recurrence. Recently, the introduction of modified surgical approaches, able to preserve the anatomical-functional structures, have brought about an improvement in post-operative hearing outcomes.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship and/or publication of this article.

ORCID iDs

Annalisa Pace  <https://orcid.org/0000-0001-6744-5280>

Antonino Maniaci  <https://orcid.org/0000-0002-1251-0185>

Giuseppe Magliulo  <https://orcid.org/0000-0003-3052-1969>

Author Contributions

Giuseppe Magliulo made a substantial contribution to the concept and design of the work; Irene Claudia Visconti, Giannicola Iannella, and Daniela Messineo: acquisition, analysis, and interpretation; Annalisa Pace and Alessandro Milani: acquisition, analysis, interpretation of data, and draft the article; Antonino Maniaci and Valeria Rossetti revised it critically for important intellectual content and approved the version to be published; Salvatore Cocuzza and Giuseppe Magliulo approved the version to be published; All authors

have participated sufficiently in the work and take public responsibility for appropriate portions of the content.

References

- Reddy P, Yan F, Liu YF, McRackan T, Rizk H. Hearing preservation in patients who undergo labyrinthectomy and translabyrinthine procedures a case report and systematic review. *JAMA Otolaryngol Head Neck Surg.* 2020;146(8):741-747. doi: [10.1001/jamaoto.2020.1292](https://doi.org/10.1001/jamaoto.2020.1292).
- er M, Zini C, Gamoletti R, et al. Petrous bone cholesteatoma. *Skull Base Surg.* 1993;3(4):201-213. doi: [10.1055/s-2008-1060585](https://doi.org/10.1055/s-2008-1060585).
- Magliulo G. Petrous bone cholesteatoma: clinical longitudinal study, *Eur Arch Oto-Rhino-Laryngol.* 2007;264(2):115-120. doi: [10.1007/s00405-006-0168-x](https://doi.org/10.1007/s00405-006-0168-x).
- Milroy CM, Michaels L. Pathology of the otic capsule. *J Laryngol Otol.* 1990;104(2):83-90. doi: [10.1017/s0022215100111946](https://doi.org/10.1017/s0022215100111946).
- Sanna M, Pandya Y, Mancini F, Sequino G, Piccirillo E. Petrous bone cholesteatoma: classification, management and review of the literature. *Audiol Neuro Otol.* 2011;16(2):124-136. doi: [10.1159/000315900](https://doi.org/10.1159/000315900).
- Prasad SC, Piras G, Piccirillo E, et al. Surgical strategy and facial nerve outcomes in petrous bone cholesteatoma. *Audiol Neuro Otol.* 2016;21(5):275-285. doi: [10.1159/000448584](https://doi.org/10.1159/000448584).
- Al Zaabi k, Hassannia F, Berrgin MJ, Rutka JA. Management of invasive intralabyrinthine cholesteatoma: can one realistically preserve hearing when disease is medial to the otic capsule? *Am J Otolaryngol.* 2020;41(3):102407. doi: [10.1016/j.amjoto.2020.102407](https://doi.org/10.1016/j.amjoto.2020.102407).
- Fisch U, Pillsbury HC. Infratemporal fossa approach to lesions in the temporal bone and base of the skull. *Arch Otolaryngol.* Feb;105(2):99-107. doi: [10.1001/archotol.1979.00790140045008](https://doi.org/10.1001/archotol.1979.00790140045008).
- Bumsted RM, Dolan KD, Sade J, McCabe BF. Preservation of cochlear function after extensive labyrinthine destruction. *Ann Otol Rhinol Laryngol.* 1977;86(2 pt. 1):131-137. doi: [10.1177/000348947708600201](https://doi.org/10.1177/000348947708600201).
- Hong SJ, Lee JH, Jung SH, Park CH, Hong SM. Can cochlear function be preserved after a modified translabyrinthine approach to eradicate a huge cholesteatoma extending to the petrous apex? *Eur Arch Oto-Rhino-Laryngol.* 2009;266(8):1191-1197. doi: [10.1007/s00405-008-0874-7](https://doi.org/10.1007/s00405-008-0874-7).
- Jahrsdoerfer RA, Johns ME, Cantrell RW. Labyrinthine trauma during ear surgery. *Laryngoscope.* 1978;88(10):1589-1595. doi: [10.1288/00005537-197810000-00004](https://doi.org/10.1288/00005537-197810000-00004).
- Palva T, Johnsson LG. Preservation of hearing after removal of the membranous canal with a cholesteatoma. *Arch Otolaryngol Head Neck Surg.* 1986;112(9):982-985. doi: [10.1001/archotol.1986.03780090078015](https://doi.org/10.1001/archotol.1986.03780090078015).
- Phelps PD. Preservation of hearing in the labyrinth invaded by cholesteatoma. *J Laryngol Otol.* 1969;83(11):1111-1114. doi: [10.1017/s0022215100071401](https://doi.org/10.1017/s0022215100071401).
- McElveen JR, Wilkins RH, Erwin AC, Wolford RD. Modifying the translabyrinthine approach to preserve hearing during acoustic tumour surgery. *J Laryngol Otol.* 1991;105(1):34-37. doi: [10.1017/s0022215100114768](https://doi.org/10.1017/s0022215100114768).
- Magliulo G, Parrotto D, Stasolla A, Marini M. Modified translabyrinthine approach and hearing preservation. *Laryngoscope.* 2004;114(6):1133-1138. doi: [10.1097/00005537-200406000-00034](https://doi.org/10.1097/00005537-200406000-00034).
- Gardner G, Robertson JH. Hearing preservation in unilateral acoustic neuroma surgery. *Ann Otol Rhinol Laryngol.* 1988; 97(1):55-66. DOI: [10.1177/000348948809700110](https://doi.org/10.1177/000348948809700110).
- House JW, Brackmann DE. Facial nerve grading system. *Otolaryngol Head Neck Surg.* 1985;93(2):146-147. doi: [10.1177/019459988509300202](https://doi.org/10.1177/019459988509300202).
- Magliulo G, Celebrini A, Cuiuli G, Parrotto D, Balsamo G, Giuzio. Partial labyrinthectomy in the treatment of labyrinthine fistula: how we do it. *Clin Otolaryngol.* 2008;33(6):607-610. doi: [10.1111/j.1749-4486.2008.01820](https://doi.org/10.1111/j.1749-4486.2008.01820).
- Mohan S, Hoeffner E, Bigelow DC, Loevner LA. Applications of magnetic resonance imaging in adult temporal bone disorders. *Magn Reson Imaging Clin N Am.* 2012;20(3):545-572. doi: [10.1016/j.mric.2012.06.001](https://doi.org/10.1016/j.mric.2012.06.001).
- Song J, An Y, Ahn S, et al. Surgical management toptions and postoperative functional outcomes of petrous apex cholesteatoma. *Acta Otolaryngol.* 2011;131(11):1142-1149. doi: [10.3109/00016489.2011.593550](https://doi.org/10.3109/00016489.2011.593550).
- Moffat D, Jones S, Smith W. Petrous temporal bone cholesteatoma: a new classificator and long-term surgical outcomes. *Skull Base.* 2008;18(2):107-115. doi: [10.1055/s-2007-991112](https://doi.org/10.1055/s-2007-991112).
- Senn P, Haeusler R, Panosetti E, Caversaccio M. Petrous bone cholesteatoma removal with hearing preservation. *Otol Neurotol.* 2011;32(2):236-241. doi: [10.1097/MAO.0b013e3182001ef4](https://doi.org/10.1097/MAO.0b013e3182001ef4).
- Orhan KS, Celik M, Polat B, et al. Endoscope-assisted surgery for petrous bone cholesteatoma with hearing preservation. *J Int Adv Otol.* 2019;15(3):391-395. doi: [10.5152/iao.2019.7212](https://doi.org/10.5152/iao.2019.7212).
- Magliulo G, Iannella G. Endoscopic versus microscopic approach in attic cholesteatoma surgery. *Am J Otolaryngol.* 2018; 39(1):25-30. doi: [10.1016/j.amjoto.2017.10.003](https://doi.org/10.1016/j.amjoto.2017.10.003).
- Gao Z, Gao G, Zhao W, et al. Petrous bone cholesteatoma: our experience of 51 patients with emphasis on cochlea preservation and use of endoscope. *Acta Otolaryngol.* 2019 Jul;139(7): 576-580. doi: [10.1080/00016489.2019.1605455](https://doi.org/10.1080/00016489.2019.1605455).