

OPEN ACCESS

$^8\text{Li} + \alpha$ resonant elastic scattering: a tool to study cluster states in ^{12}B

To cite this article: D Torresi *et al* 2014 *J. Phys.: Conf. Ser.* **569** 012024

View the [article online](#) for updates and enhancements.

You may also like

- [Alpha structure of \$^{12}\text{B}\$ studied by elastic scattering of \$^8\text{Li}\$ EXCYT beam on \$^4\text{He}\$ thick target](#)
M G Pellegriti, D Torresi, L Cosentino et al.
- [The Inverse Kinematics Thick Target scattering method as a tool to study cluster states in exotic nuclei](#)
A Di Pietro, D Torresi, M Zadro et al.
- [Quasi-elastic Scattering of a Secondary \$^6\text{He}\$ Beam on a \$^9\text{Be}\$ Target at 25 MeV/Nucleon](#)
Chen Tao, Ye Yan-Lin, Li Zhi-Huan et al.

$^8\text{Li} + \alpha$ resonant elastic scattering: a tool to study cluster states in ^{12}B

D Torresi^{1,2}, L Cosentino¹, P Descouvemont³, A Di Pietro¹, C Ducoin⁴, P Figuera¹, M Fisichella^{1,5}, M Lattuada^{1,2}, C Maiolino¹, A Musumarra^{1,2}, M Papa⁴, M G Pellegriti^{1,4}, M Rovituso², D Santonocito¹, G Scalia², V Scuderi¹, E Strano^{1,2} and M Zadro⁶

¹INFN Laboratori Nazionali del Sud, Catania, Italy

²Dipartimento di Fisica ed Astronomia, Università di Catania, Catania, Italy

³Physique Nucleaire Theorique et Physique Mathematique, Universite Libre de Bruxelles, Brussels, Belgium

⁴INFN Sezione di Catania, Catania, Italy

⁵Dipartimento di Fisica, Università di Messina, Messina, Italy

⁶Ruder Bošković Institute, Zagreb, Croatia

E-mail: domenico.torresi@lns.infn.it

Abstract. The $^8\text{Li} + ^4\text{He}$ elastic scattering excitation function was measured by using the inverse kinematic thick target method. The ^8Li beam was provided by the LNS radioactive facility EXCYT at an energy of 30 MeV and it was delivered into a large scattering chamber filled with ^4He gas. The detection system was made by three silicon telescopes and one MCP. This last detector was used to measure the number of incident particles as well as the time of flight allowing for the discrimination between elastic and inelastic scattering. The setup and the experimental technique will be described in details and the results, as well as the comparison of the data with an R-matrix calculation, will be shown and discussed.

1. Introduction

One of the interesting topics that can be explored by radioactive beams is the exotic clustering. In this type of clustering nuclei are described as formed by two or more clusters where, at least one of them, is an exotic nucleus itself.

Studying the structure of neutron rich nuclei, Kanada En'yo and collaborators [1] performed calculations in the Antisymmetrized Molecular Dynamics (AMD) framework in order to describe the structure of odd Boron isotopes. Calculations show how the cluster structure becomes more evident and more exotic with the increasing of neutron number in the case of boron isotopes. For instance, the structure of the ^{19}B ground state can be described with a very exotic di-nuclear configuration $^{11}\text{Li} - ^8\text{He}$, while the less neutron rich ^{13}B shows a single particle behavior. Analogous predictions concern also the even isotope ^{12}B . A calculation using the Generator Coordinate Method (GCM) has been performed by Descouvemont [2]. In this study the presence of a positive parity band with a large reduced width for α -decay, located just above the α -decay threshold, supports the picture of a α - ^8Li cluster configuration for ^{12}B .

In this paper it will be described an experiment aimed to the study of the resonant elastic scattering of a ^8Li beam on a ^4He target by using the Thick Target Inverse Kinematic elastic



scattering method (TTIK) [3, 4, 5].

2. The technique

The TTIK method has been used in order to investigate the α - ^8Li structure of ^{12}B in the excitation energy range between 13.5 MeV and 19.6 MeV. In this technique the thick ^4He gas target has a threefold role. It acts at the same time as a degrader, as a target and as a shield for the detector that can be placed along the beam direction. Passing through the ^4He gas, the beam decreases its energy down to zero, hence the elastic scattering is induced in a continuous range of energies between the initial beam energy (30.6 MeV) and zero. While the heavier ^8Li beam is stopped inside the target the recoiling α particles, owing to their lower stopping power, reach the detectors placed at forward angles (backward angles in center of mass (CM) system).




Figure 1. E versus ΔE plot for the telescope at 0° showing the separation of α -particles from hydrogen isotopes.

The ^8Li beam delivered at an energy of 30.6 MeV was produced by the radioactive beam facility EXCYT of INFN-LNS [6]. EXCYT uses the ISOL technique taking advantage of a K-800 superconducting cyclotron as a driver, and of a 13 MV Tandem for post-acceleration of the radioactive beams. Before entering the scattering chamber, the beam passes-through the aluminized Mylar foil ($0.8\ \mu\text{m}$ thick) of a Micro Channel Plate (MCP) detector. This last provides a reference signal for time measurements and, at the same time, allows for counting the beam particles ($\sim 5 \cdot 10^4$ pps). After the MCP detector, the ^8Li beam passes through a kapton foil ($13\ \mu\text{m}$ thick) that separates the beam line from the CT2000 scattering chamber filled with ^4He enriched gas at a pressure $P \sim 700$ mbar and temperature $T \sim 294$ K. The recoiling α particles are detected by an array of three Silicon ΔE -E telescope detectors that are placed at forward angles $\theta = -5^\circ, 0^\circ, +5^\circ$ with respect to the center of the chamber, ~ 180 cm far from the entrance of the chamber. The first stage of the telescope is made by four-quadrant-Silicon-Pad detectors (50 - $60\ \mu\text{m}$ thick) while the second stage is made by Double-Sided-Silicon-Strip-Detectors (DSSSD) 500 - $1000\ \mu\text{m}$ thick. An experimental limitation of the TTIK method is represented by the

contribution of other possible reaction mechanisms to the detected α spectrum. For example inelastic scattering if the center-of-mass energy is high enough to open the inelastic channel. Two α -particles can be detected with the same energy if one comes from an elastic event occurred in a given position of the chamber whilst the second one comes from an inelastic event occurred in a different position of the chamber. Thus, it is not possible to disentangle α coming from elastic and inelastic scattering, looking just at the detected energy of the α -particle.

This difficulty was overcome with by improving the technique with the time of flight measurement. The time elapsed between the ^8Li passing through the foil of the MCP and the detection of α particles was measured. In this case α -particles detected with the same energy but originated from different processes have different time of flight (ToF). Thus, in this way it is possible to discriminate elastic from inelastic events.

3. Analysis

In the following the results obtained from the most forward telescope $\Delta\text{E-E}$, placed at $\theta_{lab}=0^\circ$ with respect to the beam direction will be described. For this telescope, the ΔE nominal thickness is $47\text{ }\mu\text{m}$. The $\Delta\text{E-E}$ plot is shown in Fig. 1. In this plot it is possible to notice that the α -particles are well separated from hydrogen isotopes. We can also see the presence of a large yield of tritons, coming most probably from the break-up channel $^8\text{Li} \rightarrow ^4\text{He} + t + n + 4.5\text{ MeV}$.




Figure 2. ToF versus ΔE plot for the detector at 0° . Continuous line is a calculation for the elastically scattered α -particles. Dotted lines correspond from bottom to the top respectively to the inelastic scattering for the three first excited states of ^8Li .

In Fig. 2, the ToF is shown as a function of the energy deposited in the ΔE detector. The lines are calculations of the ToF as a function of the detected energy both for the elastic and inelastic scattering corresponding to ^8Li first three excited levels. The calculation includes the kinematics, energy losses of ^8Li and α -particles in ^4He gas and energy losses of α -particles in the entrance dead layer of the detector. It can be seen that the calculated locus of the elastic

events is in agreement with the data. As regards the inelastic events, we can notice that the inelastic scattering corresponding to the first excited state of ^8Li is scarcely populated. As one can observe, the elastic events are clearly distinguished from inelastic events.

A source of background comes from the ^8Li decay in two α ($^8\text{Li}(\beta^-) \rightarrow ^8\text{Be}^* \rightarrow 2\alpha$). The life time of ^8Li is 838 msec, for this reason the α particles are uncorrelated in time and hits the detectors at random time. The background of uncorrelated in time α -particles that ranges up to 5 MeV was taken into account and subtracted from the elastically scattered α particles.

For the events which stop in the ΔE detector the elastically scattered α -particles have been selected by a graphical cut on the plot ToF- ΔE . While for particle stopping in the E stage of the telescope it is possible to make a selection of the α particle locus from the ΔE -E plot and then select the elastically scattered α -particles from the ToF-E plot.




Figure 3. Elastic scattering differential cross section for most forward angles ($\sim 180^\circ$).

In order to reconstruct the elastic scattering excitation function from the spectrum of the detected α -particles it is required a precise knowledge of the stopping power of ^4He for ^8Li and α . For this reason during the same experiment a measurement of the stopping power was also performed.

The elastic scattering excitation function extracted for the telescope placed at 0° degree is shown in Fig. 3 and it manifests different structures. The analysis of the excitation functions was performed using an R-matrix approach. In order to simplify the problem the approximation of spin zero for ^8Li was considered. In this case, the spin of each level J is equal to the relative angular momentum l . In order to fit the data seven resonances were used. An additional background resonance at higher energy was introduced to fit the long tail in the high energy region. For the data fitting, the R-matrix calculation was convoluted with the experimental resolution function. The $^8\text{Li}+\alpha$ excitation function is reproduced rather well by the R-matrix calculations as shown in Fig. 4. Some of the resonances was previously known [7, 8].

The reduced α -widths of the seven resonances used to perform the fit were calculate and their low values (less than 5%) indicate that there is no clear evidence of α -cluster configurations associated to the observed resonances.




Figure 4. The elastic scattering cross section experimental data are shown as full circles, the solid red line is the R-matrix fit.

4. Conclusions

We have reported the measurements of the $^8\text{Li}-\alpha$ elastic scattering excitation function by using the TTIK method. The technique included a time of flight measurement that allows the discrimination of elastic scattering from other reaction mechanisms. The excitation function was measured for excitation energy in the range of 13.5 and 19.5 MeV and it was analyzed using an R-matrix approach. The R-matrix calculation, that reproduces rather well the experimental data, includes seven resonances plus a background term. The reduced α -widths do not show evidence of strong α cluster configurations

References

- [1] Y. Kanada En'yo et al., *Prog. Theor. Phys.* **93** (1995) 115.
- [2] P. Descouvemont, *Nucl. Phys. A* **596** (1996) 285.
- [3] K.P. Artemov et al., *Sov. J. Nucl. Phys.* **52** (1990) 408.
- [4] K.M. Kallman et al., *Nucl. Instr. and Meth. in Phys. Res. A* **338** (1994) 413.
- [5] V.Z. Goldberg et al., *Phys. Rev. C* **69** (2004) 024602.
- [6] G. Ciavola et al., *Nucl. Phys. A* **701** (2002) 54c.
- [7] F. Ajzenberg-Selove et al., *Nucl. Phys. A* **506** (1990) 1.
- [8] N. Soić et al., *Nucl. Phys. A* **738** (2004) 347c.