



FIGURE 10-1 Total Numbers and Rates of Non-rheumatic Degenerative Mitral Valve Disease: Global



Global non-rheumatic degenerative mitral valve disease count in millions and rate per 100,000 by measure for age-standardized and all-age groups with shaded 95% CI from 1990 to 2023. DALY = disability-adjusted life year.

FIGURE 10-2 Age-Standardized DALY Rates for Non-rheumatic Degenerative Mitral Valve Disease, 2023

Age-standardized non-rheumatic degenerative mitral valve disease disability-adjusted life years (DALYs) per 100,000 in 2023 (all sexes combined).

Russian Federation (2.1% [95% UI: 1.5% to 2.8%]), while locations with large decreases included Belgium (3.5% [95% UI: 2.9% to 4.1%]), the Netherlands (3.4% [95% UI: 3.0% to 3.9%]), and Singapore (2.9% [95% UI: 2.3% to 3.6%]).


CARDIOMYOPATHY AND MYOCARDITIS. Cardiomyopathy and myocarditis represented the aggregate of myocarditis, alcoholic myocarditis, and other cardiomyopathy. In 2023, DALYs from cardiomyopathy and myocarditis were mainly composed of other cardiomyopathy (76.5% [95% UI: 72.6% to 80.1%]), followed by alcoholic cardiomyopathy (18.1% [95% UI: 14.9% to 21.8%]), and last by myocarditis (5.4% [95% UI: 3.7% to 8.0%]). In 2023, cardiomyopathy and myocarditis caused 12.0 million (95% UI: 9.80 to 14.4 million) DALYs, 400,000 (95% UI: 338,000 to 465,000) deaths, and 5.35 million (95% UI: 4.39 to 6.23 million) prevalent cases (Figure 11-1). The age-standardized DALY rate for males was 186.8 (95% UI: 148.9 to 240.9) per 100,000, and for females was 97.5 (95% UI: 68.6 to 134.9) per 100,000 (Figure 11-2).

By world region, the greatest age-standardized rate of cardiomyopathy and myocarditis DALYs was in Eastern Europe (846.6 [95% UI: 768.4 to 928.8] per 100,000) and the lowest was in East Asia (29.0 [95% UI: 21.8 to 37.5]) per 100,000 (Figure 11-3). From 1990 to 2023, the total number of DALYs increased steadily by 32.8% (95% UI: 2.4% to 71.0%) (Figure 11-1); however, age-standardized DALY rates decreased by 30.4% (95% UI: 11.2% to 46.0%).


In 2023, the remainder category of other cardiomyopathy accounted for 9.19 million (95% UI: 7.12 to 11.3 million) DALYs, 321,000 (95% UI: 260,000 to 381,000) deaths, and 4.42 million (95% UI: 3.51 to 5.28) prevalent cases. The age-standardized DALY rate ranged from 16.5 DALYs (95% UI: 11.6 to 22.6 DALYs) per 100,000 in East Asia to 367.3 DALYs (95% UI: 171.9 to 661.4 DALYs) per 100,000 in Central Sub-Saharan Africa. The number of DALYs increased by 53.1% (95% UI: 8.2% to 105.8%) from 1990 to 2023.

MYOCARDITIS. In 2023, age-standardized DALYs per 100,000 population from myocarditis ranked 17th

FIGURE 11-1 Total Numbers and Rates of Cardiomyopathy and Myocarditis: Global



Global cardiomyopathy and myocarditis count in millions and rate per 100,000 by measure for age-standardized and all-age groups with shaded 95% CI from 1990 to 2023. DALY = disability-adjusted life year.

FIGURE 11-2 Global Cardiomyopathy and Myocarditis DALY Rate by Age and Sex, for 2023

Global age-specific disability-adjusted life year (DALY) rate of cardiomyopathy and myocarditis in 2023 for males and females. Specific age groups under the age of 5 years were aggregated to "under 5" for clarity.


among most detailed CVD causes, were higher in males vs females, and increased with age (Figure 12-1). From 1990 to 2023, the number of incident and prevalent cases of myocarditis increased over time, while the number of deaths and DALYs decreased. During that time, the age-standardized DALYs and deaths decreased, while incidence and prevalence flattened. In the GBD study's comparative risk assessment framework, myocarditis was only attributed to one risk factor: non-optimal temperature.

In 2023, myocarditis led to 650,000 (95% UI: 440,000 to 969,000) DALYs, 16,900 (95% UI: 11,300 to 24,100) deaths, and 390,000 (95% UI: 321,000 to 467,000) prevalent cases. Males had similar age-standardized rates of DALYs (10.1 [95% UI: 6.1 to 17.2] per 100,000) compared with females (6.7 [95%

UI: 3.4 to 13.7] per 100,000) (Figure 12-1). By world region, the greatest DALY rate was in Central Sub-Saharan Africa (22.2 [95% UI: 6.4 to 56.3] per 100,000), and the lowest in Andean Latin America (1.8 [95% UI: 1.3 to 2.6] per 100,000) (Supplemental Figure 11).

From 1990 to 2023, the total number of myocarditis DALYs decreased steadily by 46.1% (95% UI: 16.9% to 66.5%) (Supplemental Figure 12). The age-standardized DALY rate also decreased by 62.5% (95% UI: 43.0% to 76.7%). In 2023, 43,500 (95% UI: 15,600 to 85,700) DALYs from myocarditis were attributable to non-optimal temperature. These decreased by 46,500 (15,900 to 92,200) DALYs from 1990, corresponding to a reduction of 51.6% (95% UI: 23.1% to 71.7%). This decrease was mainly driven by a

FIGURE 11-3 Age-Standardized DALY Rates for Cardiomyopathy and Myocarditis, 2023




Age-standardized cardiomyopathy and myocarditis disability-adjusted life years (DALYs) per 100,000 in 2023 (all sexes combined).

reduction of 64,000 (95% UI: 21,700 to 127,000) DALYs due to risk-deleted DALY rate, followed by a decrease of 20,400 (95% UI: 7,560 to 40,400) DALYs due to population aging (Figure 12-2). The change in non-optimal temperature risk exposure globally contributed to little to no change in DALYs, changing by only -7,970 (95% UI: -20,600 to 9,580).

ALCOHOLIC CARDIOMYOPATHY. In 2023, age-standardized DALYs per 100,000 population from alcoholic cardiomyopathy ranked 13th among most detailed CVD causes. DALY rates were higher in males vs females, and by age they showed a bimodal distribution, peaking at 60 to 64 years of age and increasing again after 80 years of age. The total number and age-standardized DALYs and deaths increased between 1990 and 2005 before declining through 2023 (Figure 13-1). For prevalence, the number of cases increased over time, while the age-standardized rates decreased. Alcoholic cardiomyopathy DALYs were attributable to high alcohol use

and non-optimal temperature, with population growth and aging as the factors driving these increases.

Alcoholic cardiomyopathy was responsible for 2.16 million (95% UI: 1.93 to 2.46 million) DALYs, 62,300 (95% UI: 56,000 to 71,600) deaths, and 544,000 (95% UI: 453,000 to 647,000) prevalent cases in 2023 (Figure 13-1). Among males, the age-standardized DALY rate per 100,000 population was 39.8 (95% UI: 34.6 to 46.5), and among females it was 9.0 (95% UI: 7.7 to 10.8) (Supplemental Figure 13). By age, the greatest DALY rates were among adults 60 to 64 years of age, with 124.4 (95% UI: 111.0 to 141.9) per 100,000 for males and 25.1 (95% UI: 21.4 to 30.2) per 100,000 for females. By world region, the greatest age-standardized DALY rate was in Eastern Europe (470.6 [95% UI: 419.9 to 532.2] per 100,000) and the lowest was in Andean Latin America (0.1 [95% UI: 0.1 to 0.2] per 100,000). From 1990 to 2023, the total number of DALYs from alcoholic cardiomyopathy increased by 18.1% (95% UI: 1.6% to 38.9%), while the

FIGURE 12-1 Global Myocarditis DALY Rate by Age and Sex, for 2023

Global age-specific disability-adjusted life year (DALY) rate of myocarditis in 2023 for males and females. Specific age groups under the age of 5 years were aggregated to “under 5” for clarity.

age-standardized DALYs decreased by 41.8% (95% UI: 31.8% to 49.9%) (Figure 13-1).


From 1990 to 2023, alcoholic cardiomyopathy attributable to high alcohol use increased by 330,000 (95% UI: 30,300 to 637,000) DALYs. This increase was driven by population growth (932,000 [95% UI: 792,000 to 1,100,000] DALYs) and population aging (559,000 [95% UI: 495,000 to 651,000] DALYs) (Figure 13-2). High alcohol consumption accounts for 100% of the risk for alcoholic cardiomyopathy; however, non-optimal temperature increases the risk of alcoholic cardiomyopathy when high alcohol is present, hence this is also shown in Figure 13-2.

PULMONARY ARTERIAL HYPERTENSION. Pulmonary arterial hypertension was a less common condition among the CVD estimated in GBD, with the

16th largest number of DALYs among CVDs globally in 2023. Pulmonary arterial hypertension had a “U-shaped” age-distribution of DALYs: DALY rates globally were high in groups younger than 5 years of age, decreased near 10 through 20 years of age, then increased steadily beyond 25 years of age (Figure 14-1). Age-standardized pulmonary arterial hypertension DALY rates have decreased slightly since 1990, while the all-age DALY count was consistently level. Pulmonary arterial hypertension DALYs were not attributable to any risk measured in GBD 2023.

There were 701,000 (95% UI: 515,000 to 999,000) pulmonary arterial hypertension DALYs globally in 2023. In 1990, the age-standardized pulmonary arterial hypertension DALY rate was 17.2 (95% UI: 12.1 to 23.1) per 100,000, and since then the DALY rate has

FIGURE 12-2 Percentage Change in the Number of Global Risk-Attributable DALYs, 1990 to 2023, due to Population Growth, Population Aging, Changes in Exposures to Each Global Burden of Disease Risk Factor, and Changes in Risk-Deleted DALY Rates for All Sexes, for Myocarditis



Decomposition of change in all-age, all sexes combined myocarditis disability-adjusted life years (DALYs) attributable to risk factors from 1990 to 2023 due to population growth, population aging, risk exposure, and risk-deleted DALYs. Risk-deleted DALYs are the number of DALYs left after removing the effect of risk factors, population growth, and population aging on overall DALYs. They were calculated as the overall myocarditis DALY count multiplied by 1 minus the population attributable fraction for each risk. The dot and error bar represent the mean and 95% uncertainty interval in percentage change in number of DALYs attributable to the risk from 1990 to 2023.

decreased globally, with an annualized rate of change decrease of 2.1% (95% UI: 0.8% to 3.5%) (Figure 14-2).


DALY rates were similar for females than males globally in 2023: the age-standardized rate was 9.0 (95% UI: 5.6 to 13.6) per 100,000 for females and 8.4 (95% UI: 5.3 to 13.1) per 100,000 for males (Supplemental Table 3). This sex pattern held for both incidence and deaths as well; there were 20,800 (95% UI: 16,400 to 25,200) incident cases and 9,900 (95% UI: 6,990 to 13,700) pulmonary arterial hypertension deaths for males, while females had 24,500 (95% UI: 19,600 to 29,700) incident cases and 12,900 (95% UI: 9,200 to 18,400) pulmonary arterial hypertension deaths in 2023 globally.

ATRIAL FIBRILLATION AND FLUTTER. Among CVDs estimated in GBD, atrial fibrillation and flutter had

the eighth-highest number of DALYs. Atrial fibrillation and flutter age-standardized DALY rates varied between 45.6 and 238.4 per 100,000 globally, a more than 5-fold difference between countries. The age-standardized rates of atrial fibrillation and flutter DALYs, deaths, and prevalence all showed little change from 1990 to 2023. However, the all-age rate and number of DALYs, deaths, and prevalence have all increased globally since 1990. The increase in all-age burden, reflected by DALYs, was primarily due to population aging, while changes in exposure to harmful risks and risk-deleted burden did little to mitigate population changes.


As of 2023, there were 9.27 million (95% UI: 7.52 to 11.7 million) atrial fibrillation and flutter DALYs globally (Figure 15-1). Age-standardized atrial fibrillation and flutter burden was larger for males than

FIGURE 13-1 Total Numbers and Rates of Alcoholic Cardiomyopathy: Global



Global alcoholic cardiomyopathy count in millions and rate per 100,000 by measure for age-standardized and all-age groups with shaded 95% CI from 1990 to 2023. DALY = disability-adjusted life year.

FIGURE 13-2 Percentage Change in the Number of Global Risk-Attributable DALYs, 1990 to 2023, due to Population Growth, Population Aging, Changes in Exposures to Each Global Burden of Disease Risk Factor, and Changes in Risk-Deleted DALY Rates for All Sexes, for Alcoholic Cardiomyopathy




Decomposition of change in all-age, all sexes combined alcoholic cardiomyopathy disability-adjusted life years (DALYs) attributable to risk factors from 1990 to 2023 due to population growth, population aging, risk exposure, and risk-deleted DALYs. Risk-deleted DALYs are the number of DALYs left after removing the effect of risk factors, population growth, and population aging on overall DALYs. They were calculated as the overall alcoholic cardiomyopathy DALY count multiplied by 1 minus the population attributable fraction for each risk. The dot and error bar represent the mean and 95% uncertainty interval in percentage change in number of DALYs attributable to the risk from 1990 to 2023.

females: the age-standardized DALY rate for males was 114.5 (95% UI: 91.9 to 145.3) per 100,000 and for females was 94.4 (95% UI: 76.9 to 117.1) per 100,000 globally in 2023 (Supplemental Table 3). The GBD region with the highest DALY rate in 2023 was high-income North America at 159.9 (95% UI: 133.2 to 192.8) per 100,000, and the lowest was high-income Asia Pacific at 70.7 (95% UI: 56.5 to 88.9) per 100,000, only a 2-fold difference (Supplemental Figure 14). Atrial fibrillation and flutter was a common cause of cardiovascular deaths: there were 378,000 (95% UI: 319,000 to 424,000) deaths globally in 2023. There were 59.0 million (95% UI: 46.5 to 72.8 million) prevalent cases of atrial fibrillation and flutter globally in 2023. The prevalence rate of atrial fibrillation and flutter was slightly higher in males

than females globally; for males, the age-standardized prevalence rate was 776.3 (95% UI: 615.3 to 949.3) per 100,000 and for females, 541.9 (95% UI: 421.1 to 671.1) per 100,000.

In 2023, 3.64 million (95% UI: 1.98 to 5.62 million) atrial fibrillation and flutter DALYs were attributable to a modifiable risk factor, 39.2% (95% UI: 21.7% to 55.3%) of atrial fibrillation and flutter DALYs (Table 4). The leading risk factors for atrial fibrillation and flutter were high SBP, high BMI, and tobacco use, which accounted for 2.81 million (95% UI: 1.03 to 4.83 million), 759,000 (95% UI: 374,000 to 1.23 million), and 414,000 (95% UI: 225,000 to 603,000) atrial fibrillation DALYs, respectively. High alcohol use and lead exposure accounted for the least atrial fibrillation and flutter DALYs; high alcohol use accounted

FIGURE 14-1 Global Pulmonary Arterial Hypertension DALY Rate by Age and Sex, for 2023

Global age-specific disability-adjusted life year (DALY) rate of pulmonary arterial hypertension in 2023 for males and females. Specific age groups under the age of 5 years were aggregated to "under 5" for clarity.


for 280,000 DALYs (95% UI: 132,000 to 467,000 DALYs), while lead exposure accounted for 186,000 DALYs (95% UI: -23,600 to 470,000 DALYs).

The burden of atrial fibrillation attributable to risk factors has grown by 2.26 million (95% UI: 1.17 to 3.56 million) DALYs since 1990, a 163.8% (95% UI: 127.2% to 210.4%) increase globally. Population aging has contributed 1.55 million (95% UI: 0.825 to 2.37 million) atrial fibrillation and flutter DALYs since 1990, while population growth added 0.705 million (95% UI: 0.390 to 1.06 million) atrial fibrillation and flutter DALYs (Figure 15-2). The increase in DALYs from population growth and aging was not offset by changing exposure to harmful risk factors or risk-deleted burden. Increased exposure to high BMI and lead exposure contributed most among risk factors to the increase in atrial fibrillation DALYs. The change in exposure to high BMI led to an additional 237,000 (95% UI: 80,800 to 439,000) atrial

fibrillation and flutter DALYs since 1990. Changes in lead exposure resulted in an additional 21,300 (95% UI: -2,530 to 59,100) atrial fibrillation and flutter DALYs globally, though the slight overlap of the UI with 0 suggests the possibility of no change globally. The larger counterbalance to the increases in DALYs was from reduced exposure to tobacco. The change in exposure to tobacco usage mitigated 101,000 (95% UI: 39,300 to 188,000) atrial fibrillation and flutter DALYs.


AORTIC ANEURYSM. Aortic aneurysm was only evaluated as a cause of death, and non-fatal outcomes were not estimated. In 2023, aortic aneurysm ranked 10th among age-standardized DALY rates for all CVDs. DALY rates from aortic aneurysm were higher in males vs females and increased with age. From 1990 to 2023, the number of DALYs and deaths increased, while the age-standardized rates

FIGURE 14-2 Total Numbers and Rates of Pulmonary Arterial Hypertension: Global




Global pulmonary arterial hypertension count in millions and rate per 100,000 by measure for age-standardized and all-age groups with shaded 95% CI from 1990 to 2023. DALY = disability-adjusted life year.

FIGURE 15-1 Total Numbers and Rates of Atrial Fibrillation and Flutter: Global



Global atrial fibrillation and flutter count in millions and rate per 100,000 by measure for age-standardized and all-age groups with shaded 95% CI from 1990 to 2023. DALY = disability-adjusted life year.

FIGURE 15-2 Percentage Change in the Number of Global Risk-Attributable DALYs, 1990 to 2023, due to Population Growth, Population Aging, Changes in Exposures to Each Global Burden of Disease Risk Factor, and Changes in Risk-Deleted DALY Rates for All Sexes, for Atrial Fibrillation and Flutter



Decomposition of change in all-age, all sexes combined atrial fibrillation and flutter disability-adjusted life years (DALYs) attributable to risk factors from 1990 to 2023 due to population growth, population aging, risk exposure, and risk-deleted DALYs. Risk-deleted DALYs are the number of DALYs left after removing the effect of risk factors, population growth, and population aging on overall DALYs. They were calculated as the overall atrial fibrillation and flutter DALY count multiplied by 1 minus the population attributable fraction for each risk. The dot and error bar represent the mean and 95% uncertainty interval in percentage change in number of DALYs attributable to the risk from 1990 to 2023.


decreased. The increase in DALYs from aortic aneurysm was attributable to metabolic, behavioral, and environmental risk factors, mostly driven by population growth and population aging.

In 2023, there were 3.42 million (95% UI: 3.03 to 3.84 million) DALYs and 167,000 (95% UI: 147,000 to 187,000) deaths from aortic aneurysm (Figure 16-1). Among males, the rate of DALYs was 54.3 (95% UI: 47.7 to 63.0) per 100,000, while for females it was 22.9 (95% UI: 19.3 to 27.2) per 100,000 (Supplemental Figure 15). By world region, the greatest DALY rate was in Eastern Europe (89.0 [95% UI: 80.4 to 99.3] DALYs per 100,000) and lowest in East Asia (12.2 [95% UI: 9.3 to 15.1] DALYs per 100,000). From 1990 to 2023, the number of DALYs from aortic aneurysm increased steadily by 78.0% (95% UI: 55.9% to 101.0%). In contrast, the age-standardized DALYs and

death rates per 100,000 population decreased by 23.2% (95% UI: 13.7% to 32.6%).


In 2023, 1.63 million (95% UI: 1.37 to 1.89 million) DALYs from aortic aneurysm were attributable to measured modifiable risk factors, corresponding to 47.7% (95% UI: 42.5% to 53.8%) of all DALYs from aortic aneurysm. Tobacco exposure (1.24 million [95% UI: 1.02 to 1.48 million] DALYs) and high SBP (609,000 [95% UI: 450,000 to 790,000]) were the top risk factors. From 1990 to 2023, DALYs from aortic aneurysm attributable to all modifiable risk factors increased by 552,000 (95% UI: 328,000 to 762,000), representing an increase of 51.3% (95% UI: 29.5% to 71.4%) from 1990 (Figure 16-2). These increases were mainly due to population aging (636,000 [95% UI: 542,000 to 735,000]) and population growth (549,000 [95% UI: 472,000 to 623,000]). Out of all

FIGURE 16-1 Total Numbers and Rates of Aortic Aneurysm: Global



Global aortic aneurysm count in millions and rate per 100,000 by measure for age-standardized and all-age groups with shaded 95% CI from 1990 to 2023. DALY = disability-adjusted life year.

FIGURE 16-2 Percentage Change in the Number of Global Risk-Attributable DALYs, 1990 to 2023, due to Population Growth, Population Aging, Changes in Exposures to Each Global Burden of Disease Risk Factor, and Changes in Risk-Deleted DALY Rates for All Sexes, for Aortic Aneurysm



Decomposition of change in all-age, all sexes combined aortic aneurysm disability-adjusted life years (DALYs) attributable to risk factors from 1990 to 2023 due to population growth, population aging, risk exposure, and risk-deleted DALYs. Risk-deleted DALYs are the number of DALYs left after removing the effect of risk factors, population growth, and population aging on overall DALYs. They were calculated as the overall aortic aneurysm DALY count multiplied by 1 minus the population attributable fraction for each risk. The dot and error bar represent the mean and 95% uncertainty interval in percentage change in number of DALYs attributable to the risk from 1990 to 2023.

risk factors, lead exposure (120.6% [95% UI: 69.8% to 174.9%]) and high BMI (94.1% [95% UI: 55.2% to 144.4%]) contributed to the largest proportional increases in DALYs from aortic aneurysm.


LOWER EXTREMITY PERIPHERAL ARTERIAL DISEASE.

The global burden of peripheral arterial disease is lower than that of most CVDs, ranked as the 14th-highest cause of DALYs among CVDs. There was notable geographic variation in peripheral arterial diseases globally in 2023. The age-standardized rates of peripheral arterial disease DALYs, deaths, and prevalence all decreased from 1990 to 2023, while all-age rates all increased.

Globally there were 1.86 million (95% UI: 1.44 to 2.47 million) peripheral arterial disease DALYs in 2023. The age-standardized DALY rate was 20.6 (95% UI: 15.9 to 27.3) per 100,000 in 2023, a decrease since

1990 (28.1 [95% UI: 23.1 to 35.7] per 100,000). The burden was highest in the older ages: there were 1.23 million (95% UI: 0.928 to 1.67 million) DALYs in groups 70 years of age and older (Supplemental Figure 16). The age-standardized DALY rates in 2023 varied among GBD regions, with the highest in Eastern Europe, at 64.1 (95% UI: 55.1 to 75.1) per 100,000, and the lowest in East Asia, at 8.0 (95% UI: 4.2 to 14.2) per 100,000, an 8-fold difference (Figure 17-1). Peripheral arterial disease was also a highly prevalent cause of CVDs globally, with 122 million (95% UI: 93.9 to 157 million) prevalent cases in 2023 (47.2 million [95% UI: 36.1 to 60.8 million] in males and 74.8 million [95% UI: 57.6 to 95.8 million] in females).

In 2023, 1.24 million (95% UI: 0.937 to 1.69 million) peripheral arterial disease DALYs were attributable to a modifiable risk factor, accounting for 66.5% (95%

FIGURE 17-1 Age-Standardized DALY Rates for Lower Extremity Peripheral Arterial Disease, 2023

Age-standardized lower extremity peripheral arterial disease disability-adjusted life years (DALYs) per 100,000 in 2023 (all sexes combined).

UI: 58.2% to 74.1%) of DALYs (Table 4). Globally, high FPG, kidney dysfunction, tobacco use, and high BMI were the leading risk factors for peripheral arterial disease, accounting for 481,000 (95% UI: 338,000 to 702,000) DALYs, 580,000 (95% UI: 328,000 to 928,000) DALYs, 454,000 (95% UI: 319,000 to 649,000) DALYs, and 365,000 (95% UI: 98,400 to 765,000) DALYs in 2023, respectively.


The number of peripheral arterial disease DALYs attributable to risk factors has increased by 591,000 (95% UI: 409,000 to 857,000) since 1990. Population growth added 330,000 (95% UI: 257,000 to 428,000) DALYs, while aging added 578,000 (95% UI: 444,000 to 778,000) DALYs (Figure 17-2). Increases in the exposure to high FPG and high BMI led to increases of 62,200 (95% UI: -29,600 to 227,000) DALYs and 101,000 (95% UI: -13,400 to 289,000) DALYs since 1990, respectively. For high FPG and high BMI, the UIs were only slightly above zero, suggesting a potential contribution to the increase in DALYs. Beneficial reductions in tobacco use helped lessen the increases in peripheral arterial disease DALYs; DALYs

decreased by 191,000 (95% UI: 103,000 to 307,000) as a result.

ENDOCARDITIS. Age-standardized DALY rates from endocarditis ranked 11th among all CVD causes. DALY rates increased with age and were higher in males vs females among those younger than 90 years of age, but higher in females compared with males 90+ years of age. The number of DALYs, deaths, prevalence, and incidence increased steadily from 1990 to 2023. The age-standardized rates per 100,000 population increased for prevalence and incidence but not for deaths, in which a decrease was observed.

In 2023, there were 2.34 million (95% UI: 1.96 to 2.87 million) DALYs and 86,200 (95% UI: 74,200 to 101,000) deaths from endocarditis (Figure 18-1). Among males, the age-standardized rate of DALYs was 31.2 (95% UI: 24.3 to 41.1) DALYs per 100,000, while for females it was 24.1 (95% UI: 17.4 to 33.1) DALYs per 100,000. By world region, the greatest age-standardized DALY rate was in Oceania (78.0 [95% UI: 49.4 to 119.5] per 100,000) and the lowest

FIGURE 17-2 Percentage Change in the Number of Global Risk-Attributable DALYs, 1990 to 2023, due to Population Growth, Population Aging, Changes in Exposures to Each Global Burden of Disease Risk Factor, and Changes in Risk-Deleted DALY Rates for All Sexes, for Lower Extremity Peripheral Arterial Disease




Decomposition of change in all-age, all sexes combined lower extremity peripheral arterial disease disability-adjusted life years (DALYs) attributable to risk factors from 1990 to 2023 due to population growth, population aging, risk exposure, and risk-deleted DALYs. Risk-deleted DALYs are the number of DALYs left after removing the effect of risk factors, population growth, and population aging on overall DALYs. They were calculated as the overall lower extremity peripheral arterial disease DALY count multiplied by 1 minus the population attributable fraction for each risk. The dot and error bar represent the mean and 95% uncertainty interval in percentage change in number of DALYs attributable to the risk from 1990 to 2023.

was in East Asia (4.4 [95% UI: 3.4 to 5.6] per 100,000) (Figure 18-2, Supplemental Figure 17). From 1990 to 2023, the number of DALYs from endocarditis increased by 55.3% (95% UI: 8.0% to 111.7%). Endocarditis is not attributed to any risk factors in the GBD study’s comparative risk assessment framework.

CARDIOVASCULAR RISK FACTORS. Summary of global trends in CVD risks. In 2023, 79.6% (95% UI: 75.7% to 82.5%) of CVD DALYs were attributable to modifiable risk factors (Supplemental Table 8). This was led by metabolic risk factors, which contributed to 67.3% (95% UI: 61.7% to 72.9%) of CVD DALYs, followed by behavioral risk factors and environmental/occupational risk factors, which contributed to 44.9% (95% UI: 28.8% to 54.7%) and


35.8% (95% UI: 30.9% to 40.7%) of CVD DALYs, respectively. Since 1990, the percentage of CVD DALYs attributable to metabolic risk factors has increased from 63.1% (95% UI: 54.8% to 70.0%), primarily driven by a notable increase in DALYs attributable to high BMI and more moderate increases in DALYs attributable to high FPG, kidney dysfunction, high SBP, and high LDL-C. The percentage of CVD DALYs attributable to behavioral risk factors and environmental/occupational risk factors decreased from 47.3% (95% UI: 31.3% to 55.7%) and 38.0% (95% UI: 32.4% to 43.5%), respectively, in 1990. For behavioral risk factors, a small increase in the percentage of CVD burden attributable to low physical activity was opposed by decreases in burden attributable to high alcohol use, dietary risks, and, most

FIGURE 18-1 Total Numbers and Rates of Endocarditis: Global



Global endocarditis count in millions and rate per 100,000 by measure for age-standardized and all-age groups with shaded 95% CI from 1990 to 2023. DALY = disability-adjusted life year.

FIGURE 18-2 Age-Standardized DALY Rates for Endocarditis, 2023




Age-standardized endocarditis disability-adjusted life years (DALYs) per 100,000 in 2023 (all sexes combined).

notably, tobacco. Among environmental/occupational risk factors, the attributable percentage for CVD DALYs increased for lead exposure but decreased for air pollution and non-optimal temperature. There was considerable geographic variation in the risk factors contributing to CVD burden. Metabolic risk factors accounted for the highest percentage of CVD DALYs in middle SDI regions, followed by high-middle and high SDI regions, with the lowest percentage of CVD DALYs in low SDI regions. Similarly, behavioral risk factors accounted for the highest percentage of CVD DALYs in middle SDI regions, followed by high-middle SDI regions, with the lowest percentage of CVD DALYs in low SDI regions. Environmental/occupational risk factors accounted for the highest percentage of CVD DALYs in low-middle SDI regions, followed by high-middle and low SDI regions, with the lowest percentage of CVD DALYs in high SDI regions.

High FPG. Burden attributable to high FPG is estimated for 4 CVDs: IHD, ischemic stroke, intracerebral hemorrhage, and lower extremity peripheral arterial disease. In addition to CVDs, high FPG was also a large contributor to diabetes mellitus. High FPG

ranked ninth in terms of contributors to cardiovascular DALYs in both 1990 and 2023. The percentage of cardiovascular DALYs attributed to it rose from 6.6% (95% UI: 4.9% to 9.3%) in 1990 to 8.5% (95% UI: 7.1% to 10.7%) in 2023, in part due to the increase in global exposure to high FPG since 1990 (Supplemental Table 8).

Globally, the number of cardiovascular DALYs attributable to high FPG increased from 21.2 million (95% UI: 15.9 to 30.1 million) in 1990 to 37.3 million (95% UI: 30.3 to 47.4 million) in 2023 (Supplemental Table 7). Global age-standardized cardiovascular DALYs attributable to high FPG decreased from 578.2 (95% UI: 440.1 to 813.9) per 100,000 in 1990 to 410.0 (95% UI: 333.3 to 521.4) per 100,000 in 2023, though not all locations experienced a decrease (Figure 19-1). Western Europe and high-income Asia Pacific had the largest decreases in age-standardized attributable DALY rates with annualized rates of change of -2.9% (95% UI: -3.7% to -2.1%) and -2.9% (95% UI: -4.1% to -1.9%), respectively, between 1990 and 2023. No region showed a notable increase in attributable DALY rates from 1990 to 2023; Eastern Sub-Saharan Africa increased the most, with an annualized rate

FIGURE 19-1 Annualized Rate of Change in Age-Standardized Cardiovascular Disease Disability-Adjusted Life Year Rates Attributable to High Fasting Plasma Glucose 1990-2023

Annualized rate of change in age-standardized cardiovascular disease disability-adjusted life years attributable to high fasting plasma glucose from 1990 to 2023 (all sexes combined).


of change of 0.9% (95% UI: -0.2% to 2.1%), though the UI overlapped with zero. In 2023, age-standardized cardiovascular DALYs attributable to high FPG were highest in North Africa and the Middle East (878.6 [95% UI: 728.3 to 1,083.7] per 100,000) and Central Asia (812.7 [95% UI: 672.4 to 1,025.4] per 100,000) and lowest were in Eastern Sub-Saharan Africa (116.2 [95% UI: 82.5 to 167.2] per 100,000).

Exposure to high FPG increased globally between 1990 and 2023 (Supplemental Table 11). Among females, the age-standardized SEV increased from 13.8 (95% UI: 9.1 to 22.4) to 17.4 (95% UI: 12.7 to 25.2), while that of males increased from 14.9 (95% UI: 9.9 to 23.5) to 17.7 (95% UI: 13.3 to 24.4). This rise in high FPG exposure contributed to an increase of 6.51 million (95% UI: -4.03 to 16.5 million) cardiovascular DALYs attributable to high FPG since 1990 (Figure 19-2; Supplemental Table 8), with the UI slightly crossing zero, suggesting only a potential contribution to the increase in DALYs. Similarly, the effects of population growth and aging contributed to an increase of 10.8 million (95% UI: 8.12 to 15.0 million)

and 16.6 million (95% UI: 13.8 to 20.5 million) attributable cardiovascular DALYs, respectively. These sources of increases in DALYs were countered by changes in risk-deleted DALY rates, which contributed to a decrease of 17.8 million (95% UI: 13.3 to 23.3 million) cardiovascular DALYs attributable to high FPG. Among the 4 CVDs with which it is associated in the GBD study's comparative risk assessment framework, increasing high FPG exposure contributed to the largest percentage increase in attributable DALYs for IHD and lower extremity peripheral arterial disease.

High LDL-C. The burden attributable to high LDL-C was estimated for 2 cardiovascular outcomes: ischemic stroke and IHD. High LDL-C was the fifth- and third-largest contributor to cardiovascular DALYs in 1990 and 2023, respectively, with 20.5% (95% UI: 12.6% to 28.3%) and 20.8% (95% UI: 13.6% to 29.0%) of cardiovascular DALYs attributed to high LDL-C, respectively (Supplemental Table 8). Despite global exposure remaining fairly consistent since 1990, many locations experienced large changes,

FIGURE 19-2 Percentage Change in the Number of Global Risk-Attributable DALYs, 1990 to 2023, due to Population Growth, Population Aging, Changes in Exposures to Each Global Burden of Disease Risk Factor, and Changes in Risk-Deleted DALY Rates for All Sexes, for High Fasting Plasma Glucose




Decomposition of change in all-age, all sexes combined cardiovascular disease disability-adjusted life years (DALYs) attributable to high fasting plasma glucose from 1990 to 2023 due to population growth, population aging, risk exposure, and risk-deleted DALYs. Risk-deleted DALYs are the number of DALYs left after removing the effect of risk factors, population growth, and population aging on overall DALYs. They were calculated as the cardiovascular disease DALY count multiplied by 1 minus the population attributable fraction for high fasting plasma glucose. The dot and error bar represent the mean and 95% uncertainty interval in percentage change in number of DALYs attributable to the risk from 1990 to 2023.

with small decreases occurring in high SDI locations that continue to have high exposure to high LDL-C and increases in low SDI locations where exposure is lower.

There were a total of 90.7 million (95% UI: 59.0 to 123 million) cardiovascular DALYs attributed to high LDL-C in 2023 (Supplemental Table 7). Age-standardized attributable cardiovascular DALYs in 2023 were 1,002.6 (95% UI: 653.3 to 1,363.9) per 100,000 globally, with rates highest in Eastern Europe, at 2,252.6 (95% UI: 1,507.4 to 3,038.2) per 100,000, and in Oceania, at 2,200.7 (95% UI: 1,468.7 to 3,027.5) per 100,000, and rates lowest in high-income Asia Pacific, at 322.0 (95% UI: 208.7 to 446.6) per 100,000 (Figure 20-1). High LDL-C was attributed to 2.61 million (95% UI: 1.54 to 3.80 million) cardiovascular deaths in 1990, and age-

standardized cardiovascular mortality attributed to high LDL-C was 74.3 (95% UI: 42.7 to 109.0) per 100,000 (Supplemental Table 9). By 2023, the number of attributable cardiovascular deaths had increased to 3.63 million (95% UI: 2.23 to 5.36 million), while age-standardized attributable cardiovascular mortality decreased to 40.5 (95% UI: 24.9 to 59.9) per 100,000.

Globally, trends in LDL-C level did not contribute to a change in the associated CVD DALYs. However, change in risk-deleted DALY rates did substantially reduce burden attributable to LDL-C by 41.3 million (95% UI: 24.9 to 63.5 million) DALYs (Supplemental Table 8). Despite this, there was an overall increase in cardiovascular DALYs attributable to high LDL-C due to the effects of population growth and aging, which contributed to an increase of 33.4 million

FIGURE 20-1 Age-Standardized CVD DALY Rates Attributable to High Low-Density Lipoprotein Cholesterol, 2023

Age-standardized cardiovascular disease (CVD) disability-adjusted life years (DALYs) attributable to high low-density lipoprotein cholesterol per 100,000 in 2023 (all sexes combined).


(95% UI: 21.1 to 47.0 million) and 33.9 million (95% UI: 20.7 to 49.3 million) DALYs, respectively. However, changes in high LDL-C exposure varied by SDI (Figure 20-2). Trends in age-standardized SEVs were fairly flat for high SDI regions, with a slight decline from 52.4 (95% UI: 44.2 to 61.5) in 1990 to 49.8 (95% UI: 42.4 to 58.0) in 2023 (Supplemental Table 11). Conversely, exposure increased slightly in low SDI regions from 29.7 (95% UI: 21.7 to 39.4) to 34.6 (95% UI: 26.1 to 44.9).

High SBP. Burden attributable to high SBP was estimated for 8 CVDs, including IHD, ischemic stroke, intracerebral hemorrhage, and hypertensive heart disease, through the comparative risk assessment framework. High SBP was the largest contributor to CVD DALYs in 1990 and 2023 with 48.1% (95% UI: 36.3% to 58.0%) and 51.1% (95% UI: 41.8% to 58.5%) of CVD DALYs, respectively, attributed to it (Supplemental Table 8). The burden of high SBP has remained high, with exposure continuing to increase in many locations since 1990.

In 1990, there were 154 million (95% UI: 117 to 190 million) cardiovascular DALYs attributed to high SBP

(Supplemental Table 7). By 2023, this had increased to 223 million (95% UI: 180 to 261 million). However, age-standardized cardiovascular DALYs attributed to high SBP per 100,000 decreased from 4,044.9 (95% UI: 3,122.8 to 4,975.3) in 1990 to 2,454.9 (95% UI: 1,978.6 to 2,872.0) in 2023. Age-standardized cardiovascular mortality attributed to high SBP similarly declined from 205.8 (95% UI: 162.4 to 245.7) per 100,000 in 1990 to 117.9 (95% UI: 95.2 to 138.9) per 100,000 in 2023, while the number of attributable deaths increased from 7.07 million (95% UI: 5.55 to 8.56 million) to 10.5 million (95% UI: 8.51 to 12.4 million) (Supplemental Table 9). In 2023, age-standardized cardiovascular DALYs attributed to high SBP per 100,000 were highest in Oceania and Central Asia, followed by Eastern Europe and North Africa and the Middle East, and lowest in high-income Asia Pacific and Australasia (Figure 21-1). Globally, exposure to high SBP has risen from 1990 to 2023, from an age-standardized SEV of 28.3 (95% UI: 16.4 to 43.1) for males and 26.1 (95% UI: 15.2 to 41.4) for females to 32.6 (27.1 to 38.7) for males and 27.8 (95% UI: 22.7 to 33.4) for females

FIGURE 20-2 ARC in SEVs for High Low-Density Lipoprotein Cholesterol by Country and Territory, 1990 to 2023




Annualized rate of change in summary exposure value (SEV) for high low-density lipoprotein cholesterol from 1990 to 2023 for all sexes by Socio-demographic Index (SDI) (ranging from 0 to 1). SEVs are a measure of risk-weighted exposure in the population with a possible range of 0 to 100. A value of 0 signifies that no excess risk exists in the population due to exposure to the specified risk factor, while 100 indicates when the population is at the highest level of risk. The dashed line and the bold circle indicate the global average annualized rate of change (ARC) of the SEV. The bold circle represents the global SDI, and SEV in 2023. Labels represent the 3-digit International Organization for Standardization country code.

(Supplemental Table 11). Despite this slight global increase, there were some locations for which exposure declined, particularly high-income Asia Pacific.

The increase in number of cardiovascular DALYs attributable to high SBP globally from 1990 to 2023 was driven by population growth and aging, with an increase of 78.5 million (95% UI: 58.7 to 96.1 million) and 94.3 million (95% UI: 77.3 to 110 million) in cardiovascular DALYs, respectively. Changes in global high SBP risk exposure contributed little to no change in DALYs (Figure 21-2, Supplemental Table 8). These sources of increases were opposed by changes due to risk-deleted DALY rates, which contributed to a decrease of 111 million (95% UI: 70.2 to 156 million) cardiovascular DALYs attributable to high SBP.

Among the 8 CVDs, changes in high SBP exposure contributed to percentage increases in DALYs for intracerebral hemorrhage and subarachnoid hemorrhage and contributed to decreases in DALYs for other causes, such as hypertensive heart disease, atrial fibrillation and flutter, and aortic aneurysm.

High BMI. In the GBD study, the burden attributable to high BMI was estimated for 8 CVDs. High BMI was the seventh-largest contributor to CVD DALYs, responsible for 10.6% (95% UI: 5.9% to 14.9%) of all CVD DALYs in 2023 (Supplemental Table 8). This represents a rise from 1990, in which high BMI was responsible for 6.8% (95% UI: 3.7% to 10.3%) of all CVD DALYs. Substantial global increases in exposure to high BMI since 1990 have contributed to a rise in

FIGURE 21-1 Age-Standardized CVD DALY Rates Attributable to High Systolic Blood Pressure, 2023

Age-standardized CVD DALYs attributable to high systolic blood pressure per 100,000 in 2023 (all sexes combined). Abbreviations as in [Figure 20-1](#).

CVD DALYs, representing a major determinant in the increase in CVD DALYs along with the impacts of population growth and aging.


The total number of CVD DALYs attributable to high BMI globally increased from 21.6 million (95% UI: 11.3 to 33.1 million) in 1990 to 46.1 million (95% UI: 25.2 to 66.2 million) in 2023, while age-standardized DALYs per 100,000 decreased from 537.2 (95% UI: 281.8 to 824.1) per 100,000 to 508.5 (95% UI: 278.2 to 731.5) per 100,000 ([Supplemental Table 7](#)). Despite this global decline, there were a number of regions for which age-standardized DALYs per 100,000 increased, such as South Asia, which had an annualized rate of change of 3.7% (95% UI: 2.3% to 5.1%). North Africa and the Middle East had the highest age-standardized CVD DALY rate attributable to high BMI in 2023, at 1,527.1 (95% UI: 844.0 to 2,177.1) per 100,000, while high-income Asia Pacific had the lowest, at 104.3 (95% UI: 50.8 to 171.7) per 100,000. Exposure to high BMI has increased notably since 1990 for all countries ([Figure 22-1](#)). Globally, the age-standardized SEV for high BMI in males increased from 12.5 (95% UI: 10.5 to 14.7) in 1990 to 19.2 (95% UI: 17.5 to 21.2) in 2023 ([Supplemental Table 11](#)). For females, the summary exposure to

high BMI increased from 14.0 (95% UI: 11.7 to 16.4) in 1990 to 21.6 (95% UI: 19.7 to 23.8) in 2023.

Since 1990, the number of CVD DALYs attributable to high BMI increased by 113.8% (95% UI: 79.3% to 165.8%). A total of 17.1 million (95% UI: 7.97 to 28.3 million) of the increase in DALYs were due to increasing high BMI exposure ([Figure 22-2](#), [Supplemental Table 8](#)). Similarly, 11.0 million (95% UI: 5.88 to 16.9 million) and 16.6 million (95% UI: 8.93 to 24.0 million) of the increase in DALYs were due to population growth and aging, respectively. These increases were countered by change due to risk-deleted DALY rates, which contributed to a decrease of 20.1 million (95% UI: 10.3 to 30.4 million) DALYs. Increasing high BMI exposure contributed to the largest percentage increase in DALYs for intracerebral hemorrhage and subarachnoid hemorrhage, followed by atrial fibrillation and hypertensive heart disease.

Kidney dysfunction. In the GBD study, the burden attributable to kidney dysfunction was estimated for 4 CVDs: IHD, ischemic stroke, intracerebral hemorrhage, and lower extremity peripheral arterial disease. In addition to CVDs, kidney dysfunction was also a large contributor to chronic kidney disease burden. Kidney dysfunction was the seventh-largest

FIGURE 21-2 Percentage Change in the Number of Global Risk-Attributable DALYs, 1990 to 2023, due to Population Growth, Population Aging, Changes in Exposures to Each Global Burden of Disease Risk Factor, and Changes in Risk-Deleted DALY Rates for All Sexes, for High Systolic Blood Pressure




Decomposition of change in all-age, all sexes combined cardiovascular disease disability-adjusted life years (DALYs) attributable to high systolic blood pressure from 1990 to 2023 due to population growth, population aging, risk exposure, and risk-deleted DALYs. Risk-deleted DALYs are the number of DALYs left after removing the effect of risk factors, population growth, and population aging on overall DALYs. They were calculated as the cardiovascular disease DALY count multiplied by 1 minus the population attributable fraction for high systolic blood pressure. The dot and error bar represent the mean and 95% uncertainty interval in percentage change in number of DALYs attributable to the risk from 1990 to 2023.

contributor to cardiovascular DALYs in 1990 and the eighth largest in 2023, with 9.2% (95% UI: 6.9% to 11.6%) of cardiovascular DALYs attributed to it in 1990 and 10.2% (95% UI: 7.6% to 12.8%) in 2023 (Supplemental Table 8).

Age-standardized cardiovascular DALYs per 100,000 attributable to kidney dysfunction declined over time, decreasing from 789.1 (95% UI: 590.8 to 992.8) per 100,000 in 1990 to 491.3 (95% UI: 367.5 to 638.0) per 100,000 in 2023, despite an increase in number of DALYs from 29.5 million (95% UI: 22.4 to 37.0 million) to 44.6 million (95% UI: 33.5 to 57.8 million) (Supplemental Table 7). Age-standardized DALYs per 100,000 in 2023 were highest in Central Asia, at 1,155.2 (95% UI: 896.7 to 1,419.3) per 100,000, and Oceania, at 1,121.4 (95% UI: 848.0 to 1,495.0) per 100,000 (Figure 23-1). Rates were lowest in

Australasia, which had age-standardized attributable cardiovascular DALYs of 120.9 (95% UI: 87.1 to 159.5) per 100,000. Globally, exposure to kidney dysfunction was similar in 1990 and 2023 at an age-standardized SEV of 2.9 (95% UI: 1.9 to 4.9) and 2.9 (95% UI: 2.0 to 4.9), respectively (Supplemental Table 11). In both 1990 and 2023, the SEV was highest in Central Latin America (4.9 [95% UI: 3.0 to 7.5] and 5.1 [95% UI: 3.1 to 7.7], respectively) and the Caribbean (4.7 [95% UI: 2.9 to 7.2] and 5.0 [95% UI: 3.1 to 7.6], respectively).

Between 1990 and 2023, the number of cardiovascular DALYs attributable to kidney dysfunction globally increased by 51.0% (95% UI: 37.5% to 66.8%) (Supplemental Table 8). This increase was primarily due to the effects of population growth and aging, which contributed 15.1 million (95% UI: 11.3 to 19.1

FIGURE 22-1 ARC in SEVs for High Body Mass Index by Country and Territory, 1990 to 2023


Annualized rate of change in SEV for high body mass index from 1990 to 2023 for all sexes by SDI (ranging from 0 to 1). SEVs are a measure of risk-weighted exposure in the population with a possible range of 0 to 100. A value of 0 signifies that no excess risk exists in the population due to exposure to the specified risk factor, while 100 indicates when the population is at the highest level of risk. The dashed line and the bold circle indicate the global average ARC of the SEV. The bold circle represents the global SDI, and SEV in 2023. Labels represent the 3-digit International Organization for Standardization country code. Abbreviations as in [Figure 20-2](#).

million) and 19.3 million (95% UI: 14.5 to 24.8 million) DALYs, respectively ([Figure 23-2](#)). Changes in exposure to kidney dysfunction contributed to a smaller change of 2.03 million (95% UI: -0.0509 to 4.07 million) DALYs, the UI barely crossed zero, suggesting a potential increase in DALYs due to exposure. Decreases in DALYs were solely due to changes in risk-deleted DALY rates, which contributed to a decrease of 21.4 million (95% UI: 15.3 to 27.7 million) DALYs.

Tobacco. Tobacco causes 7 CVDs, including atrial fibrillation and flutter, lower extremity arterial disease, and aortic aneurism. In 2023, tobacco was the fifth-leading cause of CVD DALYs among all risks for CVD, decreasing from fourth place in 1990. The percentage of CVD DALYs attributable to tobacco decreased from 21.2% (95% UI: 18.5% to 24.1%) in 1990 to 17.9% (95% UI: 15.5% to 20.3%).

The number of CVD DALYs attributable to tobacco increased from 67.8 million (95% UI: 58.3 to 78.3 million) in 1990 to 78.0 million (95% UI: 67.9 to 88.8 million) in 2023, although the age-standardized rate per 100,000 population decreased from 1,650.5 (95% UI: 1,418.8 to 1,903.0) to 855.6 (95% UI: 743.7 to 971.9) ([Supplemental Figure 18](#)). By world region, the number of DALYs attributable to tobacco ranged from 118,000 (95% UI: 96,600 to 141,000) in Australasia to 22.4 million (95% UI: 18.7 to 26.1 million) in East Asia in 2023; per 100,000 population DALYs were largest in Oceania (2,066.5 [95% UI: 1,630.0 to 2,613.3]) and lowest in Andean Latin America (229.7 [95% UI: 176.8 to 296.7]) ([Figure 24-1](#)). Age-standardized DALYs attributable to tobacco per 100,000 decreased the most from 1990 to 2023 in Australasia and Western Europe, with an annualized rate of change decrease of 4.9% (95% UI: 4.6% to

FIGURE 22-2 Percentage Change in the Number of Global Risk-Attributable DALYs, 1990 to 2023, due to Population Growth, Population Aging, Changes in Exposures to Each Global Burden of Disease Risk Factor, and Changes in Risk-Deleted DALY Rates for All Sexes, for High Body Mass Index




Decomposition of change in all-age, all sexes combined cardiovascular disease disability-adjusted life years (DALYs) attributable to high body mass index from 1990 to 2023 due to population growth, population aging, risk exposure, and risk-deleted DALYs. Risk-deleted DALYs are the number of DALYs left after removing the effect of risk factors, population growth, and population aging on overall DALYs. They were calculated as the cardiovascular disease DALY count multiplied by 1 minus the population attributable fraction for high body mass index. The dot and error bar represent the mean and 95% uncertainty interval in percentage change in number of DALYs attributable to the risk from 1990 to 2023.

5.3%) and by 3.8% (95% UI: 3.5% to 4.1%), respectively.

From 1990 to 2023, tobacco exposure decreased globally from a SEV of 34.5 (95% UI: 32.4 to 37.1) for males and 23.0 (95% UI: 21.5 to 24.7) for females in 1990, to 26.1 (95% UI: 24.5 to 28.2) for males and 15.8 (95% UI: 14.5 to 17.4) for females in 2023. The decrease in tobacco exposure led to a reduction of 21.2 million (95% UI: 11.3 to 31.2 million) cardiovascular DALYs (Figure 24-2). Similarly, the risk-deleted DALY rate contributed to a decrease of 31.6 million (95% UI: 19.7 to 41.2 million) DALYs. In contrast, population growth (34.6 million [95% UI: 29.4 to 39.6 million]) and population aging (28.3 million [95% UI: 24.2 to 32.9 million]) contributed to a rise in DALYs, yielding an overall increase in the number of DALYs attributable to tobacco exposure of 15.0% (95% UI: 0.3% to 34.6%) from 1990 to 2023. The largest

proportional decreases in DALYs due to tobacco exposure only by CVD subcategory from 1990 to 2023 were for lower extremity peripheral arterial disease and aortic aneurism, where DALYs due to tobacco exposure decreased by 59.3% (95% UI: 36.7% to 82.6%) and 58.7% (95% UI: 37.2% to 81.2%) respectively.

High alcohol use. High alcohol use causes 6 CVDs: alcoholic cardiomyopathy, atrial fibrillation, hypertensive heart disease, intracerebral hemorrhage, IHD, and ischemic stroke. For IHD, the GBD study estimated a J-shaped risk curve for burden attributable to high alcohol use and therefore uses a theoretical minimum level that accounts for the possibility of protective effects. In contrast, the risks of the other outcomes are estimated as increasing from low levels of alcohol consumption. In 1990 and 2023, high alcohol use ranked 12th among all risks for CVD

FIGURE 23-1 Age-Standardized CVD DALY Rates Attributable to Kidney Dysfunction, 2023

Age-standardized CVD DALYs attributable to kidney dysfunction per 100,000 in 2023 (all sexes combined). Abbreviations as in [Figure 20-1](#).

DALYs. The percentage of CVD DALYs attributable to high alcohol use remained similar from 1990 (0.9% [95% UI: 0.0% to 2.9%]) to 2023 (0.9% [95% UI: 0.1% to 2.5%]).


The number of CVD DALYs attributable to high alcohol use increased from 2.86 million (95% UI: 0.0328 to 9.34 million) in 1990 to 3.83 million (95% UI: 0.411 to 11.0 million) in 2023 ([Supplemental Figure 19](#)). In contrast, the age-standardized rate per 100,000 population of DALYs attributable to high alcohol use decreased from 75.3 (95% UI: 8.4 to 227.5) in 1990 to 41.8 (95% UI: 4.0 to 121.4) in 2023. By world region in 2023, the number of DALYs attributable to high alcohol use was highest in Eastern Europe (1.34 million [95% UI: 1.01 to 2.37 million]) and East Asia (1.31 million [95% UI: -0.172 to 3.36 million]); per 100,000 population, DALYs were largest in Eastern Europe (445.1 [95% UI: 341.6 to 750.1]), followed by the Caribbean (106.3 [95% UI: 35.7 to 229.0]) ([Figure 25-1](#)).

From 1990 to 2023, the global exposure to high alcohol use decreased from a SEV of 9.1 (95% UI: 6.8 to 12.7) for males and 4.6 (95% UI: 3.1 to 6.4) for females, to 8.0 (95% UI: 6.0 to 10.8) and 3.4 (95% UI: 2.5

to 4.5), respectively. The decrease in high alcohol exposure contributed to a change of -0.36 million (95% UI: -2.58 to 0.983 million) DALYs ([Figure 25-2](#)), the UI largely overlapped with zero, suggesting the possibility of no notable change due to risk exposure changes. Nevertheless, there was an overall increase in the number of CVD DALYs attributable to high alcohol use driven by increases due to population aging (95% UI: 1.81 million [95% UI: 0.616 to 4.15 million]) and population growth (1.46 million [0.0168 to 4.68 million] DALYs). These increases counteracted the effect of decreased exposure, as well as that of the risk-deleted DALYs (-1.93 million [95% UI: -4.91 to -0.439 million]). By CVD subcategory, changes in exposure to high alcohol use contributed to the greatest decrease in the number of DALYs for alcoholic cardiomyopathy with a decrease of 63.6% (95% UI: 48.3% to 75.7%) DALYs from 1990 to 2023.

DIETARY RISKS. Dietary risks consist of 15 dietary risk subtypes ([Table 2](#)), 13 of which cause 8 CVDs including atrial fibrillation and flutter, aortic aneurism, and lower extremity peripheral arterial disease. Dietary risks were the second-leading attributable

FIGURE 23-2 Percentage Change in the Number of Global Risk-Attributable DALYs, 1990 to 2023, due to Population Growth, Population Aging, Changes in Exposures to Each Global Burden of Disease Risk Factor, and Changes in Risk-Deleted DALY Rates for All Sexes, for Kidney Dysfunction




Decomposition of change in all-age, all sexes combined cardiovascular disease disability-adjusted life years (DALYs) attributable to kidney dysfunction from 1990 to 2023 due to population growth, population aging, risk exposure, and risk-deleted DALYs. Risk-deleted DALYs are the number of DALYs left after removing the effect of risk factors, population growth, and population aging on overall DALYs. They were calculated as the cardiovascular disease DALY count multiplied by 1 minus the population attributable fraction for kidney dysfunction. The dot and error bar represent the mean and 95% uncertainty interval in percentage change in number of DALYs attributable to the risk from 1990 to 2023.

risk factor among all risks for CVD DALYs in both 1990 and 2023. The percentage of CVD DALYs attributable to dietary risks remained similar from 1990 (33.5% [95% UI: 13.2% to 44.5%]) to 2023 (32.3% [95% UI: 12.5% to 45.5%]).

The number of DALYs attributable to diet increased from 107 million (95% UI: 42.1 to 147 million) in 1990 to 141 million (95% UI: 56.0 to 198 million) in 2023 (Supplemental Figure 20). However, the age-standardized DALY rate per 100,000 population decreased from 2,694.9 (95% UI: 1,055.0 to 3,717.3) in 1990 to 1,557.3 (95% UI: 617.8 to 2,184.3) in 2023. In 2023, large regional variations were observed, with the number of DALYs ranging from 257,000 (95% UI: 76,700 to 379,000) DALYs in Australasia to 38.1 million (95% UI: 10.3 to 56.4 million) in South Asia and 30.1 million (95% UI: 15.5 to 42.4

million) in East Asia; age-standardized DALYs per 100,000 population were largest in Oceania (4,122.8 [95% UI: 1,434.6 to 5,961.0]) and lowest in high-income Asia Pacific (434.3 [95% UI: 224.6 to 609.7]) (Figure 26-1). CVD DALYs attributable to dietary risks per 100,000 decreased the most in Australasia and Western Europe, with an annualized rate of change decrease of 4.2% (95% UI: 3.7% to 4.6%) and 3.3% (95% UI: 2.8% to 3.7%), respectively.

From 1990 to 2023, the global SEV for dietary risks modestly decreased from 41.2 (95% UI: 32.5 to 49.0) for males and 38.8 (95% UI: 28.1 to 46.2) for females in 1990 to 39.9 (95% UI: 30.3 to 47.4) for males and 37.0 (95% UI: 27.7 to 44.1) for females in 2023. The decreased exposure in dietary risks contributed to a change of -13.1 million (95% UI: -38.4 to 15.7 million) DALYs, though a wide UI that overlaps with

FIGURE 24-1 Age-Standardized CVD DALY Rates Attributable to Tobacco, 2023

Age-standardized CVD DALYs attributable to tobacco per 100,000 in 2023 (all sexes combined). Abbreviations as in [Figure 20-1](#).


zero suggests the possibility of no change ([Figure 26-2](#)). While this value is informed by a composite of the exposure from all dietary factors causing CVD, the individual dietary risks may have an effect in opposite directions. Specifically, the harmful exposure to sugar-sweetened beverages increased since 1990, while other harmful dietary risk exposure decreased (ie, low fruit consumption). Population growth (54.8 million [95% UI: 21.7 to 75.9 million] DALYs) and population aging (53.2 million [95% UI: 21.6 to 77.4 million] DALYs) contributed to a substantial increase in CVD DALYs attributable to diet, while the risk-deleted DALY rate accounted for a decrease of 61.3 million (95% UI: 22.1 to 95.3 million) DALYs. Altogether, dietary risks led to an overall increase of attributable CVD DALYs of 31.4% (95% UI: 9.7% to 52.5%) from 1990 to 2023.

Low physical activity. In the GBD study's comparative risk assessment framework, the burden attributable to low physical activity was estimated for 3 CVDs: IHD, ischemic stroke, and lower extremity peripheral arterial disease. In both 1990 and 2023, it

was the second-smallest contributor to cardiovascular DALYs. However, it still accounted for 1.5% (95% UI: 0.6% to 2.5%) and 1.6% (95% UI: 0.6% to 2.6%) of all cardiovascular DALYs in 1990 and 2023, respectively ([Supplemental Table 8](#)).

The number of cardiovascular DALYs attributable to low physical activity increased from 4.94 million (95% UI: 1.93 to 7.87 million) in 1990 to 7.16 million (95% UI: 2.87 to 11.6 million) in 2023 ([Supplemental Table 7](#)). However, attributable age-standardized cardiovascular DALYs per 100,000 decreased from 133.1 (95% UI: 48.7 to 218.0) per 100,000 in 1990 to 78.8 (95% UI: 31.5 to 129.0) per 100,000 in 2023. In 2023, rates were highest in North Africa and Middle East with 210.0 DALYs (95% UI: 84.3 to 332.1 DALYs) per 100,000 ([Figure 27-1](#)). Andean Latin America and Southern Latin America had the lowest rates with 20.0 (95% UI: 6.6 to 33.8) and 19.0 (95% UI: 6.3 to 34.2) DALYs per 100,000, respectively. Globally, exposure to low physical activity increased over time, though this change was not substantial ([Figure 27-2](#); [Supplemental Table 11](#)). From 1990 to

FIGURE 24-2 Percentage Change in the Number of Global Risk-Attributable DALYs, 1990 to 2023, due to Population Growth, Population Aging, Changes in Exposures to Each Global Burden of Disease Risk Factor, and Changes in Risk-Deleted DALY Rates for All Sexes, for Tobacco



Decomposition of change in all-age, all sexes combined cardiovascular disease disability-adjusted life years (DALYs) attributable to tobacco from 1990 to 2023 due to population growth, population aging, risk exposure, and risk-deleted DALYs. Risk-deleted DALYs are the number of DALYs left after removing the effect of risk factors, population growth, and population aging on overall DALYs. They were calculated as the cardiovascular disease DALY count multiplied by 1 minus the population attributable fraction for tobacco. The dot and error bar represent the mean and 95% uncertainty interval in percentage change in number of DALYs attributable to the risk from 1990 to 2023.


2023, the age-standardized SEV rose from 20.6 (95% UI: 17.5 to 23.6) for females and 12.8 (95% UI: 10.3 to 15.4) for males to 23.0 (19.5 to 26.7) for females and 13.9 (11.3 to 16.6) for males.

The increase in number of cardiovascular DALYs attributable to low physical activity over time was primarily driven by the effects of population growth and aging, which contributed to an increase of 2.52 million (95% UI: 1.01 to 4.04 million) DALYs and 3.13 million (95% UI: 0.989 to 5.27 million) DALYs, respectively (Supplemental Table 8). Increasing exposure to low physical activity contributed little to no change in DALYs at only 0.218 million (95% UI: -0.623 to 1.05 million) DALYs added since 1990 due to changing exposure. These increases were countered by changes due to risk-deleted DALY rates, which contributed to a decrease of 3.65 million (95%

UI: 1.38 to 5.87 million) DALYs. Among the 3 CVDs, changes due to low physical activity contributed to the largest percentage increase for IHD as well as to a large percentage decrease for lower extremity peripheral arterial disease.

Air pollution. In the GBD study, burden attributable to air pollution is estimated for 4 CVDs: IHD, ischemic stroke, intracerebral hemorrhage, and subarachnoid hemorrhage. It was the third- and fourth-largest contributor to CVDs in 1990 and 2023, responsible for 25.5% (95% UI: 20.0% to 31.3%) and 20.7% (95% UI: 16.2% to 25.2%) of all cardiovascular DALYs, respectively (Supplemental Table 8).

In 1990, there were 81.7 million (95% UI: 61.2 to 103 million) cardiovascular DALYs and 3.46 million (95% UI: 2.59 to 4.34 million) cardiovascular deaths attributable to air pollution (Figure 28-1). By 2023,

FIGURE 25-1 Age-Standardized CVD DALY Rates Attributable to High Alcohol Use, 2023


Age-standardized CVD DALYs attributable to high alcohol use per 100,000 in 2023 (all sexes combined). Abbreviations as in [Figure 20-1](#).

this had increased to 90.5 million (95% UI: 70.7 to 110 million) DALYs and 3.97 million (95% UI: 3.09 to 4.83 million) deaths. However, despite a slight increase between 2020 and 2021, attributable age-standardized cardiovascular DALYs and deaths per 100,000 have steadily decreased from 2,077.5 (95% UI: 1,552.4 to 2,610.8) per 100,000 and 96.8 (95% UI: 73.1 to 121.2) per 100,000 in 1990 to 996.6 (95% UI: 778.6 to 1,211.2) per 100,000 and 44.1 (95% UI: 34.2 to 53.8) per 100,000 in 2023, respectively. Age-standardized rates of attributable cardiovascular DALYs in 2023 were highest in Oceania, at 2,961.0 (95% UI: 2,156.0 to 3,759.7) per 100,000, and lowest in Australasia, at 50.9 (95% UI: 26.3 to 80.8) per 100,000 ([Supplemental Table 7](#)). Although global exposure to air pollution declined from an age-standardized SEV of 51.5 (95% UI: 44.4 to 58.5) in 1990 to 37.6 (95% UI: 31.9 to 44.1) in 2023 ([Supplemental Table 11](#)), this change was not consistent by location ([Figure 28-2](#)). Decreases were largest in high SDI regions, with an annualized rate of change of -1.9% (95% UI: -2.2% to -1.4%) from a SEV of 40.7 (95% UI: 30.0 to 50.4) to 22.2 (95% UI: 15.6 to 30.0) between 1990 and 2023. Conversely, low SDI regions experienced a decrease of only 0.3% (95% UI:

0.3% to 0.4%), despite a substantially higher SEV of 69.1 (95% UI: 63.8 to 74.7) in 1990 and 61.8 (95% UI: 56.4 to 67.5) in 2023. Additionally, changes between 1990 and 2023 were not consistent for types of air pollution. From 1990 to 2023, CVD DALYs due to ambient particulate matter pollution rose from 33.3 million (95% UI: 23.4 to 42.9 million) to 60.4 million (95% UI: 47.2 to 76.0 million); in contrast, those associated with household air pollution from solid fuels decreased from 48.5 million (95% UI: 35.1 to 63.4 million) to 30.3 million (95% UI: 20.8 to 43.3 million).

Globally, the increase in number of cardiovascular DALYs attributable to air pollution was driven primarily by the effects of population growth and aging, which contributed 41.7 million (95% UI: 30.9 to 51.8) and 36.2 million (95% UI: 28.0 to 44.8 million) DALYs, respectively ([Supplemental Table 8](#)). Changes in air pollution exposure resulted in a decrease of 28.5 million (95% UI: 15.7 to 42.9 million) DALYs, which was similar to the effects of risk-deleted DALY rates, which contributed a decrease of 40.6 million (95% UI: 29.0 to 51.8 million) DALYs. Exposure contributed to the largest percentage decrease in DALYs for subarachnoid hemorrhage followed by

FIGURE 25-2 Percentage Change in the Number of Global Risk-Attributable DALYs, 1990 to 2023, due to Population Growth, Population Aging, Changes in Exposures to Each Global Burden of Disease Risk Factor, and Changes in Risk-Deleted DALY Rates for All Sexes, for High Alcohol Use




Decomposition of change in all-age, all sexes combined cardiovascular disease disability-adjusted life years (DALYs) attributable to high alcohol use from 1990 to 2023 due to population growth, population aging, risk exposure, and risk-deleted DALYs. Risk-deleted DALYs are the number of DALYs left after removing the effect of risk factors, population growth, and population aging on overall DALYs. They were calculated as the cardiovascular disease DALY count multiplied by 1 minus the population attributable fraction for high alcohol use. The dot and error bar represent the mean and 95% uncertainty interval in percentage change in number of DALYs attributable to the risk from 1990 to 2023.

intracerebral hemorrhage, which, along with the effects of risk-deleted DALY rates, were large enough to outweigh the increases from population growth and aging, resulting in a decline in the number of DALYs attributable to air pollution for these causes between 1990 and 2023.

Non-optimal temperature. Burden attributable to non-optimal temperature is estimated for 8 CVDs in the GBD study, including IHD, ischemic stroke, intracerebral hemorrhage, and hypertensive heart disease. It was the 10th-largest contributor to cardiovascular DALYs in both 1990 and 2023. A total of 5.9% (95% UI: 5.4% to 7.0%) of cardiovascular DALYs were attributed to non-optimal temperature in 1990 and 5.3% (95% UI: 4.5% to 6.5%) in 2023 (Supplemental Table 8).

In 2023, there were 22.9 million (95% UI: 19.2 to 28.7 million) cardiovascular DALYs and 1.16 million (95% UI: 0.968 to 1.41 million) cardiovascular deaths attributable to non-optimal temperature, reflecting an increase from 19.0 million (95% UI: 16.6 to 22.7 million) cardiovascular DALYs and 0.875 million (95% UI: 0.776 to 1.02 million) cardiovascular deaths in 1990 (Supplemental Tables 7 and 9). Between 1990 and 2023, global age-standardized cardiovascular DALYs attributable to non-optimal temperature per 100,000 decreased from 489.2 (95% UI: 429.8 to 577.4) to 254.7 (95% UI: 212.7 to 318.7). Among regions, rates were highest in Central Asia, North Africa and the Middle East, and Eastern Europe, with 601.3 (95% UI: 536.5 to 758.6) DALYs, 575.2 (95% UI: 433.7 to 779.8) DALYs, and 498.6 (95% UI: 463.7 to 550.9)

FIGURE 26-1 Age-Standardized CVD DALY Rates Attributable to Dietary Risks, 2023

Age-standardized CVD DALYs attributable to dietary risks per 100,000 in 2023 (all sexes combined). Abbreviations as in [Figure 20-1](#).

DALYs per 100,000, respectively ([Figure 29-1](#)). Rates were lowest in the Caribbean and Central Sub-Saharan Africa with 53.0 (95% UI: 33.8 to 74.9) DALYs and 51.9 (95% UI: -47.7 to 96.5) DALYs per 100,000, respectively. Globally, exposure to non-optimal temperature increased over time from an age-standardized SEV of 28.5 (95% UI: 27.8 to 29.6) in 1990 to 32.2 (95% UI: 31.0 to 33.2) in 2023 ([Supplemental Table 11](#)). However, global trends differed by type of non-optimal temperature, with exposure to low temperature having little to no change while exposure to high temperature increased substantially.


Between 1990 and 2023, the number of cardiovascular DALYs attributable to non-optimal temperature increased by 20.6% (95% UI: 6.5% to 36.2%) ([Supplemental Table 8](#)). This increase was driven by the effects of population growth and aging, which contributed to increases of 9.71 million (95% UI: 8.55 to 11.6 million) DALYs and 9.23 million (95% UI: 7.74 to 11.5 million) DALYs, respectively ([Figure 29-2](#)). Changes in exposure to non-optimal temperature contributed to a decrease of 3.43 million (95% UI: 2.02 to 4.87 million) DALYs, reflecting the larger risk

attribution of low temperature compared with high temperature for CVD burden. Changes due to risk-deleted DALY rate were larger, contributing to a decrease of 11.6 million (95% UI: 8.50 to 15.0 million) DALYs. Changes in exposure to non-optimal temperature contributed to the largest percentage decreases for ischemic stroke, subarachnoid hemorrhage, and hypertensive heart disease.

Lead exposure. Lead exposure is a subgroup of the category of other environmental risks in GBD. Lead exposure itself consists of lead exposure in bone and lead exposure in blood; CVD was only attributed to lead exposure in bone reflecting the long-term effect of lead accumulation. Lead exposure caused 8 CVDs in the GBD via its effect mediated through high SBP, including IHD, ischemic stroke, and intracerebral hemorrhage. Lead exposure was the sixth largest cause of CVD DALYs when compared with level 2 GBD CVD risk factors, just below tobacco. There was notable geographic and sociodemographic variation in the exposure and the global attributable burden of CVD attributable to lead exposure ([Figure 30-1](#)).

There were 70.1 million (95% UI: 53.0 to 87.5 million) cardiovascular DALYs and 3.43 million (95%

FIGURE 26-2 Percentage Change in the Number of Global Risk-Attributable DALYs, 1990 to 2023, due to Population Growth, Population Aging, Changes in Exposures to Each Global Burden of Disease Risk Factor, and Changes in Risk-Deleted DALY Rates for All Sexes, for Dietary Risks




Decomposition of change in all-age, all sexes combined cardiovascular disease disability-adjusted life years (DALYs) attributable to dietary risks from 1990 to 2023 due to population growth, population aging, risk exposure, and risk-deleted DALYs. Risk-deleted DALYs are the number of DALYs left after removing the effect of risk factors, population growth, and population aging on overall DALYs. They were calculated as the cardiovascular disease DALY count multiplied by 1 minus the population attributable fraction for dietary risks. The dot and error bar represent the mean and 95% uncertainty interval in percentage change in number of DALYs attributable to the risk from 1990 to 2023.

UI: 2.59 to 4.25 million) cardiovascular deaths attributable to lead exposure in 2023 globally. These increased from 41.2 million (95% UI: 30.8 to 53.8 million) DALYs and 1.83 million (95% UI: 1.36 to 2.35 million) deaths in 1990. In contrast, the age-standardized DALY rate decreased from 1,069.6 (95% UI: 803.2 to 1,391.5) per 100,000 in 1990 to 772.1 (95% UI: 582.4 to 963.7) per 100,000 in 2023. There was notable geographic variation in the distribution of CVD DALYs attributable to lead exposure in 2023, ranging from 1,504.1 (95% UI: 1,259.1 to 1,764.8) per 100,000 in Central Asia and 1,474.3 (95% UI: 1,154.1 to 1,801.9) per 100,000 in North Africa and the Middle East, to 126.6 (95% UI: 95.6 to 161.2) per 100,000 in high-income Asia Pacific (Figure 30-1). From 1990 to 2023, DALYs attributable to lead exposure decreased the most in Australasia (annualized rate of

change: -4.2% [95% UI: -4.4% to -3.9%]), Western Europe (-3.2% [95% UI: -3.4% to -3.0%]), high-income North America (-3.0% [95% UI: -3.3% to -2.7%]), and Southern Latin America (-2.8% [95% UI: -3.0% to -2.6%]).

The exposure to lead did not substantially change from 54.2 (95% UI: 8.8 to 65.5) per 100 among males and 44.4 (95% UI: 7.5 to 56.1) per 100 among females in 1990 to 51.8 (95% UI: 7.7 to 60.5) per 100 in males and 44.5 (95% UI: 6.8 to 52.5) per 100 in females in 2023. The rise in the number of DALYs was driven by the effects of population growth and population aging, as well as the changing exposure to lead. Population growth and aging contributed to increases of 21.0 million (95% UI: 15.5 to 27.5 million) and 28.9 million (95% UI: 21.5 to 36.2 million) CVD DALYs, respectively, while changes in lead exposure led to

FIGURE 27-1 Age-Standardized CVD DALY Rates Attributable to Low Physical Activity, 2023

Age-standardized CVD DALYs attributable to low physical activity per 100,000 in 2023 (all sexes combined). Abbreviations as in [Figure 20-1](#).

an increase of 12.5 million (95% UI: 7.59 to 18.1 million) CVD DALYs globally in 2023 ([Figure 30-2](#)). The leading subcauses of CVD that contributed to this rise were IHD, ischemic stroke, and intracerebral hemorrhage.

DISCUSSION


OVERVIEW AND KEY FINDINGS. In 2023, CVD was the leading cause of disease burden and deaths worldwide (at level 2). Low, low-middle, and middle SDI regions had the greatest age-standardized rates of DALYs due to CVD, which was almost twice the rate estimated in locations in the high SDI region. This finding, that CVD burden is substantially greater outside of the most developed settings even after accounting for differences in population age, remains among the most important messages of this analysis and provides key evidence for the upcoming Fourth High-level Meeting of the UN General Assembly on the prevention and control of non-communicable diseases. Geographic variation in CVD burden is extensive, as shown in the maps throughout our analysis, and extends beyond any simple assessment

based on economic categories. Burden varies greatly for countries at the same level of SDI, suggesting a potential for gains in cardiovascular health even before broader development goals are achieved.

The global burden of CVD is not declining. This flat trend in the decline of CVD is due to trends in low-middle and middle SDI regions where populations are growing and aging at rates outpacing improvement in modifiable risk factors and other drivers of health. These trends are occurring despite remarkable improvements in the age-standardized rate of CVD burden in higher SDI regions in the first decade of this analysis, with an inflection point for many high SDI countries around 2010, when these improving trends began to flatten. The past decade has been one of stagnation for these countries, and most have had no substantial improvement in age-standardized rates of CVD burden over that period.

In 2023, CVD burden was attributable to numerous modifiable risk factors, with high SBP the dominant contributor, followed by non-optimal diet, high LDL-C, air pollution, tobacco use, lead exposure, BMI, kidney dysfunction, FPG, non-optimal temperature, low physical activity, and unsafe alcohol use.

FIGURE 27-2 Summary Exposure Value of Low Physical Activity From 1990 to 2023




Summary exposure value for low physical activity (ranging from 0 to 100) from 1990 to 2023 for males and females. Summary exposure values are a measure of risk-weighted exposure in the population with a possible range of 0 to 100. A value of 0 signifies that no excess risk exists in the population due to exposure to the specified risk factor, while 100 indicates when the population is at the highest level of risk.

The fastest-rising threats globally for CVD burden were metabolic (high BMI and FPG) and environmental (ambient air pollution and high non-optimal temperature) risks, while improvements occurred in burden attributable to high trans fats and household air pollution.

Risk factor attribution varied considerably between regions. For example, household air pollution was a larger cause of CVD in Oceania and SSA than other regions. High alcohol use was a larger cause of CVD in Eastern Europe than other regions. High sodium consumption was a larger cause of CVD in East Asia, Southeast Asia, Central Europe, and Oceania. Given this kind of variation, disease burden

estimates offer the opportunity to tailor local health policies to target the most relevant risks.

ATHEROSCLEROTIC CVD. Trends in IHD, ischemic stroke, and PAD represent the changing global burden of ASCVD and account for the majority of CVD burden, as well as for many of its principal epidemiologic features. Our decomposition analysis shows that the leading reason for the increase in ASCVD burden after population growth and aging was increasing high BMI and FPG. The substantial increase in ASCVD burden supports the need for early detection, prompt risk stratification, and accessible paths to lifestyle modification and treatment for the




full range of cardiovascular risk factors. High SBP accounts for more than half of ASCVD burden globally, indicating that hypertension screening, delivery of cost-effective pharmacotherapies, dietary sodium reduction, and reduction in environmental lead exposure remain essential goals.^{33,34}

The role of obesity and diabetes in ASCVD requires particular attention.^{35,36} A broader framing of the risk of high BMI and FPG has been called for in which malnutrition, including obesity, is considered a global syndemic. Our findings highlight the need for wider implementation of primary prevention strategies to increase healthy eating and physical activity as well as implementation of screening and monitoring systems. Efforts to stem the increase in high BMI have focused on behavioral prevention strategies at younger ages, including healthier diets, increased physical activity levels, and more recently,

incretin-based pharmacotherapies among individuals with obesity, diabetes, or CVD.³⁷ The syndemic approach suggests a need to expand our scope to identify drivers and impacts on a societal and ecological scale. Our findings emphasize that exposure levels to these complex, intertwined risks continue to rise even in the face of innovative therapies. Given that approximately half of the world is unaware of their diabetes, substantial attention will need to be paid to out-of-pocket costs and household economic burden if any headway is to be achieved.^{38,39}

Similar challenges exist to reduce levels of both high blood pressure and cholesterol, though gains have been made for some ASCVD risks. Effective policies are helping to reduce household air pollution, eliminate trans fat, and decrease access to tobacco.⁴⁰⁻⁴² New policies will be needed to address the




threat of environmental risks, including ambient particulate matter air pollution, non-optimal high temperature, and lead exposure.

STROKE. Age-standardized DALY rates for stroke, including ischemic, intracerebral hemorrhage, and subarachnoid hemorrhage, were highest in low, low-middle, and middle SDI settings, in which there are large gaps in access to diagnostic imaging, emergency transportation and treatment, and both critical and long-term care.^{43,44} While shared risk factors suggest a parallel approach to that of ASCVD, key policy priorities have been recommended beyond risk factor control, including expanding access to acute care facilities capable of treating stroke and improving rehabilitation capabilities for stroke survivors.⁴⁵ Out-of-pocket expenses, limited training for

physiotherapy, occupational therapists, and speech-language therapists, and focus on acute care have led to limited access for rehabilitation care in many parts of the world. Future declines in the health workforce, driven by global declines in fertility, may limit capacity to provide adequate care for individuals with brain injury following stroke as well as other CVD.⁴⁶⁻⁴⁸

CARDIOMYOPATHIES, VALVULAR DISEASE, AND HYPERTENSIVE HEART DISEASE. Cardiomyopathies include a diverse collection of myocardial diseases with a broad range of etiologies, including hypertension, alcohol use, valvular heart disease, and primary genetic disease.⁴⁹ The burden of the most common of these, hypertensive heart disease, has

FIGURE 29-1 Age-Standardized CVD DALY Rates Attributable to Non-optimal Temperature, 2023

Age-standardized CVD DALYs attributable to non-optimal temperature per 100,000 in 2023 (all sexes combined). Abbreviations as in [Figure 20-1](#).


not declined significantly in the past 3 decades. Access to blood pressure-lowering drugs has become more common; however, surveys of pharmacies show that availability of multiple drug classes or combination therapies correlates with blood pressure control but is substantially lower in low- and middle-income countries.⁵⁰ We observed wide variation in the rate of change in the burden of hypertensive heart disease even among countries at the same level of SDI, suggesting that there may be best practice strategies which can be adopted by health systems even for those at similar levels of national development. For example, a single-pill combination of 3 antihypertensives has been shown to be more effective and just as safe as 2 drugs, providing a strategy to improve the treatment of hypertension.⁵¹

Alcoholic cardiomyopathy remains an important target for policy interventions. An intake of at least 80 g per day of alcohol for over 5 years is generally considered the main risk factor for alcoholic cardiomyopathy, which aligns closely with the threshold that GBD estimates for increasing risk of IHD due to alcohol consumption.⁵² Risks of stroke and atrial fibrillation are estimated by GBD to increase at lower levels of consumption. Alcohol intake tends to start

in adolescence or early adulthood, which is consistent with alcoholic cardiomyopathy DALY rates peaking at 60 years of age, a younger age than we found for other CVDs. Alcohol intake is generally higher in men vs women, particularly in countries that preserve more traditional gender roles.⁵³ This aligns with our finding of higher burden from alcoholic cardiomyopathy in men vs women. Alcohol consumption trends are changing, with substantial declines in Europe associated with adoption of effective government policies but increases in Southeast Asia.⁵³

The global burden of calcific aortic valve disease rose significantly, while degenerative mitral valve disease burden remained stable. The GBD study also estimates disease burden due to tricuspid and pulmonary valve disease (in aggregate and excluding congenital heart diseases, which are estimated separately as part of congenital diseases) and found a trend toward increasing burden for both all-age and age-standardized rates. Improvements in surveillance and reporting will be needed to better understand how the availability of diagnostic imaging and options for therapies will impact future valvular heart disease burden estimates.⁵⁴

FIGURE 29-2 Percentage Change in the Number of Global Risk-Attributable DALYs, 1990 to 2023, due to Population Growth, Population Aging, Changes in Exposures to Each Global Burden of Disease Risk Factor, and Changes in Risk-Deleted DALY Rates for All Sexes, for Non-optimal Temperature




Decomposition of change in all-age, all sexes combined cardiovascular disease disability-adjusted life years (DALYs) attributable to non-optimal temperature from 1990 to 2023 due to population growth, population aging, risk exposure, and risk-deleted DALYs. Risk-deleted DALYs are the number of DALYs left after removing the effect of risk factors, population growth, and population aging on overall DALYs. They were calculated as the cardiovascular disease DALY count multiplied by 1 minus the population attributable fraction for non-optimal temperature. The dot and error bar represent the mean and 95% uncertainty interval in percentage change in number of DALYs attributable to the risk from 1990 to 2023.

ATRIAL FIBRILLATION. We found that the burden of atrial fibrillation increased due to increasing population age and size as well as to increasing high BMI and environmental lead exposure (through its effect on blood pressure), with small improvements in other risk factors offsetting these effects. Obesity is known to increase the risk of atrial fibrillation by multiple mechanisms, and weight reduction reduces atrial fibrillation burden.⁵⁵ Pooled studies of registries and clinical trials have shown that premature death among people with atrial fibrillation is due to a cardiovascular cause in more than half of cases, most commonly sudden cardiac death, heart failure, stroke, or myocardial infarction.⁵⁶ However, identification of atrial fibrillation as the underlying cause of death on a death certificate has become common only recently, and only in higher SDI locations, as

screening and diagnostic technologies become more widely available. Educational campaigns and quality improvement interventions appear to improve both detection of atrial fibrillation and delivery of anti-coagulation.^{57,58} Access to direct oral anticoagulants remains limited in many countries due to their cost and health care provider training, despite their addition to the WHO essential medicines list in 2019.⁵⁹

CVD DUE TO INFECTION. Infection is among the most morbid causes of CVD. Burden due to rheumatic heart disease is particularly high in Oceania, Sub-Saharan Africa, and South Asia. Rheumatic heart disease burden has declined globally but still affects lower-resource settings where ongoing surveillance shows that acute rheumatic fever and its carditis

FIGURE 30-1 Age-Standardized CVD DALY Rates Attributable to Lead Exposure, 2023

Age-standardized CVD DALYs attributable to lead exposure per 100,000 in 2023 (all sexes combined). Abbreviations as in Figure 20-1.

sequelae remain common.⁶⁰ A recent prospective cohort study of acute rheumatic fever, one of the first performed in Sub-Saharan Africa, showed that over 4.3 years of follow-up, 16% of patients had died.⁶¹ A record linkage study in Fiji found a death rate of 18.5% over 2.4 years.⁶² Bacterial endocarditis carries similarly high fatality rates. In a large European registry, 17.1% of patients with endocarditis died in hospital.⁶³ Large-scale efforts to better understand the epidemiology of these high-risk conditions has advanced in recent years, but most countries lack routine surveillance.


Viral infections are also common causes of CVD. The proportion of myocarditis that is due to viral infection remains unclear; however, SARS-CoV-2 may carry as much as a 15-fold increased risk for myocarditis.⁶⁴ Influenza has been estimated to be the cause of 4% of IHD deaths globally.⁶⁵

TRENDS IN CVD RISK FACTORS. More CVD is attributable to high SBP than any other risk factor. Exposure to high SBP on average did not change since 1990, and little improvement was seen over the last 2 decades outside of the highest-income countries.⁶⁶

The addition of single-pill combination medications for the prevention of ASCVD to the WHO List of Essential Medicines is particularly important given the dominant role of high blood pressure and cholesterol as modifiable risks for CVD.⁶⁷ For hypertension, switching to pills that include 2 or 3 anti-hypertensive medications has been shown safe and more effective at achieving blood pressure targets than traditional strategies. Delivery by community health workers who both screen and treat for hypertension allows these medications to be delivered even in the most resource-limited settings.⁶⁸

From 1990 to 2023, high FPG and high BMI contributed to the greatest proportional increase in attributable CVD DALYs out of all risk factors. This worrisome trend reflects the position of non-optimal diet as the second leading risk for CVD and the contribution of artificially formulated foods such as sugar-sweetened beverages.⁶⁹ Ultra-processed foods have been identified as a particular risk for health, driven substantially by consumption of sugar-sweetened beverages and processed meats.⁷⁰ No single score or scale adequately reflects a healthy diet, and increasing attention is being paid to dietary

FIGURE 30-2 Percentage Change in the Number of Global Risk-Attributable DALYs, 1990 to 2023, due to Population Growth, Population Aging, Changes in Exposures to Each Global Burden of Disease Risk Factor, and Changes in Risk-Deleted DALY Rates for All Sexes, for Lead Exposure



Decomposition of change in all-age, all sexes combined cardiovascular disease disability-adjusted life years (DALYs) attributable to lead exposure from 1990 to 2023 due to population growth, population aging, risk exposure, and risk-deleted DALYs. Risk-deleted DALYs are the number of DALYs left after removing the effect of risk factors, population growth, and population aging on overall DALYs. They were calculated as the cardiovascular disease DALY count multiplied by 1 minus the population attributable fraction for lead exposure. The dot and error bar represent the mean and 95% uncertainty interval in percentage change in number of DALYs attributable to the risk from 1990 to 2023.

patterns that combine healthy and unhealthy foods in complex ways.⁷¹

Among the most complex risks of CVD to analyze is non-optimal temperature. The analysis presented here makes use of a global surface temperature dataset and mortality data for 64.9 million deaths since 1990. GBD defines these risks as the same-day exposure to ambient temperature that is warmer or colder than the temperature with the minimum estimated mortality risk, specific to a given location in a given year, with a separate relative risk modeled for each cause of death.⁷² Non-optimal temperature is one of the few risks where our decomposition analysis shows decreasing attributable CVD burden, due to the higher risk of low temperature for IHD in the face of increasingly higher global temperatures. These estimates approximate a direct relationship

between surface temperature and specific CVD causes of death but do not include other pathways through which climate might have affected CVD, such as dehydration, migration, or extreme weather events.⁷³

Improvements in CVD risk factors at the global level can be seen in our analysis. Decreasing exposure to tobacco, high alcohol use, dietary risks including trans fats, and household air pollution led to reductions in age-standardized rates of CVD DALYs in many locations. These gains in health can be understood as the result of highly effective global and national health policies that need to be supported, continued and expanded.

POLICIES AND INTERVENTIONS TO ADDRESS CVD BURDEN. Advances in medical diagnosis and treatment of CVD remain limited in their availability for a

large proportion of people, contributing to large global disparities in CVD.⁷⁴ Approximately half of the improvement in CVD is likely to be due to medical therapies, with risk factor modification contributing equally to improved health.⁷⁵ The contributions of health systems to past reductions in CVD burden include prehospital and hospital care, revascularization and other device-based or surgical procedures, and secondary prevention therapies. The available health care workforce trained to diagnose and treat CVD and its risk factors remains limited.

A broad range of population-level policies have been adopted to improve CVD burden beginning with the North Karelia Project launched in Finland in 1972.⁷⁶ In the current era, tobacco prevention has been among the most effective policies for the reduction of CVD burden. The WHO Framework Convention on Tobacco Control facilitates taxation for tobacco products, inclusion of package warning labels, and restricted smoking in public spaces such as restaurants, worksites, and public transportation.⁷⁷ Major improvements in tobacco control have been observed over the past decade in locations with high burden such as found in parts of India, though current trends have flattened and additional progress is not clear in all locations.⁷⁸ In comparison, global efforts to reduce exposure to the harmful effects of alcohol have varied. In 2024, WHO launched updated guidelines to curb alcohol intake, including promoting restrictions in alcohol availability, raising excise taxes on alcoholic beverages, and restricting advertising.⁷⁹

Air pollution is the leading environmental risk factor contributing to CVD burden, but there is a large degree of geographic variation, with household and ambient air pollution affecting large areas of Africa and Asia. Installation of air pollution control technologies has been a successful strategy for reducing ambient pollution in some locations such as China.⁸⁰

Improvements in diet have been limited but promising, with increasing numbers of countries banning trans fats. The effect of Mexico's tax on sugar-sweetened beverages remains unclear.^{81,82} Japan and the United Kingdom have adopted broad, effective policies to reduce sodium consumption.⁸³ Front-of-package labeling is being adopted in multiple countries to provide consumers with information on nutritional value.⁸⁴

Many barriers exist to effective implementation of these policies, including extremely limited funding for the development and implementation of CVD policies due to the extremely small share of development assistance directed to non-communicable diseases in general and CVD in particular.⁸⁵

LIMITATIONS AND FUTURE EFFORTS IN GLOBAL SURVEILLANCE. Estimates from the GBD study were limited by the quality and availability of input data, particularly in data-sparse regions. Future investments in data collection would be particularly valuable in which regional trends and country-level covariates predict a large shift in the burden of CVD, but recent surveillance data are lacking. While covariate estimates are available for all countries, vital registration or verbal autopsy studies were not available for estimation of total CVD cause of death in 56 countries including 2 countries in Latin America and the Caribbean; 5 countries within North Africa and the Middle East; 2 countries in South Asia; 14 countries in Southeast Asia, East Asia, and Oceania; and 33 countries in Sub-Saharan Africa. We report country-level details on data availability for this analysis in [Supplemental Appendix 1 \(Supplemental Figure 21, Supplemental Table 12\)](#). GBD estimates are improved by the availability of risk factor data, which are more widely available than mortality data and can be used as covariates in disease models. For example, 50 of 56 countries with no cause-of-death data for total CVD had available input data to inform estimates of mean SBP. Rates of death due to hypertensive heart disease in Sub-Saharan Africa are a result of higher measured SBP in the health examination surveys in many of those countries. In the absence of direct data on CVD burden, robust statistical models using covariates generated estimates. Some risks and conditions have particularly limited amounts of input data and rely extensively on country-level covariates, including myocarditis and atrial fibrillation. Data on mediation of risk factors by other risks and how that may vary between populations remain limited. There are also risk factors not yet included in the GBD study, such as isolated elevation in diastolic blood pressure.

IMPLICATIONS FOR HEALTH SYSTEMS AND PRIORITY SETTING. Investment in cardiovascular health policies will be challenging in the current political and economic climate. In 2025, development assistance for health has fallen to its lowest level in over 15 years as multiple countries, including Finland, France, Germany, the United Kingdom, and the United States, have cut their funding.⁸⁶ Only a small proportion of this funding was ever invested in non-communicable diseases. This May, the 78th World Health Assembly approved a resolution affirming their commitment to universal health coverage and urging increases in domestic investments to improve financial protections and reduce out-of-pocket expenditures.⁸⁷

Estimates of cardiovascular risk and disease burden such as the those presented here offer the opportunity to develop and align global health policy with “facts on the ground” in this resource-constrained environment. Burden estimates function as an aggregator of information from millions of household survey respondents, patients enrolled in disease registries, anonymized hospital and ambulatory health system records, and death certificates filled out by the decedents reporting physicians. Taken together, this data suggests that, if current trends continue, the Sustainable Development Goals Target 3.4 of a one-third reduction in premature mortality from non-communicable diseases between 2015 and 2030 is not likely to be met.

To redirect these trends, future cardiovascular health policies will need to support risk factor reduction within a comprehensive framework for universal health coverage. The leading CVD risk factors are not the same in each country and local surveillance data and burden estimates can ensure an evidence-based approach that is responsive to local needs. In addition to the most common causes of CVD such as IHD, stroke, and HHD, health systems will need to invest in diagnostic and screening tools to address the rising burden of atrial fibrillation and non-rheumatic valvular heart disease, which are often asymptomatic early in their course. Policies should also support the eradication of rheumatic heart disease, which remains a persistent health threat in the poorest countries.⁸⁸

CONCLUSIONS

CVD remains the leading cause of disease burden and death worldwide (at level 2), with the greatest

burden in low, low-middle, and middle SDI regions. Large variation exists in CVD burden even for countries at similar levels of development, a gap explained substantially by known, modifiable risk factors that are inadequately controlled. Global levels of exposure to high BMI have grown steadily since 1990, along with high FPG and low physical activity to a lesser extent, which resulted in enlarged CVD burden. High SBP, the leading risk for CVD, remains a persistent threat for the health of most countries, while exposure to high LDL-C has not improved outside a small and select number of countries. In contrast, decreases in exposure to tobacco and air pollution, specifically household air pollution, have counteracted some of the rising CVD burden. Ambient air pollution and environmental lead exposure remain serious health threats. Effective health systems and public health strategies exist to reduce these risks. Countries will need to adopt these strategies if they are to progress in achieving global goals to reduce the burden of CVD.

FUNDING SUPPORT AND AUTHOR DISCLOSURES

Funding was provided by the Bill and Melinda Gates Foundation and the American College of Cardiology Foundation. The contents and views expressed in this report are those of the authors and do not necessarily reflect the official views of the National Institutes of Health, the Department of Health and Human Services, the U.S. government, or the affiliated institutions. The authors have reported that they have no relationships relevant to the contents of this paper to disclose.

ADDRESS FOR CORRESPONDENCE: Dr Gregory A. Roth, Institute for Health Metrics and Evaluation, 3980 15th Avenue NE, UW Campus Box #351615, Seattle, Washington 98195, USA. Email: rothg@uw.edu

REFERENCES

1. World Health Organization. SDG Target 3.4 Noncommunicable diseases and mental health. Accessed January 3, 2025. <https://www.who.int/data/gho/data/themes/topics/sdg-target-3.4-noncommunicable-diseases-and-mental-health>
2. Stevens GA, Alkema L, Black RE, et al. Guidelines for Accurate and Transparent Health Estimates Reporting: the GATHER statement. *Lancet*. 2016;388:e19-e23.
3. Johnson SC, Cunningham M, Dippenaar IN, et al. Public health utility of cause of death data: applying empirical algorithms to improve data quality. *BMC Med Inform Decis Mak*. 2021;21:175. <https://doi.org/10.1186/s12911-021-01501-1>
4. Rwebembera J, Marangou J, Mwita JC, et al. 2023 World Heart Federation guidelines for the echocardiographic diagnosis of rheumatic heart disease. *Nat Rev Cardiol*. 2024;21:250-263.
5. Thygesen K, Alpert JS, Jaffe AS, et al. Fourth Universal Definition of Myocardial Infarction (2018). *Circulation*. 2018;138:e618-e651.
6. Jones WB, Riley CP, Reeves TJ, Sheffield LT. Natural history of coronary artery disease. *Bull N Y Acad Med*. 1972;48:1109-1125.
7. Raff GL, Abidov A, Achenbach S, et al. SCCT guidelines for the interpretation and reporting of coronary computed tomographic angiography. *J Cardiovasc Comput Tomogr*. 2009;3:122-136.
8. Bozkurt B, Coats AJS, Tsutsui H, et al. Universal definition and classification of heart failure: a report of the Heart Failure Society of America, Heart Failure Association of the European Society of Cardiology, Japanese Heart Failure Society and Writing Committee of the Universal Definition of Heart Failure: Endorsed by the Canadian Heart Failure Society, Heart Failure Association of India, Cardiac Society of Australia and New Zealand, and Chinese Heart Failure Association. *Eur J Heart Fail*. 2021;23:352-380.
9. Donkor ES. Stroke in the 21st century: a snapshot of the burden, epidemiology, and quality of life. *Stroke Res Treat*. 2018;2018:3238165.
10. Aho K, Harmsen P, Hatano S, Marquardsen J, Smirnov VE, Strasser T. Cerebrovascular disease in the community: results of a WHO collaborative study. *Bull World Health Organ*. 1980;58:113-130.
11. Gibson G, Blumer V, Mentz RJ, Lala A. Universal Definition and Classification of Heart Failure: a step in the right direction from failure to function. Accessed January 27, 2025. <https://www.acc.org/Latest-in-Cardiology/Articles/2021/07/12/12/31/Universal-Definition-and-Classification-of-Heart-Failure>

12. Otto CM, Nishimura RA, Bonow RO, et al. 2020 ACC/AHA guideline for the management of patients with valvular heart disease: executive summary: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol.* 2021;77(4):450-500.
13. Ammirati E, Frigerio M, Adler ED, et al. Management of acute myocarditis and chronic inflammatory cardiomyopathy: an expert consensus document. *Circ Heart Fail.* 2020;13:e007405.
14. Galiè N, Humbert M, Vachiery J-L, et al. 2015 ESC/ERS guidelines for the diagnosis and treatment of pulmonary hypertension: the Joint Task Force for the Diagnosis and Treatment of Pulmonary Hypertension of the European Society of Cardiology (ESC) and the European Respiratory Society (ERS): Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC), International Society for Heart and Lung Transplantation (ISHLT). *Eur Heart J.* 2016;37:67-119.
15. Iwasaki Y, Nishida K, Kato T, Nattel S. Atrial fibrillation pathophysiology: implications for management. *Circulation.* 2011;124:2264-2274.
16. Saleh K, Haldar S. Atrial fibrillation: a contemporary update. *Clin Med.* 2023;23:437-441.
17. Isselbacher EM, Preventza O, Hamilton Black J, et al. 2022 ACC/AHA guideline for the diagnosis and management of aortic disease: a report of the American Heart Association/American College of Cardiology Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol.* 2022;80:e223-e393.
18. Criqui MH, Matsushita K, Aboyans V, et al. Lower extremity peripheral artery disease: contemporary epidemiology, management gaps, and future directions: a Scientific Statement From the American Heart Association. *Circulation.* 2021;144:e171-e191.
19. Fowler VG Jr, Durack DT, Selson-Suty C, et al. The 2023 Duke-International Society for Cardiovascular Infectious Diseases Criteria for Infective Endocarditis: Updating the Modified Duke Criteria. *Clin Infect Dis.* 2023;77:518-526.
20. Naghavi M, Ong KL, Aali A, et al. Global burden of 288 causes of death and life expectancy decomposition in 204 countries and territories and 811 subnational locations, 1990-2021: a systematic analysis for the Global Burden of Disease Study 2021. *Lancet.* 2024;403:2100-2132.
21. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021;372:n71.
22. Stark B, Roth G, Johnson C, DeCleene N. Systematic review of the global burden of ischemic heart disease. 2022. Accessed July 29, 2025. https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42022323863
23. Lindstrom M, Roth G, Johnson C, DeCleene N, Minja N. Systematic review of the global burden of rheumatic heart disease. 2022. Accessed July 29, 2025. https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD4202235770
24. Stark B, Roth G, Johnson C, DeCleene N. Systematic review of the global burden of myocardial infarction. 2023. Accessed July 29, 2025. https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42023414474
25. DeCleene N, Roth G, Johnson C, Lindstrom M. Systematic review of global systolic blood pressure exposure from 2017 to 2022. 2024. Accessed July 29, 2025. https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42024510285
26. Lindstrom M, Roth G, Johnson C, Stark B, DeCleene N. Systematic review of the global burden of peripheral arterial disease from 2015 to 2023. 2024. Accessed July 29, 2025. https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42024510900
27. Zheng P, Barber R, Sorensen RJD, Murray CJL, Aravkin AY. Trimmed constrained mixed effects models: formulations and algorithms. *J Comput Graph Stat.* 2021;30:544-556.
28. Ferrari AJ, Santomauro DF, Aali A, et al. Global incidence, prevalence, years lived with disability (YLDs), disability-adjusted life-years (DALYs), and healthy life expectancy (HALE) for 371 diseases and injuries in 204 countries and territories and 811 subnational locations, 1990-2021: a systematic analysis for the Global Burden of Disease Study 2021. *Lancet.* 2024;403:2133-2161.
29. Zheng P, Afshin A, Biryukov S, et al. The Burden of Proof studies: assessing the evidence of risk. *Nat Med.* 2022;28:2038-2044.
30. Brauer M, Roth GA, Aravkin AY, et al. Global burden and strength of evidence for 88 risk factors in 204 countries and 811 subnational locations, 1990-2021: a systematic analysis for the Global Burden of Disease Study 2021. *Lancet.* 2024;403:2162-2203.
31. Schumacher AE, Kyu HH, Aali A, et al. Global age-sex-specific mortality, life expectancy, and population estimates in 204 countries and territories and 811 subnational locations, 1950-2021, and the impact of the COVID-19 pandemic: a comprehensive demographic analysis for the Global Burden of Disease Study 2021. *Lancet.* 2024;403:1989-2056.
32. Das Gupta P. *Standardization and Decomposition of Rates: A User's Manual.* United States Census Bureau; 1993. Accessed January 3, 2025. <https://www.census.gov/library/publications/1993/demo/p23-186.html>
33. Moran AE, Farrell M, Cazabon D, et al. Building the health-economic case for scaling up the WHO-HEARTS hypertension control package in low- and middle-income countries. *Rev Panam Salud Pub.* 2022;46:1.
34. Lanphear BP, Rauch S, Auinger P, Allen RW, Hornung RW. Low-level lead exposure and mortality in US adults: a population-based cohort study. *Lancet Public Health.* 2018;3(4):e177-e184.
35. Zhao D, Wang Y, Wong ND, Wang J. Impact of aging on cardiovascular diseases: from chronological observation to biological insights: JACC Family Series. *JACC Asia.* 2024;4:345-358.
36. Nielsen RV, Fuster V, Bundgaard H, et al. Personalized intervention based on early detection of atherosclerosis: JACC State-of-the-Art Review. *J Am Coll Cardiol.* 2024;83:2112-2127.
37. Arnett DK, Blumenthal RS, Albert MA, et al. 2019 ACC/AHA guideline on the primary prevention of cardiovascular disease: executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol.* 2019;74(10):1376-1414.
38. Ogurtsova K, Guariguata L, Barendo NC, et al. IDF diabetes Atlas: Global estimates of undiagnosed diabetes in adults for 2021. *Diabetes Res Clin Pract.* 2022;183:109118.
39. Jan S, Laba TL, Essue BM, et al. Action to address the household economic burden of non-communicable diseases. *Lancet.* 2018;391(10134):2047-2058.
40. Countdown to 2023. *WHO 5-Year Milestone Report on Global Trans Fat Elimination 2023.* 1st ed. World Health Organization; 2024.
41. WHO Global Air Quality Guidelines Particulate matter (PM2.5 and PM10), Ozone, Nitrogen Dioxide, Sulfur Dioxide and Carbon Monoxide. World Health Organization; 2021.
42. Lee K, Egbe CO, Bianco E, Arora M. The 20th anniversary of the WHO Framework Convention on Tobacco Control: hard won progress amid evolving challenges. *Lancet.* 2023;402(10402):592-594.
43. Prust ML, Forman R, Ovbiagele B. Addressing disparities in the global epidemiology of stroke. *Nat Rev Neurol.* 2024;20:207-221.
44. Owolabi MO, Thrift AG, Martins S, et al. The state of stroke services across the globe: report of World Stroke Organization-World Health Organization surveys. *Int J Stroke.* 2021;16:889-901.
45. Feigin VL, Owolabi MO, Feigin VL, et al. Pragmatic solutions to reduce the global burden of stroke: a World Stroke Organization-Lancet Neurology Commission. *Lancet Neurol.* 2023;22:1160-1206.
46. Sebastian IA, Gandhi DBC, Sylaja PN, et al. Stroke systems of care in South-East Asia Region (SEAR): commonalities and diversities. *Lancet Region Health Southeast Asia.* 2023;17:100289.
47. Bernhardt J, Urimubenshi G, Gandhi DBC, Eng JJ. Stroke rehabilitation in low-income and middle-income countries: a call to action. *Lancet.* 2020;396(10260):1452-1462.
48. Bhattacharjee NV, Schumacher AE, Aali A, et al. Global fertility in 204 countries and territories, 1950-2021, with forecasts to 2100: a comprehensive demographic analysis for the Global Burden of Disease Study 2021. *Lancet.* 2024;403(10440):2057-2099.
49. Agarwal A, Tromp J, Almahmeed W, et al. Toward a universal definition of etiologies in heart failure: categorizing causes and advancing registry science. *Circ Heart Fail.* 2024;17:e011095.
50. Attaei MW, Khatib R, McKee M, et al. Availability and affordability of blood pressure-lowering medicines and the effect on blood pressure control in high-income, middle-income, and low-income countries: an analysis of the PURE study data. *Lancet Public Health.* 2017;2:e411-e419.

51. Rodgers A, Salam A, Schutte AE, et al. Efficacy and safety of a novel low-dose triple single-pill combination of telmisartan, amlodipine and indapamide, compared with dual combinations for treatment of hypertension: a randomised, double-blind, active-controlled, international clinical trial. *Lancet*. 2024;404(10462):1536-1546.
52. Domínguez F, Adler E, García-Pavía P. Alcoholic cardiomyopathy: an update. *Eur Heart J*. 2024;45:2294-2305.
53. World Health Organization. Global status report on alcohol and health and treatment of substance use disorders. Accessed January 24, 2025. <https://www.who.int/publications/i/item/9789240096745>
54. Yadgir S, Johnson CO, Aboyans V, et al. Global, regional, and national burden of calcific aortic valve and degenerative mitral valve diseases, 1990-2017. *Circulation*. 2020;141(21):1670-1680.
55. McCauley MD, Iacobellis G, Li N, Nattel S, Goldberger JJ. Targeting the substrate for atrial fibrillation. *J Am Coll Cardiol*. 2024;83(20):2015-2027.
56. Gómez-Outes A, Suárez-Gea ML, García-Pinilla JM. Causes of death in atrial fibrillation: challenges and opportunities. *Trends Cardiovasc Med*. 2017;27:494-503.
57. Whitfield R, Ascenção R, da Silva GL, Almeida AG, Pinto FJ, Caldeira D. Screening strategies for atrial fibrillation in the elderly population: a systematic review and network meta-analysis. *Clin Res Cardiol*. 2023;112(6):705-715.
58. Suzuki A, Okamura T, Sasaki M, et al. Acceleration of opportunistic atrial fibrillation screening for elderly patients in routine primary care. *PLoS One*. 2020;15(12):e0244240.
59. Noubiap JJ, Kamtchum-Tatuene J. Addition of direct oral anticoagulants to the World Health Organization model list of essential medicines for the treatment of atrial fibrillation: An African perspective. *Br J Clin Pharmacol*. 2022;88(7):3035-3038.
60. Thandrayen J, Stacey I, Oliver J, Francia C, Katzenellenbogen JM, Wyber R. Estimating the true number of people with acute rheumatic fever and rheumatic heart disease from two data sources using capture-recapture methodology. *Aust Health Review*. 2024;49(1):AH24267.
61. Wirth SH, Pulle J, Seo J, et al. Outcomes of rheumatic fever in Uganda: a prospective cohort study. *Lancet Glob Health*. 2024;12:e500-e508.
62. Parks T, Narube L, Perman ML, et al. Population-based assessment of cardiovascular complications of rheumatic heart disease in Fiji: a record-linkage analysis. *BMJ Open*. 2023;13:e070629.
63. Habib G, Erba PA, lung B, et al. Clinical presentation, aetiology and outcome of infective endocarditis. Results of the ESC-EORP EURO-ENDO (European infective endocarditis) registry: a prospective cohort study. *Eur Heart J*. 2019;40:3222-3232.
64. Fairweather D, Beetler DJ, Di Florio DN, Musigk N, Heidecker B, Cooper LT. COVID-19, Myocarditis and Pericarditis. *Circ Res*. 2023;132:1302-1319.
65. Chaves SS, Nealon J, Burkart KG, et al. Global, regional and national estimates of influenza-attributable ischemic heart disease mortality. *EClinicalMedicine*. 2023;55:101740.
66. Moran AE, Gupta R. Implementation of global hearts hypertension control programs in 32 low- and middle-income countries. *J Am Coll Cardiol*. 2023;82(19):1868-1884.
67. Agarwal A, Huffman MD. Inclusion of polyphills for prevention of cardiovascular disease in the 23rd World Health Organization model list of essential medicines: a significant step towards reducing global cardiovascular morbidity and mortality. *Glob Heart*. 2024;19:24.
68. Tian M, Ajay VS, Dunzhu D, et al. A cluster-randomized, controlled trial of a simplified multifaceted management program for individuals at high cardiovascular risk (SimCard Trial) in rural Tibet, China, and Haryana, India. *Circulation*. 2015;132(9):815-824.
69. Afshin A, Sur PJ, Fay KA, et al. Health effects of dietary risks in 195 countries, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2019;393(10184):1958-1972.
70. Mendoza K, Smith-Warner SA, Rossato SL, et al. Ultra-processed foods and cardiovascular disease: analysis of three large US prospective cohorts and a systematic review and meta-analysis of prospective cohort studies. *Lancet Reg Health Am*. 2024;37:100859.
71. Hanley-Cook GT, Gie SM, Parraguez JP, et al. Cross-context equivalence and agreement of healthy diet metrics for national and global monitoring: a multicountry analysis of cross-sectional quantitative 24-hour dietary intake studies. *Am J Clin Nutr*. 2024;120:1093-1104.
72. Burkart KG, Brauer M, Aravkin AY, et al. Estimating the cause-specific relative risks of nonoptimal temperature on daily mortality: a two-part modelling approach applied to the Global Burden of Disease Study. *Lancet*. 2021;398(10301):685-697.
73. Burden of disease scenarios for 204 countries and territories, 2022-2050: a forecasting analysis for the Global Burden of Disease Study 2021. *Lancet*. 2024;403(10440):2204-2256.
74. Leong DP, Joseph PG, McKee M, et al. Reducing the Global Burden of Cardiovascular Disease, part 2: prevention and treatment of cardiovascular disease. *Circ Res*. 2017;121:695-710.
75. Ford ES, Ajani UA, Croft JB, et al. Explaining the decrease in U.S. deaths from coronary disease, 1980-2000. *N Engl J Med*. 2007;356(23):2388-2398.
76. Vartiainen E. The North Karelia Project: cardiovascular disease prevention in Finland. *Glob Cardiol Sci Pract*. 2018;2018(2):13.
77. World Health Organization. WHO report on the global tobacco epidemic, 2023: protect people from tobacco smoke. Accessed January 24, 2025. <https://www.who.int/publications/i/item/9789240077164>
78. Saxena A, Sharma K, Avashia V. Assessment of tobacco control policy in India: an evaluation using the World Health Organization MPOWER Framework. *Indian J Community Med*. 2020;45:543-545.
79. World Health Organization. Global alcohol action plan 2022-2030. Accessed January 24, 2025. <https://www.who.int/publications/i/item/9789240090101>
80. Geng G, Zheng Y, Zhang Q, et al. Drivers of PM2.5 air pollution deaths in China 2002-2017. *Nat Geosci*. 2021;14(9):645-650.
81. Aguilar A, Gutierrez E, Seira E. The effectiveness of sin food taxes: Evidence from Mexico. *J Health Econ*. 2021;77:102455.
82. Colchero MA, Molina M, Guerrero-López CM. After Mexico implemented a tax, purchases of sugar-sweetened beverages decreased and water increased: difference by place of residence, household composition, and income level. *J Nutr*. 2017;147(8):1552-1557.
83. Ikeda N, Yamaguchi M, Kashino I, Sugiyama T, Miura K, Nishi N. Evaluation of public health and economic impacts of dietary salt reduction initiatives on social security expenditures for cardiovascular disease control in Japan. *Hypertens Res*. 2025;48(4):1265-1273.
84. Batista MF, de Carvalho-Ferreira JP, Thimoteo da Cunha D, De Rosso VV. Front-of-package nutrition labeling as a driver for healthier food choices: lessons learned and future perspectives. *Compr Rev Food Sci Food Saf*. 2023;22(1):535-586.
85. Apeagyei AE, Lidral-Porter B, Patel N, et al. Financing health in Sub-Saharan Africa 1990-2050: donor dependence and expected domestic health spending. *PLoS Global Public Health*. 2024;4(8):e0003433.
86. Institute for Health Metrics and Evaluation. Financing Global Health 2025: Cuts in Aid and Future Outlook. Institute for Health Metrics and Evaluation. Accessed July 29, 2025. <https://www.healthdata.org/research-analysis/library/financing-global-health-2025-cuts-aid-and-future-outlook>.
87. WHO Executive Board, 156, 2025. Executive Board, 156th session, Geneva, 10 February 2025: Strengthening health financing globally. Accessed July 29, 2025. https://apps.who.int/gb/ebwha/pdf_files/EB156/B156_16-en.pdf
88. Lindstrom M, Minja NW, Stark B, et al. A systematic methodology to capture the global pattern of rheumatic heart disease: the Rheumatic Heart Disease Endemicity Index (RHDEI). *BMC Glob Public Health*. 2025;3(1):62.

KEY WORDS cardiovascular disease, epidemiology, global health

APPENDIX For supplemental information, please see the online version of this paper.