

Hippocampal CB₁ Receptors Control Incidental Associations

Highlights

- Hippocampal CB₁R is required for incidental associations leading to mediated learning
- Incidental associations enhance hippocampal CB₁R expression and inhibitory plasticity
- CB₁R-dependent control of hippocampal GABAergic cells controls mediated learning

Authors

Arnaud Busquets-Garcia,
José F. Oliveira da Cruz,
Geoffrey Terral, ..., Pierre Trifilieff,
Guillaume Ferreira,
Giovanni Marsicano

Correspondence

guillaume.ferreira@inra.fr (G.F.),
giovanni.marsicano@inserm.fr (G.M.)

In Brief

Busquets-Garcia et al. suggest that cannabinoid CB₁ receptors signaling in the hippocampus, a brain structure involved in memory processes, underlines the mammalian ability to associate randomly encountered stimuli, allowing future inferred memories and possibly explaining seemingly ungrounded responses toward certain cues.



Hippocampal CB₁ Receptors Control Incidental Associations

Arnau Busquets-Garcia,^{1,2} José F. Oliveira da Cruz,^{1,2,3,8} Geoffrey Terral,^{1,2,8} Antonio C. Pagano Zottola,^{1,2,8} Edgar Soria-Gómez,^{1,2,4,5} Andrea Contini,^{2,6} Hugo Martin,^{2,6} Bastien Redon,^{1,2} Marjorie Varilh,^{1,2} Christina Ioannidou,^{1,2} Filippo Drago,³ Federico Massa,^{1,2} Xavier Fioramonti,^{2,6} Pierre Trifilieff,^{2,6} Guillaume Ferreira,^{2,6,7,*} and Giovanni Marsicano^{1,2,7,9,*}

¹INSERM, U1215 NeuroCentre Magendie, 33000 Bordeaux, France

²University of Bordeaux, 33000 Bordeaux, France

³Department of Biomedical and Biotechnological Sciences, Section of Pharmacology, University of Catania, I-95123 Catania, Italy

⁴Department of Neurosciences, University of the Basque Country UPV/EHU, Achucarro Basque Center for Neuroscience, 48940 Leioa, Spain

⁵IKERBASQUE, Basque Foundation for Science, 48013 Bilbao, Spain

⁶INRA, Nutrition and Integrative Neurobiology, UMR 1286, Bordeaux, France

⁷Senior author

⁸These authors contributed equally

⁹Lead Contact

*Correspondence: guillaume.ferreira@inra.fr (G.F.), giovanni.marsicano@inserm.fr (G.M.)

<https://doi.org/10.1016/j.neuron.2018.08.014>

SUMMARY

By priming brain circuits, associations between low-salience stimuli often guide future behavioral choices through a process known as mediated or inferred learning. However, the precise neurobiological mechanisms of these incidental associations are largely unknown. Using sensory preconditioning procedures, we show that type 1 cannabinoid receptors (CB₁R) in hippocampal GABAergic neurons are necessary and sufficient for mediated but not direct learning. Deletion and re-expression of CB₁R in hippocampal GABAergic neurons abolishes and rescues mediated learning, respectively. Interestingly, paired presentations of low-salience sensory cues induce a specific protein synthesis-dependent enhancement of hippocampal CB₁R expression and facilitate long-term synaptic plasticity at inhibitory synapses. CB₁R blockade or chemogenetic manipulations of hippocampal GABAergic neurons upon preconditioning affect incidental associations, as revealed by impaired mediated learning. Thus, CB₁R-dependent control of inhibitory hippocampal neurotransmission mediates incidental associations, allowing future associative inference, a fundamental process for everyday life, which is altered in major neuropsychiatric diseases.

INTRODUCTION

Direct associative memories, in which a sensory stimulus is explicitly paired with a negative or rewarding outcome, can determine daily behavioral choices. Very often, however, human behavior is governed by mediated learning, based on previous

events implying incidental associations between low-salience sensory cues (Bornstein et al., 2017; Shohamy and Wagner, 2008; Wimmer and Shohamy, 2012). In other words, we are often repulsed or attracted by stimuli never explicitly paired with negative or positive outcomes but previously associated with other stimuli paired with a specific aversive or rewarding meaning (Bornstein et al., 2017; Shohamy and Wagner, 2008; Wimmer and Shohamy, 2012). These processes are well conserved in all mammals, including rodents (Gewirtz and Davis, 2000; Parkes and Westbrook, 2011). However, whereas the biological mechanisms underlying direct associative learning are under intense scrutiny (LeDoux, 2014), much less is known about the neural substrates mediating sensory stimulus-stimulus associations leading to higher-order mediated learning (Gewirtz and Davis, 2000; Parkes and Westbrook, 2011).

Sensory preconditioning is a typical behavioral procedure to study mediated learning (Gewirtz and Davis, 2000; Parkes and Westbrook, 2011). In this protocol, pairings of two low-salience stimuli (e.g., odors, tastes, lights, tones) are followed by classical conditioning of one of these stimuli with an aversive or appetitive unconditioned reinforcer (Parkes and Westbrook, 2011). As a result of these associations, subjects present aversion or preference to the stimulus never explicitly paired with the reinforcer, thereby allowing the evaluation of mediated learning (Gewirtz and Davis, 2000; Parkes and Westbrook, 2011; Wheeler et al., 2013). Thus, three distinct and temporally successive processes occur in sensory preconditioning. First, an incidental association is formed between low-salience stimuli during the preconditioning phase. Second, direct association with a reinforcer increases the salience of one of the stimuli during the conditioning phase. Finally, exposing the subjects to either of the original stimuli (the one directly associated with the reinforcer and the one never associated) reveals the retrieval of direct and mediated memories, respectively. The neurobiological mechanisms of these connected but distinct processes are poorly understood. The hippocampus has been suggested to play an important role in the conditioning and






Figure 1. CB₁R Are Necessary for Odor-Taste-Mediated Learning

(A) Schematic table of the odor-taste sensory preconditioning protocol (see [Star Methods](#) and [Figure S1](#) for more details).

(B and C) Liquid consumption under conditions of mediated taste (B) or odor (C) aversion in CB₁R-KO mice and wild-type littermates (CB₁R-WT).

(D and E) Liquid consumption under conditions of direct odor (D) or taste (E) aversion in CB₁R-KO mice and CB₁R-WT. **p* < 0.05; ****p* < 0.001 (mCS+ versus mCS- or CS+ versus CS-). For statistical details and *n*, see [Tables S1](#) and [S2](#).

retrieval phases of sensory preconditioning procedures ([Iordanova et al., 2011](#); [Talk et al., 2002](#); [Wheeler et al., 2013](#); [Wimmer and Shohamy, 2012](#)). However, its involvement in the formation of initial incidental associations leading to mediated learning is unknown.

Type 1 cannabinoid receptors (CB₁R) are key neuromodulatory elements of synapses, and they are the main targets of endogenous signaling molecules, the endocannabinoids, forming the so-called endocannabinoid system (ECS) ([Lu and Mackie, 2016](#); [Piomelli, 2003](#)). Through CB₁R, the ECS has been involved in direct conditioning such as fear conditioning ([Marsicano et al., 2002](#); [Metna-Laurent et al., 2012](#); [Resstel et al., 2009](#)) or conditioned taste aversion ([Kobilo et al., 2007](#)). Notably, the involvement of the ECS in direct conditioning appears to be more prominent in the modulation of behavioral expression of the acquired memory, rather than its formation ([Kobilo et al., 2007](#); [Marsicano et al., 2002](#); [Metna-Laurent et al., 2012](#); [Resstel et al., 2009](#)). Regarding hippocampus, CB₁R are mainly expressed in GABAergic neurons ([Marsicano and Kuner, 2008](#); [Marsicano and Lutz, 1999](#)), where they negatively control inhibitory neurotransmission ([Katona et al., 1999](#)), thereby modulating synaptic plasticity ([Araque et al., 2017](#); [Busquets-Garcia et al., 2018](#); [Castillo et al., 2012](#); [Kano et al., 2009](#)) and cognitive processes ([Busquets-Garcia et al., 2015](#)). Although hippocampal CB₁R have been implicated in the cognitive impairment produced by exogenous cannabinoids ([Busquets-Garcia et al., 2015](#); [Puighermanal et al., 2009](#)), no study has addressed the physiological role of the ECS in higher-order learning such as sensory preconditioning.

Thus, applying genetic, pharmacological, biochemical, imaging, electrophysiological, and chemogenetic approaches to sensory preconditioning procedures in mice, the present study shows how the physiological inhibition of specific hippocampal GABAergic neuronal populations by CB₁R is crucial for inci-

denal associations between low-salience stimuli, eventually leading to mediated learning.

RESULTS

CB₁R Are Necessary for Mediated Learning

Mice were exposed to a preconditioning phase with three odor-taste pairings (sucrose and maltodextrin as tastes; banana and almond as odors), followed by a conditioning phase consisting in the devaluation of one of the two stimuli ([Figures 1A](#), [S1A](#), and [S1B](#)). First, we controlled that the potential salience of taste stimuli ([Yiannakas and Rosenblum, 2017](#)) was not sufficient to induce any type of observable direct conditioning in our experimental protocol ([Figures S1C](#) and [S1D](#)). Then we observed that wild-type mice consumed lower amounts of tastes or odors that were indirectly (preconditioned) or directly devaluated, indicating the formation of reliable mediated and direct aversion learning, respectively ([Figures 1B–1E](#) and [S1E–S1H](#)). Interestingly, global CB₁R knockout mice (CB₁R-KO) displayed impaired mediated aversion independently of the sensory modality ([Figures 1B](#), [1C](#), [S1E](#), and [S1F](#)), still maintaining normal expression of direct learning ([Figures 1D](#), [1E](#), [S1G](#), and [S1H](#)), thereby demonstrating that CB₁R are specifically required for mediated learning.

Activation of CB₁R during Preconditioning Is Necessary for Mediated Learning

Long-lasting absence of CB₁R in CB₁R-KO mice might induce chronic alterations, eventually causing the observed phenotype in mediated learning. To test whether CB₁R are acutely required during incidental associations, we administered the CB₁R antagonist Rimonabant (1 mg/kg, i.p.) before each session of preconditioning ([Figures 2A](#), [S1A](#), and [S1B](#)). This treatment impaired mediated learning, both in the taste and odor modalities, without affecting either direct aversions ([Figures 2B–2E](#) and [S2A–S2D](#)). Importantly, Rimonabant treatment did not alter the total liquid consumption during preconditioning ([Figure S2E](#)), nor did it affect the lack of taste-induced odor conditioning during odor-taste pairings ([Figures S2F–S2H](#)). Importantly, Rimonabant administration immediately before the test did not affect mediated odor aversion ([Figures 2F](#), [2G](#), [S2I](#), and [S2J](#)) or mediated




Figure 2. Activation of CB₁R during Preconditioning Is Necessary for Mediated Learning

(A) Schematic table of the odor-taste sensory preconditioning protocol (see [Star Methods](#) and [Figure S1](#) for more details).

(B and C) Effect of daily preconditioning administrations of the CB₁R antagonist Rimonabant (1 mg/kg, i.p.) and its vehicle on mediated taste (B) and direct odor aversion (C) in C57BL/6-N mice.

(D and E) Effect of daily preconditioning administrations of Rimonabant (1 mg/kg, i.p.) and its vehicle on mediated odor (D) and direct taste aversion (E) in C57BL/6-N mice.

(F and G) Effect of acute administration of Rimonabant (1 mg/kg, i.p.) and its vehicle before testing on mediated odor (F) and direct taste aversion (G) in C57BL/6-N mice.

(H) Schematic representation of the light-tone sensory preconditioning protocol (see [Star Methods](#) for more details).

(I and J) Effect of daily preconditioning administrations of Rimonabant (1 mg/kg, i.p.) and its vehicle on mediated tone learning, expressed as discrimination ratio, and (J) on the conditioning to light in C57BL/6-J mice (see [Star Methods](#) for further details). **p* < 0.05; ***p* < 0.01; ****p* < 0.001 (mCS+ versus mCS- or CS+ versus CS-). ****p* < 0.001 (vehicle- versus Rimonabant-treated mice). For statistical details and *n*, see [Tables S1](#) and [S2](#).

taste aversion (Busquets-Garcia et al., 2017b), suggesting that CB₁R activation is specifically required for the formation of incidental associations, but not for the expression of mediated aversion.

Next, we evaluated whether CB₁R activation was required also in another sensory preconditioning protocol using different sensory stimuli (visual and auditory cues) and unconditioned stimulus (food reward). To this aim, we adapted an existing paradigm originally developed in rats using operant chambers and visual/auditory cues (Robinson et al., 2014; Figure 2H). Mice treated with vehicle or Rimonabant (1 mg/kg, i.p.) were exposed to two preconditioning sessions consisting in pairings between a light and a tone (preconditioned cues), and to presentations of a click alone (unpaired cue). Next, the light was paired with food delivery and the visits of the food magazine in response to tone and click presentations were evaluated in the absence of the food reward (test session). Vehicle-treated mice displayed mediated conditioning as shown by a discrimination index indicating the increased visits of the food magazine during tone presentations as compared to the click (Figure 2I). Interestingly, mediated learning was specifically prevented by the Rimonabant treatment during the preconditioning phase (Figure 2I), whereas both vehicle- and Rimonabant-treated mice exhibited comparable levels of direct conditioning to the light (Figures 2J and S2K). Altogether, these results indicate that mediated learning relies on CB₁R activation specifically during the formation of incidental associations, regardless of the sensory modalities used and of the nature (aversive or appetitive) of the reinforcers.

Hippocampal CB₁R Are Required for Mediated Learning

Inactivation and functional imaging experiments have suggested the involvement of the hippocampus in mediated learning (Iordanova et al., 2011; Wheeler et al., 2013; Wimmer and Shohamy, 2012), but the role of hippocampal CB₁R in these processes have never been investigated. To address this issue, we injected adeno-associated viruses (AAVs) expressing Cre recombinase or control green fluorescent protein (GFP) into the hippocampus of CB₁R-flox mice (Marsicano et al., 2003) to generate mice lacking CB₁R in the hippocampus and control littermates with normal expression of the receptor (called HC-CB₁R-KO and HC-CB₁R-WT, respectively; Monory et al., 2006; Figures 3A and S3A). HC-CB₁R-KO mice displayed an impairment of mediated, but not direct, aversion in the odor-taste protocol, regardless of the sensory modality (Figures 3B, 3C, and S3B–S3E). The lack of mediated learning after three odor-taste pairings was not due to slower incidental learning as mice undergoing extended preconditioning using six odor-taste pairings (Busquets-Garcia et al., 2017b) did not show mediated aversion (Figures S3F and S3G). We next investigated whether hippocampal CB₁R are sufficient to allow mediated learning. Notably, the viral re-expression of CB₁R in the hippocampus of CB₁R-KO mice (HC-CB₁R-RS mice; Figures 3D and S3A; Hebert-Chatelain et al., 2016) fully rescued mediated learning (Figures 3E, S3H, and S3I). Altogether, these data show that hippocampal CB₁R are necessary and sufficient for mediated learning.

GABAergic CB₁R Are Necessary and Sufficient for Mediated Learning

The large majority of hippocampal CB₁R are present in a specific subpopulation of GABAergic interneurons, namely the cholecystokinin (CCK)-expressing perisomatic basket cells (Kano et al., 2009; Marsicano and Kuner, 2008). To study the role of this specific population of CB₁R, we first used conditional mutant mice lacking the receptor from forebrain GABAergic neurons (Dlx5/6-CB₁R-KO mice, generally and hereafter called GABA-CB₁R-KO; Bellocchio et al., 2010; Monory et al., 2006). Notably, these animals displayed a specific impairment of mediated aversion independently of the sensory modality (Figures 4A, 4B, S4A, and S4B), accompanied by normal direct aversion (Figures 4C, 4D, S4C, and S4D). This impairment might be due to a delayed formation of mediated learning. However, when exposed to an extended preconditioning training (six odor-taste pairings; Busquets-Garcia et al., 2017b), GABA-CB₁R-KO mice were still unable to display mediated aversion (Figures S4E and S4F), strongly suggesting that “GABAergic” CB₁R are necessary for the formation of incidental learning independently of the training intensity. In addition, “rescue” mice carrying expression of the CB₁R protein exclusively in forebrain GABAergic neurons (GABA-CB₁R-RS) (Gutiérrez-Rodríguez et al., 2017; Remmers et al., 2017) displayed normal mediated aversion, both in the taste or odor modalities, in contrast to littermates globally lacking CB₁R expression (STOP-CB₁R mice; Figures 4E, 4F, S4G, and S4H; Ruehle et al., 2013), with no effect on direct aversion (Figures 4G, 4H, S4I, and S4J).

Altogether, these data indicate that, within distinct cell type-specific populations of CB₁R (Busquets-Garcia et al., 2015; Marsicano and Kuner, 2008), those present in GABAergic neurons are specifically involved in odor-taste mediated learning.

CB₁R-Dependent Hippocampal Alterations Induced by Odor-Taste Pairings

The results showed above suggest a key role of the hippocampus and of GABAergic neurons in the modulation of odor-taste mediated learning. Thus, we next aimed at determining whether preconditioning recruits classical hippocampal mechanisms associated with memory formation, such as protein synthesis and synaptic plasticity, and the involvement of inhibitory transmission in these processes.

First, we investigated whether incidental learning specifically modulates CB₁R expression in the hippocampus. Immunoblotting experiments revealed that 3 odor-taste pairings, but not the same number of presentations of odor alone, taste alone, or unpaired odor-taste, specifically resulted in enhanced hippocampal expression of CB₁R protein (Figures 5A, S5A, and S5B). Extended preconditioning training is known to suppress the expression of mediated learning (Busquets-Garcia et al., 2017b), suggesting that increasing the number of stimuli associations could trigger cellular and molecular changes. Intriguingly, we observed that hippocampal CB₁R expression was normalized by extended preconditioning training (six odor-taste pairings) (Figures 5A and S5B), further suggesting that increased levels of hippocampal CB₁R protein are reliably associated to the expression of mediated learning. To investigate the




Figure 3. Hippocampal CB_1R Are Required for Mediated Learning

(A) Representative images showing hippocampal expression of CB_1R in control mice (HC-*CB₁R*-WT) and in mice carrying virally induced deletion of the receptor (HC-*CB₁R*-KO). Scale bar, 300 μ m.

(B and C) (B) Effects of specific hippocampal deletion of CB_1R on mediated taste (left) and direct odor (right) learning and (C) on mediated odor (left) and direct taste (right) learning.

(D) Representative images showing hippocampal expression of CB_1R in *CB_1R*-KO mice injected with a control (Stop-*CB_1R*-KO) or with a CB_1R -expressing virus (HC-*CB_1R*-RS). Scale bar, 300 μ m.

(E) Effects of exclusive hippocampal re-expression of CB_1R on mediated taste (left) and direct odor (right) learning. * $p < 0.05$; ** $p < 0.01$ (mCS+ versus mCS- or CS+ versus CS-). For statistical details and n , see [Tables S1](#) and [S2](#).

molecular mechanisms of this effect, we asked whether the enhancement of CB_1R expression was due to new protein synthesis during the preconditioning phase. The administration of the protein synthesis inhibitor Anisomycin (18 mg/kg, i.p.; Puighermanal et al., 2009) before each odor-taste pairing blunted the increase of CB_1R expression (Figure S5C).

Intracellular CB_1R signaling involves many different pathways, including extracellular-regulated kinases (ERKs), mechanistic target of rapamycin (mTOR), and the cAMP response element-binding protein (CREB) (Piomelli, 2003; Puighermanal et al., 2009). Neither the phosphorylation of ERKs nor the activation of the mTOR pathway was affected by exposure to three odor-taste pairings (data not shown). Conversely, CREB phosphorylation was enhanced after three pairings, but not after extended training (Figure S5D). However, whereas exposure to taste alone did not affect CREB phosphorylation, odor alone promoted it (Figure S5D).

CB_1R activation participates in the regulation of synaptic transmission and plasticity (Busquets-Garcia et al., 2018; Castillo et al., 2012; Chevalleyre et al., 2006; Kano et al., 2009). Therefore, we asked whether odor-taste pairings impacted *in vivo* hippocampal long-term potentiation (LTP) induced by high-frequency stimulation (HFS) of the Schaffer collateral-CA1 pathway (Robin et al., 2018). Exposure to three odor-taste pairings enhanced *in vivo* hippocampal LTP in wild-type mice, but not in GABA-*CB_1R*-KO (Figures 5B, 5C, and S5E). However, control experiments indicated that exposure to the taste alone also increased *in vivo* hippocampal LTP in WT, but not in GABA-*CB_1R*-KO mice (Figures 5B and 5C).

Hippocampal CB_1R in GABAergic neurons control inhibitory transmission and plasticity, such as short-term depolarization-induced suppression (DSI) or long-term depression (L-LTD) of hippocampal evoked inhibitory post-synaptic currents (eIPSCs; Araque et al., 2017; Busquets-Garcia et al., 2018; Castillo et al., 2012;




Figure 4. GABAergic *CB₁R* Are Necessary and Sufficient for Mediated Learning

(A–D) Liquid consumption under conditions of mediated taste (A) and odor (B) aversion and direct odor (C) and taste (D) aversion in mice lacking *CB₁R* in forebrain GABAergic neurons (GABA-*CB₁R*-KO) and wild-type littermates (GABA-*CB₁R*-WT).

(E–H) Liquid consumption under conditions of mediated taste (E) or odor (F) aversion and direct odor (G) or taste aversion (H) in mice carrying exclusive expression of *CB₁R* in forebrain GABAergic neurons (GABA-*CB₁R*-RS) and in littermates lacking *CB₁R* expression (STOP-*CB₁R*). **p* < 0.05; ***p* < 0.01; ****p* < 0.001 (mCS+ versus mCS– or CS+ versus CS). For statistical details, see Tables S1 and S2.

Chevalyere et al., 2006; Kano et al., 2009). Hippocampal slices were prepared from mice exposed to three odor-taste pairings or control conditions, and inhibitory synaptic plasticity was analyzed. Preconditioning altered neither basal inhibitory neurotransmission as assessed by measuring miniature IPSCs (Figure S5F) nor short-term plasticity as determined by DSI measurements (Figure S5G). As previously described (Chevalyere and Castillo, 2003), we observed that the application of two HFS trains induced a reliable I-LTD in 71% of pyramidal neurons from naive control mice (Figures 5D and 5E). Odor-taste preconditioning procedures involve limited access to water (1 hr per day; Busquets-Garcia et al., 2017a, 2017b; see Star Methods). Surprisingly, only 26% of pyramidal neurons in hippocampal slices from water-restricted control mice only exposed to plain water underwent I-LTD after HFS (Figure 5E), indicating an impact of limited access to water on this form of synaptic plasticity. However, paired odor-taste exposures in equally water-restricted mice rescued I-LTD expression to levels indistinguishable from naive animals in both amplitude (Figure 5D) and percentage of responsive cells (Figure 5E). Importantly, these effects were not present in mice exposed either to taste or odor alone (Figures 5D and 5E).

Altogether, these results show that exposure to three odor-taste pairings increases protein synthesis-dependent expression

of *CB₁R*, CREB phosphorylation, *in vivo* LTP, and cellular sensitivity to endocannabinoid-dependent I-LTD in the hippocampus. However, whereas the effects on CREB phosphorylation and LTP are also triggered by the mere exposure to low-salience stimuli such as odors or tastes alone, the enhancements of *CB₁R* expression and I-LTD amplitude and sensitivity are specifically related to preconditioning odor-taste pairings, suggesting their involvement in the formation of incidental associations and the expression of mediated learning.

Specific Hippocampal GABAergic Interneurons Control the Formation of Incidental Associations

Considering the inhibitory role of *CB₁R* on hippocampal GABAergic neurotransmission (Chevalyere et al., 2006; Katona et al., 1999; Ohno-Shosaku et al., 2012) and the enhancement of I-LTD sensitivity induced by preconditioning training, lack of *CB₁R*-dependent inhibition of hippocampal GABAergic transmission during incidental associations might be responsible for the impairment of mediated learning found in *CB₁R*-KO, HC-*CB₁R*-KO, and GABA-*CB₁R*-KO mice. In other words, the lack of *CB₁R*-dependent control might induce excessive inhibitory transmission, impairing incidental learning. To test this hypothesis, we aimed at reducing hippocampal GABAergic transmission




Figure 5. Hippocampal Alterations Induced by Odor-Taste Pairings

(A) Optical densitometric quantification of CB₁R levels in mice that received water, three odor-taste pairings (three OT pairings), odor alone (Odor), taste alone (Taste), unpaired odor-taste exposures (Unp. OT), or six odor-taste pairings (6 OT) (see Figure S5 for the representative gels and controls).

(B) Summary traces (top) and time course plots of normalized fEPSPs recorded *in vivo* before (1) and after (2) high-frequency stimulation (HFS) in the CA1 hippocampal region of GABA-CB₁R-WT mice (left) and GABA-CB₁R-KO littermates (right). Animals were anesthetized for recording immediately after the last exposure to water, odor-taste (3 OT pairings), taste alone, or odor alone, respectively.

(C) Bar histogram representing LTP amplitude during the last 3 min of recording in the same groups as described in (B).

(D) Left, time course plots showing eIPSCs amplitude before and after HFS in hippocampal slices obtained from non-water-restricted mice (Naive), or mice that received water, three odor-taste pairings (3 OT pairings), odor alone (Odor), or taste alone (Taste). Right, averaged eIPSCs recorded 15–20 min after HFS in the same groups.

(E) Top, traces of eIPSCs before and after HFS in representative recordings resulting in I-LTD (left) or no I-LTD (right). I-LTD was defined as a $\geq 15\%$ reduction of eIPSCs as compared to baseline. Bottom, black bar histograms representing the proportion of cells displaying I-LTD in the same groups as described in (D). * $p < 0.05$; ** $p < 0.01$ and *** $p < 0.001$ as compared to Water or KO conditions. # $p < 0.05$ and ## $p < 0.01$ as compared to baseline. NS, not significant. For statistical details and n , see Tables S1 and S2.

in GABA-CB₁R-KO mutant mice during preconditioning, i.e., during each odor-taste pairing, using a chemogenetic approach (Figure 6A). An adeno-associated viral vector carrying Cre-dependent expression of an inhibitory DREADD (DIO-hM4DGi, hereafter called DREADD-Gi) (Robinson and Adelman, 2015) was infused into the hippocampi of GABA-CB₁R-KO mice, leading to specific expression of DREADD-Gi in hippocampal GABAergic neurons to generate GABA-CB₁R-KO (DREADD-Gi) (Figure 6B). To control for the potential effects of CNO in virally infected mice lacking DREADD receptors (Gomez et al., 2017), GABA-CB₁R-WT mice were infused with a control AAV ex-

pressing the mCherry protein (GABA-CB₁R-WT [mCherry]). The DREADD ligand clozapine-N-oxide (CNO) enhanced hippocampal *in vivo* LTP in GABA-CB₁R-KO (DREADD-Gi) mice as compared to saline-treated mice (Figure S6A), thereby indicating the effectiveness of the chemogenetic approach. Importantly, however, CNO injections into GABA-CB₁R-WT (mCherry) mice did not affect the consumption during preconditioning (Figure S6B) and had no effect on mediated (Figures 6B and S6C) or direct aversion (Figures 6B and S6D), showing that the potential unspecific effects of the drug or its metabolites (Gomez et al., 2017) were not present in our experimental conditions. Strikingly,




Figure 6. Specific Hippocampal GABAergic Neurons Control the Formation of Incidental Associations

(A) Scheme of CNO treatments in mice injected with different DREADD-expressing viruses.

(B) Top, representative micrograph image showing the viral expression of Cre-dependent DREADD-Gi in the hippocampus of GABA-CB₁R-KO mice. Middle bottom, effect of chemogenetic inhibition of hippocampal GABAergic neurons during preconditioning on mediated (middle) and direct (bottom) aversion in GABA-CB₁R-KO mice (see Figure S6 for further details; scale bar, 300 μ m). Note the reversal of the mediated learning impairment in the mutants after CNO treatment.

(C) Top, representative micrograph image showing the viral expression of Cre-dependent DREADD-Gq in the hippocampus of GABA-CRE mice. Middle bottom, effect of chemogenetic activation of hippocampal GABAergic neurons during preconditioning on mediated (middle) and direct (bottom) aversion in GABA-DREADD-Gq mice (see Figure S6 for further details; scale bar, 300 μ m). Note mediated learning impairment after CNO treatment.

(D) Top, representative micrograph image showing the mCherry viral expression, the PV interneurons and the merged image in the CA1 region of the hippocampus of PV-CRE mice (see Figure S6 for further details; scale bar, 100 μ m). Middle bottom, effect of chemogenetic activation of hippocampal PV neurons during preconditioning on mediated (middle) and direct (bottom) aversion in PV-DREADD-Gq mice. Note the lack of impairment after CNO treatment. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ (mCS+ versus mCS- or CS+ versus CS). For statistical details, see Tables S1 and S2.

however, the same treatment in GABA-CB₁R-KO (DREADD-Gi) mice fully rescued mediated learning (Figures 6B and S6C), without altering preconditioning consumption or direct aversion (Figures 6B, S6B, and S6D). These results suggest that an excess of inhibitory neurotransmission during the formation of incidental stimulus-stimulus associations might be the cause of the mediated learning impairment in GABA-CB₁R-KO mice.

The data presented so far imply a key role of the inhibition of hippocampal GABAergic transmission during the processing of incidental associations between low-salience stimuli and related synaptic plasticity. Therefore, stimulation of GABAergic transmission in the hippocampus of mice expressing normal levels

of CB₁R should impair these processes and, thereby, block mediated learning. To test this idea, an adeno-associated viral vector carrying Cre-dependent expression of an excitatory DREADD (DIO-hM3DGq, hereafter called DREADD-Gq) (Robinson and Adelman, 2015) was infused into the hippocampi of mice expressing the Cre recombinase in Dlx5/6-positive cells (Figure 6C), encompassing all GABAergic interneurons (GABA-Cre mice; Monory et al., 2006) to obtain GABA-DREADD-Gq mice. We first verified that CNO administration was able to increase the number of cFos-positive neurons in the hippocampus of these mice (Figure S6E) and that CNO treatment did not alter mediated or direct aversion in GABA-Cre mice infused with a

control Cre-dependent virus (GABA-mCherry; Figure S6F). Notably, however, CNO administration during preconditioning fully blocked mediated, but not direct, aversion in GABA-DREADD-Gq mice (Figures 6C, S6G, and S6H).

These data are in agreement with the idea that hippocampal inhibitory neurotransmission plays a key role in the formation of incidental associations. However, generalized activation of GABAergic neurons by DREADD-Gq does not allow determination of whether there is a specific population of GABAergic neurons modulating mediated learning. In this context, it is important to note that CB₁R are expressed in specific interneuronal subpopulations in the hippocampus, CCK-positive but not in parvalbumin (PV)-positive basket cells (Katona et al., 1999; Marsicano and Kuner, 2008; Marsicano and Lutz, 1999). CCK- and PV-positive interneurons are two functionally and anatomically distinct cell populations, which together encompass the large majority of inhibitory basket cells innervating somas and proximal dendrites of hippocampal pyramidal neurons (Whissell et al., 2015). Due to the additional expression of CCK in pyramidal hippocampal neurons, the specific targeting of CCK- and CB₁R-positive interneurons is technically very challenging and laborious (Dimidschstein et al., 2016). Conversely, the use of PV-Cre mice (Courtin et al., 2014) allows a reliable targeting of PV-positive interneurons, which represent approximately half of the hippocampal basket cells (Whissell et al., 2015). Therefore, we injected adeno-associated viruses expressing Cre-dependent DREADD-Gq into the hippocampi of PV-Cre mice, obtaining the localization of the receptor in PV-positive neurons (>85% colocalization, Figure 6D), thereby generating PV-DREADD-Gq mice. First, we verified the functionality of the approach, by showing that CNO administration was able to increase the number of c-Fos-positive neurons in the hippocampus of these mice (Figure S6I). Strikingly, the specific activation of PV interneurons by CNO administration during preconditioning did not affect mediated learning (Figures 6D, S6J, and S6K). These data suggest that specific PV-negative subpopulations of GABAergic hippocampal interneurons are involved in the development of incidental associations between low-salience stimuli, eventually enabling the successive formation of mediated learning and memory.

DISCUSSION

Our daily behavioral choices are mainly based on past experiences. These behaviors can depend on direct associative memories, where sensory stimuli are directly associated with specific aversive or rewarding situations. However, they often originate from previous incidental stimulus-stimulus associations between low-salience sensory cues, which are able to assign new value to stimuli that were never directly reinforced (Bornstein et al., 2017; Shohamy and Wagner, 2008; Wimmer and Shohamy, 2012). Stimulus-stimulus associations can be evaluated through the sensory preconditioning paradigm in humans (Bornstein et al., 2017; Shohamy and Wagner, 2008; Wimmer and Shohamy, 2012) and animals (Gewirtz and Davis, 2000; Parkes and Westbrook, 2011). Using this task, our data strongly suggest that the CB₁R-dependent control of discrete subpopulations of hippocampal GABAergic interneurons is necessary

and sufficient for the processing of incidental stimulus-stimulus associations. Accordingly, incidental associations induce a specific protein synthesis-dependent increase of hippocampal CB₁R expression, which is accompanied by an enhancement of hippocampal sensitivity to CB₁R-dependent synaptic plasticity. Thus, our data are compatible with a scenario in which incidental associations of different stimuli would trigger CB₁R activity and induce CB₁R-dependent synaptic plasticity, which would “prime” hippocampal circuits allowing mediated learning. In this context, our data reveal that the CB₁R-dependent control of hippocampal inhibitory synaptic neurotransmission is a key element of high-order cognitive processes, which allow the flexible use of the complex and changing patterns of sensory information that characterize daily individual experiences. Thus, the hippocampal ECS appears to enhance the repertoire of possible behavioral choices dictated by previous experiences, thereby increasing the survival potential of individuals.

Importantly, these mechanisms do not appear to be limited to specific sensory modalities of the stimuli. Despite the fact that tastes can also act as primary reinforcers (Yiannakas and Rosenblum, 2017), gustatory cues have been widely used in sensory preconditioning protocols as low-salience stimuli (Wheeler et al., 2013). Important control experiments showed that these cues are unable to elicit direct odor conditioning per se in our experimental conditions. Moreover, inhibition of CB₁R signaling impairs mediated learning when either tastes or odors are used as cues to elicit aversion. Thus, we can conclude that our odor-taste preconditioning paradigm is suitable to identify the formation of incidental associations between stimuli that are of low salience for the mice. In line with this idea, the general implication of the ECS in incidental learning extends also to other sensory preconditioning paradigms, in which low-salience auditory and visual stimuli are used and appetitive behavior is evaluated. Thus, the common involvement of the ECS in different experimental conditions suggests that similar mechanisms might underline higher-order cognitive processes independently of the sensory modalities used and of the nature (aversive or appetitive) of the reinforcer. Future studies will address this intriguing hypothesis that would lead to a unified vision of complex higher-order learning processes in the brain.

The hippocampus has been suggested to be necessary for associating and temporarily maintaining an internal record of different stimuli simultaneously presented (Voss et al., 2017), making it a key brain region for the integration of different sensory information. Accordingly, previous studies supported the importance of the hippocampus in sensory preconditioning both in humans and in animals (Iordanova et al., 2011; Talk et al., 2002; Wheeler et al., 2013; Wimmer and Shohamy, 2012). However, these studies focused on the role of the hippocampus during the second phase of the procedure (conditioning) or during the retrieval test, when incidental associations have already formed. Thus, these studies underlined the role of the hippocampus in enabling the value of the reinforcer to spread across the associated and not explicitly reactivated item, rather than in the processing and recording of incidental associations per se. Conversely, the present results reveal that the hippocampus is also involved in the preconditioning phase, when the coincidence of low-salience stimuli is recorded and stored for future

potential use. Specifically, our data provide an unforeseen physiological link between hippocampal GABAergic signaling and associative memory between low-salience events.

CB₁R blockade (or activation) before testing does not affect the expression of mediated aversion (Busquets-Garcia et al., 2017b), strongly arguing for a specific role of hippocampal endocannabinoid signaling in the processing of incidental stimulus-stimulus associations. Interestingly, recent work pointed to the importance of other brain regions, such as the orbitofrontal, retrosplenial, and perirhinal cortices for the processing of incidental associations between low-salience sensory cues (Holmes et al., 2013, 2018; Robinson et al., 2014; Sadacca et al., 2018). Interestingly, these regions are highly connected to the hippocampus (Agster and Burwell, 2013; Ritchey et al., 2015; Witter et al., 2017), suggesting that so-far-unexplored network activities between neocortical structures and CB₁R-controlled inhibitory circuits in the hippocampus allow the brain to record and store associations between different external stimuli that are constantly present in the environment.

The cellular and molecular mechanisms underlying ECS-dependent control of incidental learning in the hippocampus appear to be quite complex. Exposure to low-salience stimuli enhances *in vivo* hippocampal LTP and CREB phosphorylation. However, these effects are not specific to incidental associations, because they are also induced by the mere exposure to gustatory or olfactory stimuli alone, respectively. On the other hand, the simultaneous presentation of stimuli induces specific molecular and cellular effects. Incidental learning causes a protein synthesis-dependent increase of CB₁R expression and higher hippocampal sensitivity to ECS-dependent plasticity of inhibitory neurotransmission. The increased sensitivity of hippocampal pyramidal neurons to undergo I-LTD might be due to the increase of CB₁R expression. However, the fact that preconditioning does not alter other forms of ECS-dependent plasticity, such as DSI, suggests that the enhanced I-LTD sensitivity might result from other mechanisms, such as, for instance, increased endocannabinoid mobilization following HFS. Whereas enhancement of hippocampal CB₁R expression and I-LTD sensitivity are specific to odor-taste preconditioning (i.e., they do not occur under exposure to odors or tastes alone), the increased *in vivo* LTP induced by both odor-taste and taste alone is abolished in GABA-CB₁R-KO mice. Differently from I-LTD, this ECS-dependent phenomenon and the enhanced phosphorylation of CREB cannot be linked to increased CB₁R expression levels, because they occur in conditions where these levels are not changed (i.e., under exposure to single stimulus). Future experiments will address these complex molecular and cellular interactions. However, the present data allow speculating that incidental learning might occur via a “two-step” hippocampal CB₁R-dependent process: (1) First, animals might record the presence of stimuli independently of their possible association. CB₁R-dependent increase of LTP might underline this process, at least for taste-related information. (2) The simultaneous presence of odors and tastes then triggers additional processes, again involving the ECS (increased CB₁R expression and I-LTD sensitivity). In this frame, step (1) might represent a preliminary event (stimulus detection) that is necessary to further develop step (2), which would underline the “actual” inci-

dental learning. Future studies will address these intriguing possibilities.

Our previous data revealed that extended preconditioning training (six odor-taste pairings instead of three) suppresses mediated learning, installing the so-called “reality testing,” through which the animals are able to successfully discriminate between stimuli that they initially reacted to in the same way (Busquets-Garcia et al., 2017b; McLaren and Mackintosh, 2002). Notably, extended training also normalizes the hippocampal expression of CB₁R. Thus, it will be extremely interesting to investigate the causes and the consequences of these molecular adaptations in the training-dependent behavioral switch from mediated learning to “reality testing” responses (McDannald and Schoenbaum, 2009).

It has been suggested that the numerous and largely diverse types of hippocampal GABAergic interneurons are differentially required for specific functions and contribute to the computing of cognitive processes in different conditions (Caroni, 2015; Chevalyere and Piskorowski, 2014; Klausberger and Somogyi, 2008). An important corollary of our data is that selective subpopulations of hippocampal interneurons are specifically involved in the processing of incidental stimulus-stimulus associations. Hippocampal CB₁R are abundantly expressed in a specific subset of basket cells, the inhibitory interneurons specialized in the innervation of somas and proximal dendrites of pyramidal neurons (Katona et al., 1999, 2000; Marsicano and Kuner, 2008; Marsicano and Lutz, 1999). The expression of PV or CCK characterizes two non-overlapping subpopulations of basket cells (Klausberger et al., 2005), with CB₁R almost exclusively expressed in CCK-positive cells (Katona et al., 1999, 2000; Marsicano and Kuner, 2008; Marsicano and Lutz, 1999). Notably, whereas global activation of GABAergic neurons in the hippocampus during preconditioning blocks mediated learning, selective activation of PV-positive hippocampal neurons does not alter sensory incidental associations nor mediated responses. Thus, our data suggest that the activity of CCK- and CB₁R-expressing cells plays a key role in the processing of incidental learning. According to the large inhibition exerted by both PV- and CCK-positive interneurons on the firing properties of pyramidal cells, this conclusion might appear surprising. However, the known wide functional differences between the microcircuits formed by these two interneuronal subpopulations (Armstrong and Soltesz, 2012; Bartos and Elgueta, 2012) might underline this intriguing dichotomy. Thus, our data suggest that activity- and endocannabinoid-dependent reduction of neurotransmission of a selected population of hippocampal interneurons is required during incidental associations of low-salience stimuli, to eventually enable mediated learning.

In conclusion, the tight regulation of hippocampal GABAergic interneurons by CB₁R can explain how people integrate and associate different low-salience stimuli randomly encountered to develop seemingly ungrounded attraction or aversion toward particular objects, places, or people. Importantly, alterations of mediated learning, also known as associative inference, can be found in several psychiatric diseases (Armstrong et al., 2012a, 2012b). Thus, these results suggest that modulating specific inhibitory hippocampal transmission via CB₁R might

represent a potential therapeutic target against cognitive dysfunctions in neuropsychiatric conditions.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- **KEY RESOURCES TABLE**
- **CONTACT FOR REAGENT AND RESOURCE SHARING**
- **EXPERIMENTAL MODEL AND SUBJECT DETAILS**
 - Animals
- **METHOD DETAILS**
 - Drug preparation and administration
 - Chemical odors and tastes
 - Sensory preconditioning tasks
 - Western Blot analysis
 - Virus generation
 - Surgery and viral administration
 - Immunohistochemistry and fluorescence detection
 - *In vivo* electrophysiology in anesthetized mice
 - *Ex vivo* electrophysiology
- **QUANTIFICATION AND STATISTICAL ANALYSIS**
 - Data collection
 - Statistical analyses

SUPPLEMENTAL INFORMATION

Supplemental Information includes six figures and two tables and can be found with this article at <https://doi.org/10.1016/j.neuron.2018.08.014>.

A video abstract is available at <https://doi.org/10.1016/j.neuron.2018.08.014#mmc3>.

ACKNOWLEDGMENTS

We thank Delphine Gonzales, Nathalie Aubailly, and all the personnel of the Animal Facility of the NeuroCentre Magendie for mouse care. We also thank all the members of the Marsicano lab for useful discussions, Virginie Morales for invaluable help with administrative work, and Isabel Matias and Ilaria Beluomo for their help in processing the samples for the immunoblot. We thank Melissa Sharpe for helpful suggestions regarding the light-tone sensory preconditioning task and Cyril Herry for providing the PV-Cre mice. We also thank Shauna Parkes and Etienne Coutureau for their useful and critical reading on an early version of the manuscript. This work was supported by INSERM (to G.M.), INRA (to G.F., P.T., and X.F.), EU-FP7 (PAINCAGE, HEALTH-603191 to G.M.; FP7-PEOPLE-2013-IEF-623638 to A.B.-G.; and PRESTIGE-2017-2-0031 to A.C.), European Research Council (Endofood, ERC-2010-StG-260515 and CannaPreg, ERC-2014-PoC-640923, to G.M.), Fondation pour la Recherche Medicale (DRM20101220445 to G.M. and FDT20160435664 to J.F.O.C.; FDT20170436845 to G.T.). Human Frontiers Science Program (to G.M.), Region Aquitaine (to G.M.), French State/Agence Nationale de la Recherche (LABEX BRAIN ANR-10-LABX-43 to G.M., G.F. and A.B.G., ANR-10-IDEX-03-02 to A.B.-G., NeuroNutriSens ANR-13-BSV4-0006-02 to G.M., Orups ANR-16-CE37-0010 to G.M. and G.F., and SynLip ANR-16-CE16-0022 to P.T.), Fyssen Foundation (to E.S.-G.), and CONACyT (to E.S.-G.).

AUTHOR CONTRIBUTIONS

A.B.-G., G.F., and G.M. designed research; A.B.-G., J.F.O.C., E.S.-G., A.C.P.Z., G.T., A.C., B.R., M.V., and C.I. performed research; A.B.-G., G.F., and G.M. supervised research; A.B.-G., J.F.O.C., A.C.P.Z., G.T., P.T., G.F.,

and G.M. analyzed data; A.B.-G., G.F., and G.M. wrote the manuscript. All authors edited and approved the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing financial interests.

Received: August 7, 2017

Revised: March 16, 2018

Accepted: August 9, 2018

Published: August 30, 2018

REFERENCES

- Agster, K.L., and Burwell, R.D. (2013). Hippocampal and subicular efferents and afferents of the perirhinal, postrhinal, and entorhinal cortices of the rat. *Behav. Brain Res.* *254*, 50–64.
- Araque, A., Castillo, P.E., Manzoni, O.J., and Tonini, R. (2017). Synaptic functions of endocannabinoid signaling in health and disease. *Neuropharmacology* *124*, 13–24.
- Armstrong, C., and Soltesz, I. (2012). Basket cell dichotomy in microcircuit function. *J. Physiol.* *590*, 683–694.
- Armstrong, K., Kose, S., Williams, L., Woolard, A., and Heckers, S. (2012a). Impaired associative inference in patients with schizophrenia. *Schizophr. Bull.* *38*, 622–629.
- Armstrong, K., Williams, L.E., and Heckers, S. (2012b). Revised associative inference paradigm confirms relational memory impairment in schizophrenia. *Neuropsychology* *26*, 451–458.
- Bartos, M., and Elgueta, C. (2012). Functional characteristics of parvalbumin- and cholecystokinin-expressing basket cells. *J. Physiol.* *590*, 669–681.
- Bellochio, L., Lafenêtre, P., Cannich, A., Cota, D., Puente, N., Grandes, P., Chaouloff, F., Piazza, P.V., and Marsicano, G. (2010). Bimodal control of stimulated food intake by the endocannabinoid system. *Nat. Neurosci.* *13*, 281–283.
- Bellochio, L., Ruiz-Calvo, A., Chiarlone, A., Cabanas, M., Resel, E., Cazalets, J.R., Blázquez, C., Cho, Y.H., Galve-Roperh, I., and Guzmán, M. (2016). Sustained Gq-protein signaling disrupts striatal circuits via JNK. *J. Neurosci.* *36*, 10611–10624.
- Bornstein, A.M., Khaw, M.W., Shohamy, D., and Daw, N.D. (2017). Reminders of past choices bias decisions for reward in humans. *Nat. Commun.* *8*, 15958.
- Busquets-García, A., Desprez, T., Metna-Laurent, M., Bellochio, L., Marsicano, G., and Soria-Gomez, E. (2015). Dissecting the cannabinergic control of behavior: The where matters. *BioEssays* *37*, 1215–1225.
- Busquets-García, A., Gomis-González, M., Srivastava, R.K., Cutando, L., Ortega-Alvaro, A., Ruehle, S., Remmers, F., Bindila, L., Bellochio, L., Marsicano, G., et al. (2016). Peripheral and central CB1 cannabinoid receptors control stress-induced impairment of memory consolidation. *Proc. Natl. Acad. Sci. USA* *113*, 9904–9909.
- Busquets-García, A., Soria-Gómez, E., Ferreira, G., and Marsicano, G. (2017a). Representation-mediated aversion as a model to study psychotic-like states in mice. *Bio-protocol* *7*, e2358.
- Busquets-García, A., Soria-Gómez, E., Redon, B., Mackenbach, Y., Vallée, M., Chaouloff, F., Varilh, M., Ferreira, G., Piazza, P.V., and Marsicano, G. (2017b). Pregnenolone blocks cannabinoid-induced acute psychotic-like states in mice. *Mol. Psychiatry* *22*, 1594–1603.
- Busquets-García, A., Bains, J., and Marsicano, G. (2018). CB₁ receptor signaling in the brain: extracting specificity from ubiquity. *Neuropsychopharmacology* *43*, 4–20.
- Caroni, P. (2015). Inhibitory microcircuit modules in hippocampal learning. *Curr. Opin. Neurobiol.* *35*, 66–73.
- Castillo, P.E., Younts, T.J., Chávez, A.E., and Hashimoto, Y. (2012). Endocannabinoid signaling and synaptic function. *Neuron* *76*, 70–81.
- Chevalyère, V., and Castillo, P.E. (2003). Heterosynaptic LTD of hippocampal GABAergic synapses: a novel role of endocannabinoids in regulating excitability. *Neuron* *38*, 461–472.

- Chevalyere, V., and Piskorowski, R. (2014). Modulating excitation through plasticity at inhibitory synapses. *Front. Cell. Neurosci.* 8, 93.
- Chevalyere, V., Takahashi, K.A., and Castillo, P.E. (2006). Endocannabinoid-mediated synaptic plasticity in the CNS. *Annu. Rev. Neurosci.* 29, 37–76.
- Courtin, J., Chaudun, F., Rozeske, R.R., Karalis, N., Gonzalez-Campo, C., Wurtz, H., Abdi, A., Baufreton, J., Bienvenu, T.C., and Herry, C. (2014). Prefrontal parvalbumin interneurons shape neuronal activity to drive fear expression. *Nature* 505, 92–96.
- Dimidschstein, J., Chen, Q., Tremblay, R., Rogers, S.L., Saldi, G.A., Guo, L., Xu, Q., Liu, R., Lu, C., Chu, J., et al. (2016). A viral strategy for targeting and manipulating interneurons across vertebrate species. *Nat. Neurosci.* 19, 1743–1749.
- Gewirtz, J.C., and Davis, M. (2000). Using pavlovian higher-order conditioning paradigms to investigate the neural substrates of emotional learning and memory. *Learn. Mem.* 7, 257–266.
- Gomez, J.L., Bonaventura, J., Lesniak, W., Mathews, W.B., Sysa-Shah, P., Rodriguez, L.A., Ellis, R.J., Richie, C.T., Harvey, B.K., Dannals, R.F., et al. (2017). Chemogenetics revealed: DREADD occupancy and activation via converted clozapine. *Science* 357, 503–507.
- Gutiérrez-Rodríguez, A., Puente, N., Elezgarai, I., Ruehle, S., Lutz, B., Reguero, L., Gerrikagoitia, I., Marsicano, G., and Grandes, P. (2017). Anatomical characterization of the cannabinoid CB₁ receptor in cell-type-specific mutant mouse rescue models. *J. Comp. Neurol.* 525, 302–318.
- Hebert-Chatelain, E., Desprez, T., Serrat, R., Bellocchio, L., Soria-Gomez, E., Busquets-Garcia, A., Pagano Zottola, A.C., Delamarre, A., Cannich, A., Vincent, P., et al. (2016). A cannabinoid link between mitochondria and memory. *Nature* 539, 555–559.
- Holmes, N.M., Parkes, S.L., Killcross, A.S., and Westbrook, R.F. (2013). The basolateral amygdala is critical for learning about neutral stimuli in the presence of danger, and the perirhinal cortex is critical in the absence of danger. *J. Neurosci.* 33, 13112–13125.
- Holmes, N.M., Raipuria, M., Qureshi, O.A., Killcross, S., and Westbrook, R.F. (2018). Danger changes the way the mammalian brain stores information about innocuous events: a study of sensory preconditioning in rats. *eNeuro* 5, ENEURO.0381-17.2017.
- Iordanova, M.D., Good, M., and Honey, R.C. (2011). Retrieval-mediated learning involving episodes requires synaptic plasticity in the hippocampus. *J. Neurosci.* 31, 7156–7162.
- Kano, M., Ohno-Shosaku, T., Hashimoto-dani, Y., Uchigashima, M., and Watanabe, M. (2009). Endocannabinoid-mediated control of synaptic transmission. *Physiol. Rev.* 89, 309–380.
- Kato, A., Punnakkal, P., Pernia-Andrade, A.J., von Scholtz, C., Sharopov, S., Nyilas, R., Katona, I., and Zeilhofer, H.U. (2012). Endocannabinoid-dependent plasticity at spinal nociceptor synapses. *J. Physiol.* 590, 4717–4733.
- Katona, I., Sperlách, B., Sík, A., Káfalvi, A., Vizi, E.S., Mackie, K., and Freund, T.F. (1999). Presynaptically located CB₁ cannabinoid receptors regulate GABA release from axon terminals of specific hippocampal interneurons. *J. Neurosci.* 19, 4544–4558.
- Katona, I., Sperlách, B., Maglóczy, Z., Sántha, E., Kófalvi, A., Czirják, S., Mackie, K., Vizi, E.S., and Freund, T.F. (2000). GABAergic interneurons are the targets of cannabinoid actions in the human hippocampus. *Neuroscience* 100, 797–804.
- Kawamura, Y., Fukaya, M., Maejima, T., Yoshida, T., Miura, E., Watanabe, M., Ohno-Shosaku, T., and Kano, M. (2006). The CB₁ cannabinoid receptor is the major cannabinoid receptor at excitatory presynaptic sites in the hippocampus and cerebellum. *J. Neurosci.* 26, 2991–3001.
- Klausberger, T., and Somogyi, P. (2008). Neuronal diversity and temporal dynamics: the unity of hippocampal circuit operations. *Science* 321, 53–57.
- Klausberger, T., Marton, L.F., O’Neill, J., Huck, J.H., Dalezios, Y., Fuentealba, P., Suen, W.Y., Papp, E., Kaneko, T., Watanabe, M., et al. (2005). Complementary roles of cholecystokinin- and parvalbumin-expressing GABAergic neurons in hippocampal network oscillations. *J. Neurosci.* 25, 9782–9793.
- Kobilo, T., Hazvi, S., and Dudai, Y. (2007). Role of cortical cannabinoid CB₁ receptor in conditioned taste aversion memory. *Eur. J. Neurosci.* 25, 3417–3421.
- LeDoux, J.E. (2014). Coming to terms with fear. *Proc. Natl. Acad. Sci. USA* 111, 2871–2878.
- Lu, H.C., and Mackie, K. (2016). An introduction to the endogenous cannabinoid system. *Biol. Psychiatry* 79, 516–525.
- Marsicano, G., and Kuner, R. (2008). Anatomical distribution of receptors, ligands and enzymes in the brain and the spinal cord: circuitries and neurochemistry. In *Cannabinoids and the Brain*, A. Kofalvi, ed. (New York: Springer), pp. 161–202.
- Marsicano, G., and Lutz, B. (1999). Expression of the cannabinoid receptor CB₁ in distinct neuronal subpopulations in the adult mouse forebrain. *Eur. J. Neurosci.* 11, 4213–4225.
- Marsicano, G., Wotjak, C.T., Azad, S.C., Bisogno, T., Rammes, G., Cascio, M.G., Hermann, H., Tang, J., Hofmann, C., Zieglgänsberger, W., et al. (2002). The endogenous cannabinoid system controls extinction of aversive memories. *Nature* 418, 530–534.
- Marsicano, G., Goodenough, S., Monory, K., Hermann, H., Eder, M., Cannich, A., Azad, S.C., Cascio, M.G., Gutiérrez, S.O., van der Stelt, M., et al. (2003). CB₁ cannabinoid receptors and on-demand defense against excitotoxicity. *Science* 302, 84–88.
- McDannald, M., and Schoenbaum, G. (2009). Toward a model of impaired reality testing in rats. *Schizophr. Bull.* 35, 664–667.
- McLaren, I.P., and Mackintosh, N.J. (2002). Associative learning and elemental representation: II. Generalization and discrimination. *Anim. Learn. Behav.* 30, 177–200.
- Metna-Laurent, M., Soria-Gómez, E., Verrier, D., Conforzi, M., Jégo, P., Lafenêtre, P., and Marsicano, G. (2012). Bimodal control of fear-coping strategies by CB₁ cannabinoid receptors. *J. Neurosci.* 32, 7109–7118.
- Monory, K., Massa, F., Egertová, M., Eder, M., Blaudzun, H., Westenbroek, R., Kelsch, W., Jacob, W., Marsch, R., Ekker, M., et al. (2006). The endocannabinoid system controls key epileptogenic circuits in the hippocampus. *Neuron* 51, 455–466.
- Ohno-Shosaku, T., Tanimura, A., Hashimoto-dani, Y., and Kano, M. (2012). Endocannabinoids and retrograde modulation of synaptic transmission. *Neuroscientist* 18, 119–132.
- Parkes, S.L., and Westbrook, R.F. (2011). Role of the basolateral amygdala and NMDA receptors in higher-order conditioned fear. *Rev. Neurosci.* 22, 317–333.
- Paxinos, G., and Franklin, K.B.J. (2001). *The Mouse Brain in Stereotaxic Coordinates* (Academic Press).
- Piomelli, D. (2003). The molecular logic of endocannabinoid signalling. *Nat. Rev. Neurosci.* 4, 873–884.
- Puighermanal, E., Marsicano, G., Busquets-Garcia, A., Lutz, B., Maldonado, R., and Ozaita, A. (2009). Cannabinoid modulation of hippocampal long-term memory is mediated by mTOR signaling. *Nat. Neurosci.* 12, 1152–1158.
- Remmers, F., Lange, M.D., Hamann, M., Ruehle, S., Pape, H.C., and Lutz, B. (2017). Addressing sufficiency of the CB₁ receptor for endocannabinoid-mediated functions through conditional genetic rescue in forebrain GABAergic neurons. *Brain Struct. Funct.* 222, 3431–3452.
- Resstel, L.B., Moreira, F.A., and Guimarães, F.S. (2009). Endocannabinoid system and fear conditioning. *Vitam. Horm.* 81, 421–440.
- Ritchey, M., Libby, L.A., and Ranganath, C. (2015). Cortico-hippocampal systems involved in memory and cognition: the PMAT framework. *Prog. Brain Res.* 219, 45–64.
- Robin, L.M., Oliveira da Cruz, J.F., Langlais, V.C., Martin-Fernandez, M., Metna-Laurent, M., Busquets-Garcia, A., Bellocchio, L., Soria-Gomez, E., Papouin, T., Variilh, M., et al. (2018). Astroglial CB₁ receptors determine synaptic D-serine availability to enable recognition memory. *Neuron* 98, 935–944.
- Robinson, S., and Adelman, J.S. (2015). A method for remotely silencing neural activity in rodents during discrete phases of learning. *J. Vis. Exp.* Published online June 22, 2015. <https://doi.org/10.3791/52859>.

- Robinson, S., Todd, T.P., Pasternak, A.R., Luikart, B.W., Skelton, P.D., Urban, D.J., and Bucci, D.J. (2014). Chemogenetic silencing of neurons in retrosplenial cortex disrupts sensory preconditioning. *J. Neurosci.* *34*, 10982–10988.
- Ruehle, S., Remmers, F., Romo-Parra, H., Massa, F., Wickert, M., Wörtge, S., Häring, M., Kaiser, N., Marsicano, G., Pape, H.C., and Lutz, B. (2013). Cannabinoid CB1 receptor in dorsal telencephalic glutamatergic neurons: distinctive sufficiency for hippocampus-dependent and amygdala-dependent synaptic and behavioral functions. *J. Neurosci.* *33*, 10264–10277.
- Sadacca, B.F., Wied, H.M., Lopatina, N., Saini, G.K., Nemirovsky, D., and Schoenbaum, G. (2018). Orbitofrontal neurons signal sensory associations underlying model-based inference in a sensory preconditioning task. *eLife* *7*, e30373.
- Shohamy, D., and Wagner, A.D. (2008). Integrating memories in the human brain: hippocampal-midbrain encoding of overlapping events. *Neuron* *60*, 378–389.
- Soria-Gómez, E., Bellocchio, L., Reguero, L., Lepousez, G., Martin, C., Bendahmane, M., Ruehle, S., Remmers, F., Desprez, T., Matias, I., et al. (2014). The endocannabinoid system controls food intake via olfactory processes. *Nat. Neurosci.* *17*, 407–415.
- Talk, A.C., Gandhi, C.C., and Matzel, L.D. (2002). Hippocampal function during behaviorally silent associative learning: dissociation of memory storage and expression. *Hippocampus* *12*, 648–656.
- Voss, J.L., Bridge, D.J., Cohen, N.J., and Walker, J.A. (2017). A closer look at the hippocampus and memory. *Trends Cogn. Sci.* *21*, 577–588.
- Wheeler, D.S., Chang, S.E., and Holland, P.C. (2013). Odor-mediated taste learning requires dorsal hippocampus, but not basolateral amygdala activity. *Neurobiol. Learn. Mem.* *101*, 1–7.
- Whissell, P.D., Cajanding, J.D., Fogel, N., and Kim, J.C. (2015). Comparative density of CCK- and PV-GABA cells within the cortex and hippocampus. *Front. Neuroanat.* *9*, 124.
- Wimmer, G.E., and Shohamy, D. (2012). Preference by association: how memory mechanisms in the hippocampus bias decisions. *Science* *338*, 270–273.
- Witter, M.P., Doan, T.P., Jacobsen, B., Nilssen, E.S., and Ohara, S. (2017). Architecture of the Entorhinal Cortex A Review of Entorhinal Anatomy in Rodents with Some Comparative Notes. *Front. Syst. Neurosci.* *11*, 46.
- Yiannakas, A., and Rosenblum, K. (2017). The Insula and Taste Learning. *Front. Mol. Neurosci.* *10*, 335.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-goat Alexa 488	Fisher Scientific	CAT #A21467; RRID: AB_10055703
Anti-Guinea pig Alexa 488	Fisher Scientific	CAT # A-11073; RRID: AB_2534117
Anti-rabbit Alexa 488	Fisher Scientific	CAT #PA5-23091; RRID: AB_2540618
Anti-rabbit Alexa 594	Fisher Scientific	CAT # A21211; RRID: AB_10375432
cAMP response element-binding protein (CREB)	Cell Signaling Technology	CAT #9104; RRID: AB_490881
CB1 receptor (Western Blot)	Abcam	CAT Ab23703; RRID:AB_447623
cFOS polyclonal antibody	Santa Cruz Biotechnology	CAT #Sc-52; RRID: AB_2106783
DAPI	Life Technologies	CAT #D3571; RRID: AB_2307445
DsRED	Clontech Laboratories	CAT #632496; RRID: AB_10013483
Glyceraldehyde-3-phosphate dehydrogenase (GAPDH)	Cell Signaling Technology Danvers	CAT #2118; RRID: AB_561053
Goat polyclonal CB1 receptor (Immunofluorescence)	Frontier Science Co	CB1-Go-Af450-1; RRID: AB_2571530
Guinea pig polyclonal parvalbumin	Synaptic Systems	CAT #1955004; RRID: AB_2156476
HRP-linked antibodies	Cell Signaling Technology	CAT #7074 (Anti Rabbit); RRID: AB_2099233; CAT #7076 (Anti Mouse); RRID: AB_330924
Phospho-cAMP response element-binding protein (pCREB)	Cell Signaling Technology Danvers	CAT #9198; RRID:AB_2561044
Bacterial and Virus Strains		
AAV-CAG-CRECre	Hebert-Chatelain et al., 2016	Virus n3 lab stock
AAV-CBA-GFP	Hebert-Chatelain et al., 2016	Virus n4 lab stock
AAV-CBA-mCB1	Hebert-Chatelain et al., 2016	Virus n6 lab stock
AAV-DIO-hM3Dq	UNC Vector Core	N/A
AAV-DIO-hM4Di	UNC Vector Core	N/A
AAV-DIO-mCherryCHERRY	UNC Vector Core	N/A
Chemicals, Peptides, and Recombinant Proteins		
2-Mercaptoethanol (as component of Laemmli Buffer 4X, 5%)	Sigma-Aldrich	CAT#3148
4-20% Mini-PROTEAN TGX Stain-Free Precast Gels	Bio-Rad	CAT#4568096
Anisomycin	Sigma-Aldrich	CAT#A9789
Anti-Fluorescein-POD	Sigma Aldrich	11426346910
Benzaldehyde	Sigma-Aldrich	CAT#418099
Bromophenol Blue (as component of Laemmli Buffer 4X, 0.02%)	Sigma-Aldrich	CAT#114391
CaCl ₂	Sigma-Aldrich	CAT#C3881
Clarity Western ECL Substrates	Bio-Rad	CAT#1705060
Clozapine-N-oxide (CNO)	Tocris Bioscience	CAT#4936
Complete Protease inhibitors cocktail	Roche	CAT#000000011697498001
Cremophor EL	Sigma-Aldrich	CAT#C5135
DAPI	Fisher Scientific	CAT #11530306
D-APV	Abcam	CAT# ab120003
DMSO	Sigma-Aldrich	CAT#D5879-M
Donkey serum	Sigma Aldrich	S30-100ML
EDTA	Sigma-Aldrich	CAT#EDS
EGTA	Sigma-Aldrich	CAT#E4378
Glucose	Sigma-Aldrich	CAT#G57-67

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Glycerol (as component of Laemmli Buffer 4X, 40%)	VWR	CAT#24386.298
HEPES	Sigma-Aldrich	CAT#H3375
Hydrochloric acid (HCl)	Sigma-Aldrich	CAT#
Isoamyl acetate	Sigma-Aldrich	CAT#W205508
Isoflurane (Vetflurane)	Virbac	N/A
KCL	Merck Millipore	CAT#104936
Ketamine (IMALGENE 1000)	Merck	N/A
Lidocaine (Lurocaine)	Vetoquinol	N/A
Lithium Chloride	Sigma-Aldrich	CAT#L4408
Maltodextrin	Sigma-Aldrich	CAT#419672
Membrane Immobilon-P, PVDF 0.45 μ m	Merck	CAT#IPVH00010
Mg-ATP	Sigma-Aldrich	CAT#A9187
MgCl ₂	VWR chemicals	CAT#36226
NaCl	Sigma-Aldrich	CAT#746398
NaCl	VWR	CAT#0241
Na-GTP	Sigma-Aldrich	CAT#G8877
NaH ₂ PO ₄	Merck Millipore	CAT#106345
NaHCO ₃	Sigma-Aldrich	CAT#S6014
NBQX	Abcam	CAT#ab120046
Non-fat milk (skim milk powder)	Régilat	N/A
Paraformaldehyde	Sigma-Aldrich	CAT#HT501128
Phosphatase inhibitors cocktail (PhosSTOP)	Roche	CAT#00000004906845001
Phosphocreatin	Sigma-Aldrich	CAT#P7936
Pontamine sky blue	Sigma-Aldrich	CAT#C8679
Roti-Nanoquant	Carl Roth	CAT#K880.2
SDS (as component of Laemmli Buffer 4X, 8%)	Carl Roth	CAT#2326.3
Sodium acetate	Sigma-Aldrich	CAT# S2889
Sodium chloride 0,9%	Cooper	N/A
SR101	Cayman Chemical	CAT#9000484
Sucrose	Sigma-Aldrich	CAT#S7903
Tris(hydroxymethyl)aminomethane (Tris)	Research Organics	CAT#30955T
Triton X-100	Sigma-Aldrich	CAT#X100
TSA Plus Fluorescein	Perkin Elmer	NEL741001KT
TTX	Tocris Bioscience	CAT#1069
Tween20	Sigma-Aldrich	CAT#P1379
Experimental Models: Organisms/Strains		
Mouse: C57BL/6	JANVIER Labs	C57BL/6
Mouse: <i>CB₁R</i> -flox	Marsicano et al., 2003	N/A
Mouse: <i>CB₁R</i> -KO	Marsicano et al., 2003	N/A
Mouse: DLX- <i>CB₁R</i> -KO	Monory et al., 2006 , Bellocchio et al., 2010	N/A
Mouse: DLX-CRECre	Bellocchio et al., 2010	N/A
Mouse: GABA- <i>CB₁R</i> -Rescue	Ruehle et al., 2013	N/A
Mouse: PV-Cre	Courtin et al., 2014	N/A
Software and Algorithms		
Axograph	AxoGraph Software	N/A
Behavioral Scoring Panel	A. Dubreug	N/A
Clampfit, pClamp10	Molecular devices	N/A

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Digidata 1440A	Molecular devices	N/A
GrabBee	N/A	N/A
GraphPad prism 6.0	GraphPad Software	N/A
ImageJ	NIH	N/A
Spike2	Cambridge Electronic Design	N/A
Other		
Amplifier	DAGAN Corporation	2400A
Concentric bipolar electrode model	FHC	CBARC50
Constant Current Stimulator	Digitimer	DS3
Data acquisition unit	Cambridge Electronic Design	CED 1401
Homeothermic system model	Harvard Apparatus	50-7087-F
Hydraulic micropositioner	Kopf instruments	Model 2650,
MultiClamp 700B amplifier	Molecular devices	N/A
Stereotaxic frame	Kopf instruments	Model 900
Vibratome VT1200S	Leica	N/A

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for sources should be directed to the Lead Contact, Giovanni Marsicano (giovanni.marsicano@inserm.fr).

EXPERIMENTAL MODEL AND SUBJECT DETAILS**Animals**

All experimental procedures were approved by the Committee on Animal Health and Care of INSERM and the French Ministry of Agriculture and Forestry (authorization numbers, A501350, A3310035, C33063096). Maximal efforts were made to reduce the suffering and the number of mice used. All behavioral experiments were performed during the light phase (from 9am to 1pm) and animals were kept in individual cages.

Male wild-type C57BL/6 mice purchased from Janvier (France) were used for the pharmacological, biochemical and electrophysiological experiments. Male *CB₁R*-floxed mice (Marsicano et al., 2003), *CB₁R* full knockout (*CB₁R*-KO) mice (Marsicano et al., 2003), conditional knockout mice lacking *CB₁R* in forebrain GABAergic *Dlx5/6* positive neurons (*GABA-CB₁R*-KO) (Bellocchio et al., 2010; Monory et al., 2006), mice expressing Cre recombinase under the regulatory elements of the *Dlx5/6* gene (*GABA-Cre*) or the *parvalbumin* (*PV*) gene (*PV-Cre*) and their wild-type littermates mice were obtained, maintained and genotyped as described before (Bellocchio et al., 2010; Courtin et al., 2014; Marsicano et al., 2003; Monory et al., 2006). The rescue line (*GABA-CB₁R*-KO) was generated as described before (Busquets-Garcia et al., 2016; Gutiérrez-Rodríguez et al., 2017; Ruehle et al., 2013; Soria-Gómez et al., 2014). All the mice used in this study were 9-10 weeks old at the beginning of the experiments.

METHOD DETAILS**Drug preparation and administration**

Rimonabant (1 mg/kg) was purchased from Cayman Chemical (Michigan, USA) and was dissolved in a mixture of 4% ethanol, 4% Cremophor-EL and 92% of saline (NaCl 0.9%). Anisomycin (18 mg/kg) was purchased from Sigma-Aldrich (Saint-Quentin-Fallavier, France) and was dissolved in 2% DMSO 2% Tween and 96% saline (NaCl 0.9%). The exogenous DREADD ligand clozapine-N-oxide CNO (2 mg/kg) was purchased from Tocris Bioscience (Bristol, UK) and dissolved in saline after gently mixing with a vortex. All drugs were injected intraperitoneally (i.p.) in a volume of 10 ml/kg. Vehicle injection was composed of 4% ethanol, 4% Cremophor-EL and 92% saline for the experiment using Rimonabant, 2% DMSO, 2% Tween and 96% saline for the experiment using Anisomycin and saline injection for experiments with CNO.

Chemical odors and tastes

The solutions used in the sensory preconditioning task were presented in 50-mL drinking tubes in the home cage with either banana (0.05%, isoamyl acetate) or almond (0.01%, benzaldehyde) for odors, and sucrose (5%) or maltodextrin (5%) for tastes. All compounds were obtained from Sigma-Aldrich (St. Quentin Fallavier Cedex, France). The concentrations of odors and tastes were chosen to be equally preferred (Busquets-Garcia et al., 2017a, 2017b). In all experimental groups (see above) half of the animals

were conditioned with sucrose/banana and half of the mice with maltodextrin/almond. No differences were observed between both conditions in all the experiments performed.

Sensory preconditioning tasks

Odor-taste sensory precondition test. Mice were water deprived in the same room where the whole protocol occurred (Figure S1) as described before (Busquets-Garcia et al., 2017a, 2017b).

Habituation. All subjects received 1 hr access to water during three consecutive days.

Preconditioning phase. This consisted in 6 days of 3 odor-taste pairings (incidental associations). Each pairing consisted in two days: on the first day, the subjects had 1 hr access to a flavored solution containing a new taste (either 5% sucrose or 5% maltodextrin; Taste 1, T1) and a new odorant (0.05% Banana or 0.01% Almond; Odor 1, O1) mixed with water in order to pair T1 with O1. On day two, the animals received the taste and odor not given during the previous day, i.e., Taste 2 (T2) and Odor 2 (O2).

Conditioning or devaluation phase. This phase consisted in 6 days where O1 (or T1) was devaluated as to become the conditioned stimulus (CS+). On days 1, 3 and 5 of this phase, subjects received 1 hr access to O2 (or T2) followed by i.p. injection of saline (CS-). On days 2, 4 and 6, they received 1 hr access to O1 (or T1) immediately followed by an i.p. injection of lithium chloride (LiCl, 0.3 M, 1% b.w.) which induces gastric malaise to the mice (CS+). After conditioning, subjects were given a recovery day in which they could only drink water during 1 hr. In each experimental condition, half of the mice were conditioned with one stimulus (O1 or T1) or the other one (O2 or T2).

Test phase. On the next 2 days, mediated and direct aversions were assessed using a 1 hr two-choice test. Mediated aversion was always evaluated on the first test day with a choice between the stimulus T1 (or O1) previously associated with the CS+ but never directly paired with LiCl (thereby called mediated CS+, mCS+) and the stimulus T2 (or O2) previously associated with the CS- (mediated CS-, mCS-). On the second test day, the direct aversion was evaluated with a choice between the CS+ and CS-. Data are presented as absolute liquid intake.

The pharmacological treatments with Rimonabant (1 mg/kg, i.p.) and CNO (2 mg/kg, i.p.) and their respective vehicles were administered 1 hr before each session during the preconditioning phase. Moreover, Rimonabant (1 mg/kg, i.p.) and its vehicle were also administered 1 hr before the mediated odor learning test to check the effect of CB₁R blockade in this specific test phase. For the experiments using viral approaches, the behavioral protocol started five weeks after surgery.

Additional experiments

We evaluate whether sucrose and maltodextrin used during 3 odor-taste pairings (i.e., preconditioning sessions) induced appetitive conditioning to the odors and what is the effect of Rimonabant treatments during preconditioning on this response. Three different groups of C57BL6 mice underwent the 3-odor-taste pairings (one group injected with vehicle 1 hr before each session, one group receiving Rimonabant before all session and one group injected with Rimonabant before each O1/T1 session and vehicle before each O2/T2 session and). Then, different two-choice tests were performed in consecutive days: choice between odors (O1 versus O2) and between each odor and water (O1 versus water and O2 versus water).

In order to know whether impairment of mediated aversion could be due to delayed formation of mediated learning, mice with a specific deletion of CB₁R in the hippocampus and their controls (HC-CB₁R-KO and HC-CB₁R-WT, see below) and GABA-CB₁R-KO and their littermates underwent an extended preconditioning protocol receiving 6 odor-taste pairings instead of 3 (Busquets-Garcia et al., 2017a, 2017b).

Tone-light sensory preconditioning test

Twenty-two animals were used in this experiment. 10 animals were administered with Rimonabant (1 mg/kg, i.p.) and 12 with its vehicle. Mice were food-deprived to be maintained at 85%–90% of their *ad libitum* weight throughout the duration of the experiment.

Apparatus. Operant chambers (Imétronic, France) had internal dimensions 30x40x36 (cm) and located in a light- and sound-attenuated cabinet with a floor consisting of metal rods. Each operant chamber had two opaque panels at the right and left walls and two clear Plexiglas panels at the back and front walls. A feeder trough was centered on one wall of the chamber. Inside the trough, an infrared photocell detector was used to record head entries into the trough. Food pellets (dehydrated milk) were used as a reward.

Habituation. Animals were exposed to the operant chambers during 20 min for 2 consecutive days, 3 sessions a day, with free access to 4–5 pellets. Saline injection (i.p.) was performed 30 min before the beginning of the first session.

Preconditioning phase. This consisted in two 30 min sessions per day during 2 consecutive days. The “preconditioning session” consisted in the simultaneous presentation of a 10 s tone (65 db, 3000 Hz) with a light (house light located at the top of the chamber), 6 times, with an average intertrial interval (ITI) of 5 min. This was followed 30 min later by a “control” session in which a 10 s click (65 db, 10 Hz) was presented 6 times (average ITI 5 min), with no light. Preconditioning and control sessions were counterbalanced between the 2 days of the preconditioning phase. Rimonabant (1 mg/kg, i.p.) or saline were injected 30 min before the preconditioning phase.

Conditioning phase. This consisted in 1 hr session per day for 6 consecutive days during which the light previously used during the preconditioning phase was presented 12 times (average ITI 5 min) and resulted in the delivery of a food pellet. Head entries in the food trough were recorded during the tone, the click and the ITI. The strength of sensory preconditioning was assessed using a discrimination ratio defined as the amount of responding observed during the first 4 presentations of the tone divided by the sum of responding during the first 4 presentations of the tone and the click.

Test phases. 1 hr session was performed during which the tone and the click were randomly presented 6 times each (average ITI 5 min) in the absence of any food reward. Head entries in the food trough were recorded.

The next day, response to the light cue was assessed through a 1 hr session during which the light was presented 12 times (average ITI 5 min) with no food delivery and head entries in the food trough were recorded.

Western Blot analysis

Just after the last preconditioning session (3 or 6 odor-taste pairings and control experimental groups), C57BL6 mice were sacrificed and used for the Immunoblot. For these experiments, different control groups were used: mice receiving water during all the “preconditioning” phase, mice receiving only taste presentations (sucrose on days 1, 3, 5 and maltodextrin on days 2, 4, 6 or vice versa), mice receiving only odor presentations (banana on days 1, 3, 5 and almond on days 2, 4, 6 or vice versa) and mice receiving unpaired associations of tastes and odors (tastes on days 1, 3, 5 and odors on days 2, 4, 6 or vice versa). Additional groups of animals were injected with vehicle or the protein synthesis inhibitor anisomycin (18 mg/kg, i.p.) 1 hr before each preconditioning session (3 odor-taste pairings) or water presentation. Immediately after the last 1 hr session of liquid presentation, mice were sacrificed and hippocampal tissues extracted, frozen in dry ice and kept in -80°C . Samples were homogenized in lysis buffer (0.05M Tris-HCl pH 7.4, 0.15M NaCl, 0.001M EDTA, 10% Glycerol, 1% Triton X-100) supplemented with protease and phosphatase inhibitors purchased from Roche (Basel, Switzerland), using the Tissue Lyser (Quiagen, Hilden, Germany). After 10 min incubation at 4°C , samples were centrifuged at 17 000 g for 20 min at 4°C to remove insoluble debris. The protein contents of the total solubilized fractions were determined with Roti-Nanoquant protein quantitation assay, following manufacturer’s instruction (Carl Roth, Karlsruhe, Germany). Protein extracts were mixed with denaturing 4x Laemmli loading buffer and warmed for 30 min at 37°C . Samples (25 μg per lane) were analyzed on 4%–20% precast polyacrylamide gels (Bio-Rad, Hercules, California) and transferred onto PVDF membranes 0.45 μm (Merk Millipore, Billerica, MA). Membranes were blocked in a mixture of Tris-buffered saline and polysorbate 20 (20mM Tris-HCl pH 7.6, 150mM NaCl, 0.05% Tween 20) containing 5% of non-fat milk for 1 h at room temperature (RT). For immunoblotting we used antibodies against CB₁R, (ab23703; 1:200, 1h RT, Abcam, Cambridge, UK), phospho - cAMP response element-binding protein (pCREB, Ser133, #9198; 1:1000, overnight 4°C , Cell Signaling Technology Danvers, MA), glyceraldehyde-3-phosphate dehydrogenase (GAPDH) (#2118; 1:1000, 30 min RT, Cell Signaling Technology Danvers, MA) or CREB (#9104; 1:1000 overnight 4°C , Cell Signaling Technology Danvers, MA). Bound primary antibodies were detected with HRP-linked antibodies (1:2000, Cell Signaling Technology, Danvers, MA) and visualized by enhanced chemiluminescence detection (Clarity Western ECL Substrate, Bio-Rad, Hercules, California). The Optical densities of immunoreactive bands were quantified by the Image Lab software (Bio-Rad, Hercules, California) after acquisition on ChemiDoc Touch (Bio-Rad, Hercules, California). The CB₁R expression levels normalized to the amount of GAPDH (loading control) in the same sample have been expressed as a percentage of control condition (water group). pCREB levels normalized to the amount of total CREB in the same sample have been expressed as a percentage of control condition (water group).

Virus generation

GFP (as control virus), CRE (to delete CB₁R) and CB₁R viral (to re-express CB₁R) constructs were of a mixed serotype AAV1/AAV2 and were generated by Calcium Phosphate transfection of HEK293T cells and purified as described (Bellocchio et al., 2016; Monory et al., 2006). Cre-dependent DIO-hM4Di (DREADD-Gi) and DIO-hM3Dq (DREADD-Gq) fused to mCherry (AAV8 serotype) was provided by UNC Gene Therapy Center (University of North Carolina). In the DREADD experiments, mCherry viral construct for GABA-CB₁R-WT and DIO-mCherry for GABA-Cre mice in order to check for the viral expression and to control the potential effects of the expression of a fluorescent protein in the hippocampus were used.

Surgery and viral administration

Mice were anesthetized by i.p. injection of a mixture of ketamine (100mg/kg, Imalgene 500[®], Merial, France) and xylazine (10mg/kg, Rompun[®], Bayer, France) and placed into a stereotaxic apparatus (Model 900, Kopf instruments, CA, USA) with mouse adaptor and lateral ear bars. For viral intra-hippocampus delivery, AAV vectors were injected with the help of a microsyringe attached to a pump (UMP3-1, World Precision Instruments, FL, USA). Mice were injected with AAV-GFP, AAV-Cre, AAV-CB₁R, AAV-DIO-hM4Di or AAV-DIO-hM3Dq and their control viral vectors (DIO-mCherry or AAV-mCherry) directly into the hippocampus (2 injections of 1 μl per side), with the following coordinates: AP -2 , L ± 1 , DV -2 (1st injection) and $-1,5$ (2nd injection), according to Paxinos and Franklin (Paxinos and Franklin, 2001). Animals were used for behavioral or electrophysiological experiments five weeks after injections in order to get an optimal expression of the viruses. CB₁R deletion or reexpression were verified by immunohistochemistry or fluorescent *in situ* hybridization (see below) and mCherry expression was checked by epifluorescence (see below). All mice used in the behavioral experiments were checked and representative images are shown in Figure 6.

Immunohistochemistry and fluorescence detection

After the behavioral experiment, mice were anesthetized with chloral hydrate (400 mg/kg body weight), transcardially perfused first with phosphate-buffered solution (PB 0.1M, pH 7.4) and then with 4% formaldehyde prepared at 4°C to fix tissues. After brain extraction, serial brain coronal sections were cut at 40 μm and collected in 0.1M PB (pH 7.4) at RT. Sections were permeabilized in a blocking solution of 10% donkey serum, 0.3% Triton X-100 and 0.02% sodium azide prepared in 0.1M PB for 1 hr at RT. Free-floating

sections were incubated with goat CB1R polyclonal primary antibodies raised against a 31 aminoacid C-terminal sequence (NM007726) of the mouse CB1R (CB1-Go-Af450-1; 1:2000, Frontier Science Co. Shinko-nishi, Ishikari, Hokkaido, Japan) for 48h at 4°C. The antibody was prepared in the blocking solution. After several washes, the tissue was incubated with a secondary antibody anti-goat Alexa Fluor 488 (10246392, 1:500, Fisher Scientific) for 2 hr and then washed in 0.1 M PB at RT. All sections were counterstained with 4',6-diamidino-2-phenylindole (DAPI, 1:20000) to visualize cell nuclei, and then were mounted, dried and coverslipped. The sections were analyzed with an epifluorescence Leica DM6000 microscope (Leica, France). For c-FOS staining, GABA-Cre and PV-Cre mice infused with DREADD-Gq were sacrificed 90 min after treatment with saline or CNO (2 mg/kg, i.p.) and the same protocol described above was used. After permeabilization, free-floating sections were incubated with rabbit c-FOS polyclonal primary antibody (sc-52; 1:500; Santa Cruz Biotechnology) and revealed with a secondary antibody anti-rabbit Alexa Fluor 488 (10424752; 1/500; Fisher Scientific). The quantitative analysis of c-FOS expression on sections was conducted on images using ImageJ software from black and white images captured by an epifluorescence Leica DM6000 microscope at 10X (Leica, France). Moreover, the merge between the expression of mCherry and the c-FOS staining was performed in some slices. Brains from animals expressing mCherry were processed in the same way, but omitting antibody incubations. For PV-Cre mice, the same protocol described above was used, with the following modifications. Briefly, after the perfusion, brains were cryoprotected in sucrose 30% PB 0.1M, pH 7.4, and were frozen and kept at -80°C . Free-floating frozen brain sections were collected with a cryostat (40 μm , Microm HM 500M, Microm Microtech). Slices were double stained with guinea pig parvalbumin (195004; 1:1000, Synaptic Systems) and rabbit DsRed (632496; 1:1000, Clontech Laboratories, Inc. Mountain View, CA, USA) polyclonal primary antibodies revealed respectively with secondary antibodies anti-guinea pig Alexa Fluor 488 (10193752, 1:500, Fisher Scientific) and anti-rabbit Alexa Fluor 594 (10266352, 1:500, Fisher Scientific).

Fluorescent *in situ* hybridization

The procedure was performed as described (Bellocchio et al., 2010; Marsicano and Lutz, 1999). After behavioral experiments, mice were killed by cervical dislocation. Their brains were isolated, quickly frozen on dry ice and stored at -80°C until sectioning in a cryostat (14 μm , Microm HM 500M, Microm Microtech). FITC-labeled riboprobes against mouse CB₁R were prepared as described (Kato et al., 2012; Kawamura et al., 2006). We used the tyramide signal amplification (TSA) method to improve the detection sensitivity of the *in situ* hybridization. Slides were incubated with FITC-conjugated tyramide (NEL741001KT, 1:80, PerkinElmer). Blocking buffer and wash buffer TNT were prepared according to the manufacturer's protocol. Slides were incubated in DAPI solution (1:20000, Fisher Scientific, NH, USA), washed, mounted, coverslipped and analyzed by an epifluorescence Leica DM6000 microscope (Leica, Germany).

In vivo electrophysiology in anesthetized mice

Experiments were performed as described in Robin et al. (2018). Immediately after the last session of preconditioning (3 odor-taste pairings), or control procedure (either water, odor alone or taste alone, see above Western Blot Analysis) mice were anesthetized in a box containing 5% Isoflurane (VIRBAC, France) before being placed in a stereotaxic frame (Model 900, Kopf instruments, CA, USA) in which 1.0% to 1.5% of Isoflurane was continuously supplied via an anesthetic mask during the whole duration of the experiment. The body temperature was maintained at $\sim 36.5^{\circ}\text{C}$ using a homeothermic system (model 50-7087-F, Harvard Apparatus, MA, USA) and the state of anesthesia was assessed by mild tail pinch. Before surgery, 100 μl of the local anesthetic lurocaine (vetoquinol, France) was injected in the scalp region. Surgical procedure started with a longitudinal incision of 1.5 cm in length aimed to expose Bregma and Lambda. After ensuring the correct alignment of the head, two holes were drilled in the skull for electrode placement. Electrodes were: a glass recording electrode, inserted in the CA1 stratum radiatum, and a concentric stimulating bipolar electrode (Model CBARC50, FHC, ME, USA) placed in the CA3 region. Coordinates were as follows: CA1 stratum radiatum: A/P -1.5 , M/L -1.0 , DV 1.20 ; CA3: A/P -2.2 , M/L -2.8 , D/V -1.3 (20° insertion angle). The recording electrode (tip diameter = $1-2 \mu\text{m}$, $4-6 \text{M}\Omega$) was filled with a 2% pontamine sky blue solution in 0.5M sodium acetate. At first the recording electrode was placed by hand until it reached the surface of the brain and then to the final depth using a hydraulic micropositioner (Model 2650, KOPF instruments, CA, USA). The stimulation electrode was placed in the correct area using a standard manipulator. Both electrodes were adjusted to find the area with maximum response. *In vivo* recordings of evoked field excitatory postsynaptic potentials (fEPSPs) were amplified 1000 times and filtered (low-pass at 1Hz and high-pass 3000Hz) by a DAGAN 2400A amplifier (DAGAN Corporation, MN, USA). fEPSPs were digitized and collected on-line using a laboratory interface and software (CED 1401, SPIKE 2; Cambridge Electronic Design, Cambridge, UK). Test pulses were generated through an Isolated Constant Current Stimulator (DS3, Digitimer, Hertfordshire, UK) triggered by the SPIKE 2 output sequencer via CED 1401 and collected every 2 s at a 10 kHz sampling frequency and then averaged every 180 s. Test pulse intensities were typically between 40–250 μA with a duration of 50 μs . Basal stimulation intensity was adjusted to 30%–50% of the current intensity that evoked a maximum field response. All responses were expressed as percent from the average responses recorded during the 15 min before high frequency stimulation (HFS). HFS was induced by applying 3 trains of 100 Hz (1 s each), separated by 20 s interval. fEPSP were then recorded for a period of 30 or 39 min, depending on the experiment. C57BL/6 mice underwent this *in vivo* LTP induction immediately after the last exposition to odor-taste pairing, taste alone or odor alone of the behavioral protocol. Mice were exposed to control conditions (9 days of a daily presentation of water during 1 hr), “odor-taste pairings” (3 days of water habituation+6 days of odor-taste pairings as described above), “taste alone” (3 days of water habituation+6 days of taste presentations as described above) and “taste alone” (3 days of water habituation+6 days odor presentations as described above). On another group of GABA-CB₁R-KO mice, 5 weeks after the injection with the DIO-hM4Di virus into the

hippocampus, the following treatments were applied: CNO (2 mg/kg, i.p., 60-90 min before HFS) or vehicle (saline, i.p.). At the end of each experiment, the position of the electrodes was marked (recording area: iontophoretic infusion of the recording solution during 180 s at $-20\mu\text{A}$; stimulation area: continuous current discharge over 20 s at $+20\mu\text{A}$) and histological verification was performed *ex vivo* as shown in Supplementary information.

Ex vivo electrophysiology

Immediately after the last session of preconditioning (3 odor-taste pairings), or control procedure (either water, odor alone or taste alone, see above Western Blot Analysis) C57BL/6 mice were sacrificed by dislocation and the brain was immediately immersed in ice-cold oxygenated cutting solution containing in mM: 180 Sucrose, 26 NaHCO_3 , 12 MgSO_4 , 11 Glucose, 2.5 KCL, 1.25 NaH_2PO_4 , 0.2 CaCl_2 , oxygenated with 95% O_2 -5% $\text{CO}_2 \approx 300\text{mOsm}$. Parasagittal hippocampal slices (300 μm thick) were obtained using a vibratome (VT1200S, Leica, Germany) and transferred for 30min into a 34°C bath of oxygenated ACSF containing in mM: 123 NaCl, 26 NaHCO_3 , 11 Glucose, 2.5 KCL, 2.5 CaCl_2 , 1.3 MgCl_2 , 1.25 $\text{NaH}_2\text{PO}_4 \approx 305\text{mOsm}$. After a minimum of 1h recovery at room temperature (22 - 25°C), slices were transferred to a recording chamber in ACSF at 32°C .

Whole-cell recordings of IPSCs were made using a MultiClamp 700B amplifier (Molecular devices, UK) in CA1 pyramidal neurons voltage clamped at -70mV with a pipette (3-5 $\text{M}\Omega$) containing in mM: 130 KCl, 10 HEPES, 1 EGTA, 2 MgCl_2 , 0.3 CaCl_2 , 7 Phosphocreatin, 3 Mg-ATP, 0.3 Na-GTP; pH = 7.2; 290mOsm. Evoked IPSCs were performed by a monopolar stimulating patch pipette filled with ACSF in *stratum radiatum* in presence of NMDA and AMPA/Kainate receptor antagonists (50 μM D-APV and 10 μM NBQX). Miniature IPSCs were isolated in presence of the voltage-gated sodium channels blocker, tetrodotoxin (1 μM TTX) and collected for 5min.

Progressive DSIs were performed by depolarizing pyramidal neurons from -70mV to 0mV for 1 s, 3 s and 5 s. For every voltage step, DSIs' magnitude were measured as the average of 3 DSIs with 2min apart and represented the percentage of change between the mean of the 5 consecutive IPSCs preceding the depolarization and the first three IPSCs following the depolarization, with IPSCs evoked every 3 s.

iLTD were induced by High-Frequency-Stimulation (HFS) with 2 trains of 100 pulses at 100Hz with 20 s apart after a minimum of 10min of stable baseline, with IPSCs evoked every 20 s. The magnitude of iLTD was evaluated by the percentage of change between the mean of the 10min baseline with the percentage of responses averaged between 15 to 20min after HFS. Weak or no iLTD were assessed with a percentage of reduction less or equal to 15% whereas strong iLTD were estimated with a percentage of reduction higher than 15%.

Currents were filtered at 4kHz by a Digidata 1440A (Molecular devices, UK) and were analyzed using either Clampfit software (pClamp10) or Axograph for eIPSCs and mIPSCs, respectively.

QUANTIFICATION AND STATISTICAL ANALYSIS

Data collection

No statistical methods were used to pre-determine sample sizes, but they are similar to those reported in previous publications. Data collection and analysis were performed blind to the conditions of the experiment. All mice were assigned randomly to the different experimental conditions.

Statistical analyses

For the behavioral experiments, we performed the D'Agostino&Pearson normality test using the Prism 6 Software. When all groups of an experiment passed the normality test, ANOVA (Two-way or Three-way, where appropriate) analysis was performed and when interaction was significant Bonferroni's post hoc analysis was used. When the normality was rejected, the non-parametric Wilcoxon test was used for within-group comparisons and Mann-Whitney test for inter-group comparisons. For the *in vivo* electrophysiological and western blot experiments, data were analyzed by One-way ANOVA followed, when it was significant, by Dunnet's post hoc test. For *ex vivo* electrophysiological experiments, data were analyzed by unpaired t test for DSI or by Chi-square test for I-LTD. For detailed statistical analysis, see statistical [Tables S1](#) and [S2](#).