

Reduction of Carryover of Aflatoxin from Cow Feed to Milk by Addition of Activated Carbons

FABIO GALVANO,¹ AMEDEO PIETRI,² TERENCEO BERTUZZI,² GIORGIO FUSCONI,³ MARCO GALVANO,⁴ ANDREA PIVA,⁵ and GIANFRANCO PIVA²

¹Istituto di Zootecnica, Facoltà di Agraria, Università di Reggio Calabria, Piazza S. Francesco 7, 89061 Reggio Calabria; ²Istituto di Scienze degli Alimenti e della Nutrizione, Facoltà di Agraria, Università Cattolica del Sacro Cuore, Via Emilia Parmense 84, 29100 Piacenza; ³CERZOO, Centro Ricerche Zootecniche, S. Bonico, Piacenza; ⁴Dipartimento di Scienze Chimiche, Università di Catania, Viale A. Doria 6, 95125 Catania; and ⁵Facoltà di Medicina Veterinaria, Università de Bologna, Via Tolara di Spora 50, 40064 Ozzano Emilia, Bologna, Italy

(MS 95-168: Received 17 July 1995/Accepted 19 October 1995)

ABSTRACT

According to a double-reversal experimental design on 12 late-lactation Friesian cows the effect of two activated carbons (ACs) (CAC1 and CAC2) and a hydrated sodium calcium aluminosilicate (HSCAS) on carryover of aflatoxin B₁ (AFB₁) from feed to aflatoxin M₁ (AFM₁) in milk was determined. Cows were fed a basal diet containing AFB₁ naturally contaminated corn meal and copra. During week 1 cows were fed diets containing AFB₁ alone (11.28 µg of AFB₁/kg of feed); in week 2 the diets contained AFB₁ plus 2.0% sorbent; and in week 3 the diets again contained AFB₁ alone (13.43 µg of AFB₁/kg of feed). ACs reduced the analytical content of AFB₁ in the pelleted feed by from 40.6% to 73.6%, whereas reduction by HSCAS was 59.2%. The AFM₁ concentrations in milk in weeks 1 and 3 were higher than that in week 2. Decreases in the AFM₁ excreted in the milk by addition to feed of 2% of the sorbents ranged from 22% to 45%. CAC1 and HSCAS were significantly different from each other in reducing the AFM₁ concentration in milk (45.3% versus 32.5%); these reductions were significantly higher than that of CAC2 (22.0%). Carryover reduction by addition of CAC1 (50%) was significantly higher than that of HSCAS (36%). Addition of 2% CAC2 did not allow pelleting of feed because of the caking action of this carbon. The lower performance of CAC2 could be related to the unsuccessful pelleting. The addition of ACs did not influence feed intake, milk production, milk composition, or body weight. Our results suggest that ACs, high-affinity sorbents for AFB₁ in vitro, are efficacious in reducing AFB₁ carryover from cow feed to milk. Further in vivo investigations should establish lower amounts of ACs which can be efficacious.

Key words: Aflatoxins, milk, activated carbons, carryover, feed

Aflatoxins (AF) are a group of highly toxic secondary metabolic products of *Aspergillus flavus* and *Aspergillus parasiticus* that easily occur on feeds and foods. As AF are carcinogenic, teratogenic, and mutagenic to animals, including humans, detoxification of aflatoxin-contaminated foods

and feeds is one of the most serious problems of food hygiene.

Dairy cows fed aflatoxin B₁ (AFB₁)-contaminated diets secrete into milk amounts of the main hepatic AFB₁ metabolite, aflatoxin M₁ (AFM₁), as a function of AFB₁ intake and lactation stage (33). AFM₁ toxicity is similar to that of the original molecule as shown by trials carried out on ducks (25), rats (23), and trout (27), whereas its mutagenic potential seems to be lower (1). Recently AFB₁ and AFM₁ were categorized as class 1 and 2B (or probable) human carcinogens, respectively (26). As milk is one of the most important human foods and the main nutrient for growing young, who are notably vulnerable and potentially more sensitive than adults, numerous methods of detoxification have been tested (22).

The addition of sorbents to contaminated feeds seems to be the most efficacious strategy for preventing aflatoxicosis. Sorbents act as inabsorbable carriers that adsorb toxic molecules, thereby preventing or reducing their absorption in the intestinal tract. Some in vitro tests (20) showed that hydrated sodium calcium aluminosilicates (HSCASs) were particularly efficacious in binding aflatoxin. Harvey et al. (10) conducted a detoxification test by adding HSCAS to dairy cow feeds having 200 µg of AFB₁/kg and obtained a significant reduction in carryover, also confirming the efficacy of the treatment even at the low contamination levels frequently found in practice.

Pietri et al. (21) tested the effectiveness of a zeolite in reducing carryover of AFB₁ and did not detect any effect on carryover. Veldman (32) revealed a 33% and 0% carryover reduction by bentonite and HSCAS, respectively, and detected the formation of an inert, stable complex between bentonite and AFB₁ capable of preventing absorption in the intestine.

Although activated carbons (ACs) have been recognized as one of the most effective and nontoxic adsorbents (4), few studies about their employment as sorbents for AFB₁ are reported in the literature. Some authors observed low (5, 6) or no (14) ability of activated charcoal to reduce

* Author for correspondence. Tel: 965.682816; Fax: 965.682453.

AFB₁ toxic effects in chickens. In contrast, detoxification tests carried out on ducklings (28), mink (2), and chickens (13, 30) showed that activated charcoal was effective in reducing the toxic effects of AFB₁. These contrasts in the data can be ascribed to the type of activated charcoal used.

No data on the efficacy of ACs in reducing AFB₁ carryover into milk are available. Galvano et al. (12) evaluated AFB₁ affinity of ACs in vitro, revealing very high adsorption and higher affinity for AFB₁ than HSCAS. The aims of our investigation were to evaluate the ability of ACs and an HSCAS to reduce the carryover of AFB₁ from feed to milk and to establish whether AC could be added to contaminated feeds in practical conditions.

MATERIALS AND METHODS

Sorbents

The following finely pulverized sorbents were used: two commercial ACs (CAC1 and CAC2) previously tested and demonstrated to have an high affinity in vitro for AFB₁ (12), and an HSCAS demonstrated to reduce aflatoxin toxicity (10, 14, 20).

Experimental design

According to a double-reversal experimental design (34), 12 multiparous late-lactation (30 to 33 weeks) Friesian cows were randomly distributed into three groups, each receiving only one type of sorbent and used as its own control. Cows were fed a basal diet (15 kg of corn silage, 4 kg of triticale (*Triticum secale*) silage, 4 kg of alfalfa hay, 2 kg of dehydrated beet pulp) to which 5 kg of pelleted feed containing AFB₁ naturally contaminated corn meal and copra was added. Compound feed samples of each dietary treatment were collected during the experimental period to determine the AFB₁ content.

The experimental period was divided into three weeks: during week 1, cows were fed diets containing AFB₁ alone; in week 2 the diets contained AFB₁ plus 2.0% sorbent, and in week 3 the diets again contained AFB₁ alone. AFM₁ excretion into the milk was determined only on days 3 to 7 of each week. Daily morning and evening milk samples from each cow were collected and mixed in proportion to morning and evening milk production. These samples were again combined in proportion to the daily milk production of each cow to constitute a representative bulk milk sample of each group. The AFM₁ concentration in the milk from cows fed sorbents were compared with those when sorbents were not fed. Moreover, a comparison between the three sorbents was carried out.

Methods of analysis

Determination of AFB₁ in feed was carried out according to the method of Howell and Taylor (11) as regards extraction and column cleanup; 250 µl of the purified extract were treated to convert AFB₁ to AFB₂A by trifluoroacetic acid-catalyzed hydration according to Cohen and Lapointe (3).

AFM₁ in milk was extracted by immunoaffinity column (Easi-Extract[®], Biocode Ltd, University Road, Heslington, York, UK) according to the method reported by Mortimer et al. (17). For both AFB₁ and AFM₁, the final extract was evaporated under a nitrogen stream and finally redissolved in the mobile phase (500 µl) and filtered (0.45-µm pore size); 50 µl of extract were usually injected.

Determination of AFB₁ in feed and AFM₁ in milk were carried out by high-performance liquid chromatography (HPLC) under the following experimental conditions. A Perkin-Elmer Series 4 HPLC (Perkin-Elmer Corp., Norwalk, CT, USA) equipped with ISS 100 sampling system and 1020 computing integrator was used. The

detector was a Perkin-Elmer LS 4 fluorescence spectrophotometer with excitation and emission wavelengths of 365 nm (slit 15 nm) and 440 nm (slit 20 nm), respectively. The stationary phase was a LiChrosorb RP-18 (125 by 4 mm (i.d.) column) 5 µm (Merck, Darmstadt, Germany).

The mobile phase for AFB₁ determination was acetonitrile (A) plus water (B). The program was 20A plus 80B to 50A plus 50B, 0 to 12 min; 100A plus 0B, 13 to 14 min; and 20A plus 80B, 15 to 21 min. The flow rate was 1.5 ml/min. Retention time for AFB₂A was 8.4–8.5 min. Mobile phase for AFM₁ determination was isocratic, acetonitrile-water (28:72). The flow rate was 1.5 ml/min. The retention time for AFM₁ was 5.4 to 5.5 min.

Statistics

AFB₁ and AFM₁ data are presented as means ± SEM and *P* < 0.05 values were considered to be significant according to the Duncan multiple range test (7).

RESULTS AND DISCUSSION

Owing to the addition of 2% AC the feeds become black, but no effects on feed intake were revealed, as the diets were entirely consumed. The addition of 2% CAC2 did not allow pelleting of feed because of the caking action of this carbon.

AFB₁ reduction in feed

The addition of ACs to the feed achieved a substantial reduction in the analytically detected AFB₁. As reported in Figure 1, the addition of CAC1 and CAC2 reduced detectable AFB₁ from 11.28 µg/kg to 2.68 µg/kg and 6.70 µg/kg respectively, whereas HSCAS reduced AFB₁ to 4.60 µg/kg.

Phillips et al. (20) in in vitro studies showed that HSCAS had a great affinity for AFB₁, whereas others (32, 35) reported a reduction of detectable AFB₁ in feed by the addition of bentonite. The authors explained these findings by the formation of a stable complex between the sorbents




FIGURE 1. Reduction of AFB₁ in feed and AFM₁ in milk as an effect of the addition of sorbents. Each bar represents the mean (*n* = 4 for AFB₁; *n* = 10 for AFM₁); error bars denote standard error of the mean. Different letters on the tops of bars mean significantly different values, *P* < 0.05.

and AFB₁, resistant against solvent extraction. Since addition of ACs reduced the analytically recovered AFB₁ from the feed by 40.6% to 73.6%, the formation of an analogous inert, stable complex, AC-AFB₁, resistant to solvent extraction (chloroform), must occur.

Under the conditions of the present study, CAC1 showed greater reduction of the analytically recovered AFB₁ from feed than HSCAS (73.6% versus 59.2%) (Figure 1). This agrees with our previous study (12) in which *in vitro* affinity for AFB₁ of ACs compared to HSCAS was tested. In contrast, the performance of CAC2 was far from that expected. As pelleting of feed probably improves the adsorption of AFB₁ by ACs, the lower reduction of analytically recovered AFB₁ from feed with CAC2 could be related to the unsuccessful pelleting.

AFM₁ reduction in milk

The AFM₁ content of milk related to AFB₁ intake was lower than values reported by Munksgaard et al. (18). Veldman et al. (33) reported that the average daily AFB₁ intake for dairy cows has to be below 40 µg/day to produce milk with less than 0.05 µg of AFM₁/kg (European Communities tolerance). In the present study, despite the average AFB₁ intake of cows being over 55 µg/day, the AFM₁ content of milk, with or without addition of sorbents was substantially below the EC tolerance level.

Mean values for AFM₁ of week 2 (sorbents present) were significantly lower than those of weeks 1 and 3 (Table 1). In agreement with the observations of Harvey et al. (10), the absence of sorbents during week 3 allowed AFM₁ concentration in milk to reach the same or greater values than in week 1. Nevertheless, these results could be also related to the increase of AFB₁ intake in week 3. In contrast, AFM₁ secretion showed the same values from week 1 to week 3 because of decreased milk yield. However, data pooled for the 3 test weeks showed that both the ACs and the HSCAS were effective in reducing AFM₁ concentration in milk. In agreement with the results of analytic recovery of AFB₁ from feed, addition of CAC1 to feed caused a significantly higher decline of AFM₁ concentration in milk than HSCAS (45.3% vs 32.5%); furthermore, CAC2, with a

22.0% reduction, was less effective than CAC1 and HSCAS in reducing AFM₁ concentration in milk (Figure 1).

Carryover from feed to milk

Carryover of AFM₁ from feed to milk in weeks 1 and 3 (controls) ranged from 0.45% to 0.55%, whereas in week 2 (sorbents added) it ranged from 0.27% to 0.41%. Carryover data from the present study were similar to those (0.14% to 6.2%) commonly reported for dairy cows (8–10, 15, 16, 18, 19, 24, 29, 31–33). Nevertheless, carryover values related to milk yield in both control and treatment weeks were quite low if compared to the values observed by other authors (31–33). This could be partially explained by the late lactation stage of the cows, in agreement with the observations of Lafont et al. (16) and Veldman et al. (33). Moreover, these latter authors (33) reported that a wide variability of carryover could be ascribed to animal factors related to the conversion of AFB₁ to AFM₁ by the mixed-function oxidase system.

The HSCAS was confirmed as a protective agent against AFB₁. Under our experimental conditions, 2% HSCAS addition reduced carryover by 36%. This reduction was lower than those reported in studies on dairy goats (2% HSCAS, 60% carryover reduction) (10), (1, 2, and 4% HSCAS addition, 52, 82, and 87% carryover reduction, respectively) (29), and on dairy cows (1% HSCAS, 44% carryover reduction) (10), but higher if compared to the studies on dairy cows of Harvey et al. (10) (0.5% HSCAS, 24% carry-over reduction), and Veldman (32), who reported no carryover reduction by addition of 1% HSCAS. Nevertheless, differences in species, dosages of HSCAS and AFB₁, and duration of AFB₁ exposure should be considered when comparing published results.

We are aware of no reports on carryover reduction by addition of ACs. CAC1 was more efficient at reducing carryover than HSCAS (50% versus 36%). In contrast, CAC2 (27%) was less effective than HSCAS in reducing carryover. This latter result, as well as the lower reduction of detectable AFB₁ in feed and AFM₁ in milk, could be ascribed to the unsuccessful pelleting of feed. Our data

TABLE 1. Reduction in carryover of AFB₁ in feed to AFM₁ in milk over a 3-week period in three groups of cows with or without sorbents added to the feed

Week (group)	Sorbent in feed	AFB ₁ intake (µg/day)	Milk yield (kg/day)	AFM ₁ in milk		Carryover AFB ₁ to AFM ₁ (%)	Reduction of carryover (%)
				(ng/l) ^a	(µg/day)		
1 (1)	none	56.40	19.40	15.52 ± 0.73A	0.30	0.53	NA ^b
1 (2)	none	56.40	19.34	15.88 ± 0.44B	0.31	0.54	NA
1 (3)	none	56.40	19.66	15.83 ± 0.65B	0.31	0.55	NA
2 (1)	HSCAS	56.40	18.10	10.48 ± 0.33B	0.19	0.34	36
2 (2)	CAC1	56.40	17.72	8.68 ± 0.22C	0.15	0.27	50
2 (3)	CAC2	56.40	18.78	12.35 ± 0.70C	0.23	0.41	27
3 (1)	none	67.15	17.58	17.25 ± 0.73A	0.30	0.45	NA
3 (2)	none	67.15	17.66	17.83 ± 0.48A	0.31	0.47	NA
3 (3)	none	67.15	17.44	18.41 ± 0.69A	0.32	0.45	NA

^a Mean ± SEM; n = 10. Within each group (see first column) of cows, values followed by different letters are statistically different (P < 0.05).

^b NA, not applicable.

suggest that the complex AC-AFB₁ should decrease the bioavailability of AFB₁ for absorption in the intestine and could reduce carryover up to 50%. Measure of fecal AFB₁ excretion would provide additional evidence. Adding a lower amount (from 0.1% to 0.5%) may attenuate both pelleting difficulties and the color change of the feed. Since addition of ACs did not show effects on milk yield and composition or body weight (data not shown) our study demonstrates that ACs could be added to feed in practical conditions. As regards toxic effects related to addition of ACs, no obvious adverse effects were observed, but our data do not allow us to affirm that there would never be adverse effects on health status. Undesired preferential adsorption of useful organic molecules (e.g., vitamins or minerals) should be investigated in long-term studies.

In summary, our results demonstrate that ACs, high-affinity sorbents for AFB₁ in vitro, could reduce the AFM₁ content of milk to safe levels and are potential protective agents against aflatoxicosis at low contamination levels. Further in vivo investigations should establish lower amounts of ACs added to feed which can be efficacious. Moreover, studies in progress in our laboratory on the aflatoxin binding process could allow functional improvements to the adsorption ability of ACs.

ACKNOWLEDGMENTS

This study was supported by Regione Emilia Romagna, Assessorato Agricoltura, Piano Stralcio 1994, LR 19/88, prog. no. 51. The authors would like to express their gratitude to Mr. Vittorio Galafaro for his technical assistance.

REFERENCES

- Barnes, J. M. 1970. Aflatoxin as health hazard. *J. Appl. Bacteriol.* 33:285-298.
- Bonna, R. J., R. J. Aulerich, S. J. Bursan, R. H. Poppenga, W. E. Braselton, G. L. Watson. 1991. Efficacy of hydrated sodium calcium aluminosilicate and activated charcoal in reducing the toxicity of dietary aflatoxin to mink. *Arch. Environ. Contam. Toxicol.* 20:441-447.
- Cohen, H., and M. Lapointe. 1981. High pressure liquid chromatographic determination and fluorescence detection of aflatoxins in corn and dairy feeds. *J. Assoc. Off. Anal. Chem.* 64:1372-1376.
- Cooney, D. O. 1980. Activated charcoal: antidotal and other medical uses. Marcel Dekker Inc., New York.
- Dalvi, R. R., and A. Ademoyero. 1984. Toxic effects of aflatoxin B₁ in chickens given feed contaminated with *Aspergillus flavus* and reduction of the toxicity by activated charcoal and some chemical agents. *Avian Dis.* 28:61-69.
- Dalvi, R. R., and C. McGowan. 1984. Experimental induction of chronic aflatoxicosis in chickens by purified aflatoxin B₁ and its reversal by activated charcoal, phenobarbital, and reduced glutathione. *Poult. Sci.* 63:485-491.
- Duncan, D. B. 1955. Multiple range and multiple F tests. *Biometrics* 11:1-42.
- Fremy, J. M., J. P. Gautier, M. P. Henry, C. Terrier, and C. Calet. 1987. Effects of ammoniation on the "carry-over" of aflatoxins into bovine milk. *Food Addit. Contam.* 5:39-44.
- Frobish, R. A., B. D. Bradley, D. D. Wagner, P. E. Long-Bradley, and H. Hairstone. 1986. Aflatoxin residues in milk of dairy cows after ingestion of naturally contaminated grain. *J. Food Prot.* 49:781-785.
- Harvey, R. B., T. D. Phillips, J. A. Ellis, L. F. Kubena, W. E. Huff, and H. D. Petersen. 1991. Effect on aflatoxin M₁ residues in milk by addition of hydrated sodium calcium aluminosilicate to aflatoxin contaminated diets of dairy cows. *Am. J. Vet. Res.* 52:1556-1559.
- Howell, M. V., and P. W. Taylor. 1981. Determination of aflatoxins, ochratoxin A, and zearalenone in mixed feeds, with detection by thin layer chromatography or high performance liquid chromatography. *J. Assoc. Off. Anal. Chem.* 64:1356-1363.
- Galvano, F., A. Pietri, B. Fallico, T. Bertuzzi, S. Scirè, M. Galvano, and R. Maggiore. 1996. Activated carbons: in vitro affinity for aflatoxin B₁ and relation of adsorption ability to physicochemical parameters. *J. Food Prot.* 59:545-550.
- Jindal, N., S. K. Mahial, and N. K. Mahajan. 1994. Toxicity of aflatoxin B₁ in broiler chicks and its reduction by activated charcoal. *Res. Vet. Sci.* 56:37-40.
- Kubena, L. F., R. B. Harvey, D. E. Corrier, T. D. Phillips, and W. E. Huff. 1990. Diminution of aflatoxicosis in growing chickens by dietary addition of hydrated sodium calcium aluminosilicate. *Poult. Sci.* 69:727-735.
- Kulczycki, J., M. Lesiak, and K. Polkowski. 1985. Effect of subtoxic doses of aflatoxins fed to cow on serum and milk composition. *Bull. Vet. Inst. Pulawy* 29:142-150.
- Lafont, P., J. Lafont, S. Mousset, and C. Frayssinet. 1980. Étude de la contamination du lait de vache lors de l'ingestion de faibles quantités d'aflatoxine. *Ann. Nutr. Aliment.* 34:699-708.
- Mortimer, D. N., J. Gilbert, and M. J. Shepherd. 1987. Rapid and highly sensitive analysis of aflatoxin M₁ in liquid and powdered milk using an affinity column cleanup. *J. Chromatogr.* 407:393-398.
- Munksgaard, L., J. Larsen, H. Werner, P. E. Anderson, and B. T. Viuf. 1987. Carryover of aflatoxin from cows' feed to milk and milk products. *Milchwissenschaft* 42:165-167.
- Petterson, H., J. Bertilsson, and O. Wennberg. 1989. Carry-over of aflatoxin from dairy cattle feed to milk, p. 97-102. *In Proc. Healthy animals, safe foods, healthy man. World Association of Veterinary Food Hygienists Xth (Jubilee), International Symposium in Stockholm 2-7 July.*
- Phillips, T. D., L. F. Kubena, R. B. Harvey, D. R. Taylor, and N. D. Heidelbaugh. 1988. Hydrated sodium calcium aluminosilicate: a high affinity sorbent for aflatoxin. *Poult. Sci.* 67:243-247.
- Pietri, A., G. Fusconi, P. Blasi, and M. Moschini. 1993. Azione sequestrante di una zeolite nei confronti di radionuclidi e aflatoxina in diete per ruminanti e monogastrici. Xth Congresso ASPA, Bologna, Italy, 31 maggio-6 giugno. *Proc.* 141-146.
- Piva, G., F. Galvano, A. Pietri, and A. Piva. 1995. Detoxification methods of aflatoxins. A review. *Nutr. Res.* 15:689-715.
- Pong, R. S., and G. N. Wogan. 1971. Toxicity and biochemical and fine structural effects of synthetic aflatoxin M₁ and B₁ in rat liver. *J. Natl. Cancer Inst.* 47:585-590.
- Price, R. L., J. H. Paulson, O. G. Lough, C. Gingg, and A. G. Kurtz. 1985. Aflatoxin conversion by dairy cattle consuming naturally contaminated whole cottonseed. *J. Food Prot.* 48:11-15.
- Purchase, I. F. H. 1967. Acute toxicity of aflatoxins M₁ and M₂ in one-day-old ducklings. *Food Cosmetol. Toxicol.* 5:339-342.
- Rothschild, L., Jr. 1992. IARC classes AFB₁ as class 1 human carcinogen. *Food Chem. News* 34:62-66.
- Sinnhuber, R. O., D. J. Lee, J. K. Wales, M. K. Landers, and A. C. Keyl. 1974. Aflatoxin M₁, a potent liver carcinogen for rainbow trout. *Fed. Proc.* 29:568.
- Sjamsul, B., P. Zahari, and H. Hamid. 1990. The use of activated charcoal on the prevention of aflatoxicosis in duckling (in Indonesian). *Penyakit Hewan (Bogor)* 22:122-127.
- Smith, E. E., T. D. Phillips, R. B. Harvey, L. F. Kubena, J. Thompson, and G. Newton. 1994. Dietary hydrated sodium calcium aluminosilicate reduction of aflatoxin M₁ residues in dairy goat milk and effects on milk production and components. *J. Anim. Sci.* 72:677-682.
- Umesh, D., V. N. Rao, and H. C. Joshi. 1991. Chemoprophylactic effect of certain drugs against chronic aflatoxicosis in chicken. *Ind. J. Vet. Med.* 11:58-59.
- Van Egmond, H. P. 1989. Carry-over of aflatoxin B₁ into aflatoxin M₁ in milk. *In H. P. van Egmond (ed.), Mycotoxins in dairy products.* Elsevier Applied Science Publishers, Amsterdam.
- Veldman, A. 1992. Effect of sorbentia on carry-over of aflatoxin from cow feed to milk. *Milchwissenschaft* 47:777-780.
- Veldman, A., A. C. Meijst, G. J. Borggreve, and J. J. Heeres-van der Tol. 1992. Carry-over of aflatoxin from cows' food to milk. *Anim. Prod.* 55:163-168.
- Winer, B. J. 1971. Statistical principles in experimental design, 2nd ed., p. 514-599. McGraw-Hill Book Co., New York.
- Winfree, R. A., and A. Allred. 1992. Bentonite reduces measurable aflatoxin B₁ in fish feed. *Prog. Fish-Cult.* 54:157-162.