

Dietary Inflammatory Index and Risk of Bladder Cancer in a Large Italian Case-control Study



Nitin Shivappa, James R. Hébert, Valentina Rosato, Marta Rossi, Massimo Libra, Maurizio Montella, Diego Serraino, and Carlo La Vecchia

OBJECTIVE	To evaluate the association between diet in relation to its inflammatory property and bladder cancer (BC) risk.
METHODS	In this study we explored the association between the dietary inflammatory index (DII) and BC risk in an Italian case-control study conducted between 2003 and 2014. Cases were 690 patients with incident and histologically confirmed BC from 4 areas in Italy. Controls were 665 cancer-free subjects admitted to the same network of hospitals as cases for a wide spectrum of acute, non-neoplastic conditions. The DII was computed based on dietary intake assessed using a reproducible and valid 80-item food frequency questionnaire. Odds ratios (OR) were estimated through logistic regression models adjusting for age, sex, total energy intake, and other recognized confounding factors.
RESULTS	Subjects in the highest quartile of DII scores (ie, with a more pro-inflammatory diet) had a higher risk of BC compared to subjects in the lowest quartile (ie, with an anti-inflammatory diet) ($OR_{\text{Quartile4vs1}} = 1.97$; 95% [confidence interval], 1.28, 3.03; P trend = .003). Stratified analyses produced stronger associations between DII and BC risk among females ($OR_{\text{Quartile4vs1}} = 5.73$; 95% CI = 1.46, 22.44), older ≥ 65 years ($OR_{\text{Quartile4vs1}} = 2.45$; 95% CI = 1.38, 4.34), subjects with higher education ≥ 7 years ($OR_{\text{Quartile4vs1}} = 2.22$; 95% CI = 1.27, 3.88), and never smokers ($OR_{\text{Quartile4vs1}} = 4.04$; 95% CI = 1.51, 10.80).
CONCLUSION	A pro-inflammatory diet as indicated by higher DII scores is associated with increased BC risk. UROLOGY 100: 84–89, 2017. © 2016 Elsevier Inc.

Financial Disclosure: Dr. James R. Hébert owns controlling interest in Connecting Health Innovations LLC (CHI), a company planning to license the right to his invention of the dietary inflammatory index (DII) from the University of South Carolina to develop computer and smartphone applications for patient counseling and dietary intervention in clinical settings. Dr. Nitin Shivappa is an employee of CHI. The remaining authors declare that they have no relevant financial interests.

Funding Support: This study was supported by the Italian Foundation for Research on Cancer (FIRC) and by the Italian Ministry of Health, General Directorate of European and International Relations (Grant No. 8942). Drs. Shivappa and Hébert were supported by grant number R44DK103377 from the United States National Institute of Diabetes and Digestive and Kidney Diseases.

From the Cancer Prevention and Control Program, University of South Carolina, Columbia, SC; the Department of Epidemiology and Biostatistics, Arnold School of Public Health, University of South Carolina, Columbia, SC; the Department of Family and Preventive Medicine, University of South Carolina School of Medicine, Columbia, SC; the Connecting Health Innovations LLC, Columbia, SC; the Department of Clinical Sciences and Community Health, Università degli Studi di Milano, Milan, Italy; the Laboratory of Translational Oncology & Functional Genomics, Department of Biomedical and Biotechnological Sciences, Università di Catania, Catania, Italy; the Unit of Epidemiology, "Fondazione G. Pascale", Istituto Nazionale Tumori, Naples, Italy; and the Unit of Cancer Epidemiology, CRO Aviano National Cancer Institute IRCCS, Aviano, Italy

Address correspondence to: Nitin Shivappa, Ph.D., South Carolina Statewide Cancer Prevention and Control Program, 915 Greene Street, Columbia, SC 29205. E-mail: shivappa@mailbox.sc.edu

Submitted: April 27, 2016, accepted (with revisions): September 14, 2016

Bladder cancer (BC) is the ninth most common cancer worldwide,¹ the sixth most common cancer worldwide in men, and the 19th most common cancer in women, with approximately 38,000 deaths per year in the European Union.^{1,2} Major recognized risk factors for BC are tobacco smoking and past occupational exposure to aromatic amines.²

Considerable evidence has been gathered over the past few years linking increased cancer risk with chronic inflammation, and several clinical and experimental studies have linked tumor progression with the upregulation of pro-inflammatory molecules, especially during late stages of the disease.³ In addition to chronic inflammation, intrinsic properties of premalignant cells and other factors promote tumor initiation and promotion.⁴ Substantial evidence from developed countries shows the role of chronic inflammation in urological cancers, including BC.⁵

Dietary components such as fruits, vegetables, and coffee have been studied in relation to incident BC in various studies.^{6,7} Dietary patterns characterizing the

Western-type diet, that is, including high consumption of red meat, high-fat dairy products, and refined grains, have been associated with higher levels of C-reactive protein, interleukin-6 (IL-6), and fibrinogen.⁸ On the other hand, the Mediterranean diet—characterized by a high consumption of whole grains, fruit and green vegetables, fish, and olive oil, a low consumption of red meat and butter, and a moderate alcohol and dairy products consumption—has been associated with lower levels of inflammation.⁹ Despite the circumstantial evidence, the possible relation between inflammation deriving from dietary exposure and BC risk has been investigated widely.

The literature-derived dietary inflammatory index (DII) was developed to assess the inflammatory potential of an individual's diet.¹⁰ Higher DII scores indicate increasing inflammatory potential of diet. The DII has been validated with various inflammatory markers, including C-reactive protein,¹¹ IL-6,¹² and tumor necrosis factor.¹²

The DII has been associated with a variety of cancers, including urological cancers, such as prostate,^{13,14} colorectal,¹⁵ esophageal,¹⁶ and breast.¹⁷ DII and urothelial cancer association has been explored in 1 Australian prospective study before.¹⁸ This large case-control study conducted in Italy^{19,20} provides us the opportunity to examine the association between DII scores and BC risk. Our working hypothesis is that the increasing inflammatory potential of diet is associated with increased risk of BC.

METHODS

Design and Participants

Between 2003 and 2014, we conducted a case-control study on BC in 4 Italian centers: Aviano and Milan (in Northern Italy), and Naples and Catania (in Southern Italy).^{19,20} Controls were patients admitted to the same network of hospitals as cases for a wide spectrum of acute, non-neoplastic conditions unrelated to tobacco smoking and alcohol consumption or long-term diet modification (eg, bronchitis, myocardial infarction). Overall, 28.9% of controls were admitted for traumas, 22.1% for nontraumatic orthopedic disorders, 39.3% for acute surgical conditions, and 9.8% for other miscellaneous illnesses. The study protocol was approved by the Institutional Ethics Committees of each of the hospitals, and all subjects signed an informed consent.

Centrally trained and supervised interviewers collected information on socio-demographic characteristics, anthropometric measures, lifestyle habits, including tobacco smoking, and history of selected diseases during their hospital stay using a structured questionnaire. Each subject's usual diet during the 2 years prior to cancer diagnosis (for cases) or hospital admission (for controls) was assessed using an interviewer-administered food frequency questionnaire (FFQ), consisting of 80 items on food and 15 (alcoholic and non-alcoholic) on beverages. Subjects were asked to indicate the weekly average frequency of consumption of each dietary item; intakes lower than once a week, but

at least once a month, were coded as 0.5 per week. Nutrient and total energy intake was determined using an Italian food composition database.²¹ The FFQ showed a satisfactory validity²² and reproducibility.²³

Details of the steps involved in DII calculation is described elsewhere.^{10,24} To compute the DII score, dietary information for each study participant was first linked to the regionally representative database that provided a robust estimate of a mean and a standard deviation (SD) for each of the 45 parameters (ie, food, nutrients, and other food components) considered in the DII definition.¹⁰ These parameters were then used to derive the subject's exposure relative to the standard global mean as a z -score, derived by subtracting the mean of the regionally representative database from the amount reported, and dividing this value by the parameter's SD. To minimize the effect of "right skewing," this value was converted to a centered percentile score, which was computed by doubling the raw percentile score and then subtracting 1. This score was then multiplied by the respective food parameter effect score (derived from a literature review on the basis of 1943 articles).^{10,24} All of these food parameter-specific DII scores were then summed to create the overall DII score for every subject in the study. Higher scores indicate a pro-inflammatory diet whereas lower scores indicate a more anti-inflammatory diet. The DII computed on this study's FFQ includes data on 31 of the 45 possible food parameters comprising the DII: carbohydrates, proteins, fats, alcohol, fibers, cholesterol, saturated fatty acids, monounsaturated fatty acids, polyunsaturated fatty acids, omega 3, omega 6, niacin, thiamin, riboflavin, vitamin B6, iron, zinc, vitamin A, vitamin C, vitamin D, vitamin E, folic acid, beta carotene, anthocyanidins, flavan-3-ols, flavonols, flavanones, flavones, isoflavones, caffeine, and tea.

The DII was analyzed as continuous and by quartiles computed among controls. Odds ratios (ORs) and the corresponding 95% confidence intervals (CIs) were estimated using unconditional logistic regression models including terms for age (quinquennial), sex, year of interview, study center, education (<7; 7-11; ≥ 12 years), tobacco smoking (never; former; current: <15; 15-24, ≥ 25 cigarettes/day), and total energy intake (quintiles of the distribution of the controls). We further adjusted for the most frequent weekly occupational exposure to carcinogens in our study (ie, aromatic amines, ammonia, sulfuric acid or hydrochloric, tar, pitch, asphalt, and soot, mineral oils and lubricants, gasoline, diesel engine fumes, foundry fumes, aluminum powder, concrete, lead, benzene, and other solvents) and selected diseases (ie, diabetes, hypercholesterolemia, and obesity); however, the estimates did not change materially, and thus these factors were not included in the final, most parsimonious, model. Tests for linear trend were performed using the median value within each quartile as an ordinal variable. Stratified analyses were carried out according to sex, age (<65, ≥ 65 years), education (<7, ≥ 7 years), and tobacco smoking (never smokers and ever smokers), and heterogeneity across strata was tested computing the difference in the $-2 \log$ likelihood of the models with and without

the interaction terms. Statistical analyses were performed using SAS 9.3 (SAS Institute Inc., Cary, NC).

RESULTS

Cases were 690 patients aged 25-80 years (median 67 years) with incident BC (mostly transitional cell carcinoma), admitted to major general hospitals in the study areas. Nearly all BCs ($n = 642$, 93.0%) were confirmed histologically on tumor tissue specimen from biopsy or surgery, whereas 3 additional cases were confirmed by cytology. Overall, 460 patients (66.7%) were nonmuscle-invasive (NMIBC, TNM pTis/Ta/T1) and 159 (23.0%) were muscle-invasive (MIBC, other T) BCs; 307 (44.5%) were well differentiated or low grade (G1, G2), and 312 (45.2%) were poorly differentiated or high grade (G3, G4). Stage and grade were undefined for 71 (10.3%) patients. The controls were 665 patients frequency-matched to cases by study center, sex, and 5-year age group (median age 66 years). The distribution of BC cases and controls according to selected factors is given in Table 1. Cases and controls had a similar distribution by study center, sex, age, and education. Cases were more frequently smokers than controls.

The mean DII value was -0.63 (SD = 1.94, range -5.42 to $+4.48$) among cases and -0.93 (SD = 2.00, range -5.94 to $+4.58$) among controls, indicating a more pro-inflammatory diet for cases. The characteristics of control

Table 1. Distribution of 690 cases of bladder cancer and 665 controls according to center, sex, age and other selected variables (Italy, 2003-2014)

	Cases		Controls	
	No.	%	No.	%
Center				
Pordenone	242	35.1	250	37.6
Milan	241	34.9	238	35.8
Naples	129	18.7	100	15.0
Catania	78	11.3	77	11.6
Sex				
Men	595	86.2	561	84.4
Women	95	13.8	104	15.6
Age (years)				
<60	148	21.5	178	26.8
60-64	107	15.5	119	17.9
65-69	164	23.8	147	22.1
70-74	155	22.5	124	18.7
≥75	116	16.8	97	14.6
Education (years)*				
<7	292	42.4	273	41.1
7-11	224	32.5	215	32.3
≥12	173	25.1	177	26.6
Tobacco smoking*				
Never smokers	96	14.1	237	35.6
Ex-smokers	310	45.5	284	42.7
Current smokers				
<15 cigarettes/day	79	11.6	53	8.0
15-24 cigarettes/day	127	18.7	68	10.2
≥25 cigarettes/day	69	10.1	23	3.5

* The sum does not add up to the total because of some missing values.

subjects across quartiles of DII are provided in Table 2. There were significant differences in tobacco smoking across quartiles of DII. Subjects in the highest quartile of DII were more likely to be current smokers.

Table 3 shows OR of BC and 95% CI according to continuous and quartiles of DII score. When used as a continuous exposure, one-unit increase in DII score (corresponding to 9% change in DII in the current study) was associated with increased odds for BC (OR_{Continuous} = 1.11, 95% CI = 1.03, 1.20), and when used as quartiles, subjects in the highest DII score quartiles had 97% increased odds for BC compared to subjects in the lowest quartile (OR_{Quartile4vs1} = 1.97, 95% CI = 1.28, 3.03; P trend = .003).

Table 4 shows ORs of BC in strata of selected covariates. A stronger association was observed among females (OR_{Quartile4vs1} = 5.73, 95% CI 1.46, 22.44) than males (OR_{Quartile4vs1} = 1.83, 95% CI 1.14, 2.91) (P value for interaction .002), although the former estimate was based on many fewer cases and controls (24 cases and 28 controls in the highest quartile). Stronger associations—although in the absence of significant heterogeneity (P values $>.10$)—were observed among subjects aged ≥ 65 years (OR_{Quartile4vs1} = 2.45, 95% CI 1.38, 4.34), education ≥ 7 years (OR_{Quartile4vs1} = 2.22, 95% CI 1.27, 3.88), and never smokers (OR_{Quartile4vs1} = 4.04, 95% CI 1.51, 10.80). We also performed analysis separated by NMIBC and MIBC patients, and found DII to be associated with both NMIBC (OR_{Quartile4vs1} = 1.92, 95% CI: 1.19, 3.08) and MIBC (OR_{Quartile4vs1} = 2.42, 95% CI: 1.18, 4.93). When we performed separated analysis by grade, we found DII to be associated with both grade 1 BC (OR_{Quartile4vs1} = 1.79, 95% CI: 1.04, 3.09) and grade 2 BC (OR_{Quartile4vs1} = 2.13, 95% CI: 1.24, 3.67).

DISCUSSION

This large multicentric Italian case-control study is the first attempt to investigate the association between inflammatory potential of diet and BC. We observed a 2-fold excess risk of BC among individuals with a pro-inflammatory diet as expressed by high DII scores.

High consumption of fruits and vegetables may reduce BC risk whereas high coffee consumption is inconsistently related to the increased risk of BC.^{6,7} Fruits and vegetables are rich in various vitamins such as thiamin, niacin, vitamin E, and carotenoids, which have anti-inflammatory effects and therefore contribute to lower DII scores.¹⁰ In a large European cohort study, consumption of carotenoids, such as beta-carotene, was found to reduce the risk of BC.²⁵ In a prospective study conducted in Australia, increasing DII was observed to have a nonsignificant association with urothelial cancer.¹⁸

We observed a stronger association between DII scores and BC among nonsmokers in this study. It is known that tobacco is a strongly pro-inflammatory agent.²⁶ It could be that among smokers, or people exposed occupationally to pro-inflammatory carcinogens (or both), the pro-inflammatory state induced by smoking, or occupational

Table 2. Participants' characteristics across quartiles of dietary inflammatory index (DII) among 665 controls (Italy, 2003-2014)

Characteristics	DII Quartiles				P Value*
	-5.94, -2.41 No. (%)	-2.40, -0.92 No. (%)	-0.91, 0.41 No. (%)	0.42, 4.58 No. (%)	
Sex					.15
Male	139 (83.7)	135 (80.8)	148 (89.7)	139 (83.2)	
Female	27 (16.3)	32 (19.2)	17 (10.3)	28 (16.8)	
Age (years)					.66
<60	37 (22.3)	41 (24.6)	52 (31.5)	48 (28.7)	
60-64	33 (19.9)	30 (18.0)	27 (16.4)	29 (17.4)	
65-69	44 (26.5)	36 (21.6)	38 (23.0)	29 (17.4)	
70-74	28 (16.9)	34 (20.4)	29 (17.6)	33 (19.8)	
≥75	24 (14.5)	26 (15.6)	19 (11.5)	28 (16.8)	
Education (years)					.29
<7	74 (44.6)	70 (41.9)	61 (37.0)	68 (40.7)	
7-11	53 (31.9)	49 (29.3)	65 (39.4)	48 (28.7)	
>11	39 (23.5)	48 (28.7)	39 (23.6)	51 (30.5)	
Tobacco smoking†					.007
Never smokers	63 (37.9)	68 (40.7)	54 (32.7)	52 (31.1)	
Ex-smokers	76 (45.8)	71 (42.5)	72 (43.6)	65 (38.9)	
Current smokers					
<15 cigarettes/day	17 (10.2)	10 (5.9)	13 (7.9)	13 (7.8)	
15-24 cigarettes/day	9 (5.4)	14 (8.4)	21 (12.7)	24 (14.4)	
≥25 cigarettes/day	1 (0.6)	4 (2.4)	5 (3.0)	13 (7.8)	

* P value for chi-square test.

† The sum does not add up to the total because of some missing values.

Table 3. Odds ratio (OR) of bladder cancer and corresponding 95% confidence intervals (CI) according to dietary inflammatory index (DII) among 690 cases and 665 controls (Italy, 2003-2014)

	DII Quartiles, OR (95% CI)				P Value for Trend	DII Continuous
	-5.94, -2.41	-2.40, -0.92	-0.91, 0.41	0.42, 4.58		
Cases/controls	124/166	180/167	179/165	207/167		690/665
Model 1*	1†	1.56 (1.11, 2.17)	1.64 (1.11, 2.35)	2.11 (1.41, 3.14)	<.0001	1.17 (1.08, 1.26)
Model 2†	1†	1.46 (1.02, 2.08)	1.46 (0.99, 2.14)	1.97 (1.28, 3.03)	.003	1.11 (1.03, 1.20)

* Adjusted for age, sex, year of interview, study center, and total energy intake.

† Reference category.

† Model 1 additionally adjusted for education (<7; 7-11; ≥12 years) and tobacco smoking (never; former; current: <15, 15-24, ≥25 cigarettes/day).

carcinogens, overwhelms the countervailing anti-inflammatory effects of diet²⁷; further research should be conducted to explore this mechanism. It is important to note, however, that heterogeneity across strata of tobacco smoking was nonsignificant.

A significant positive association between DII and BC risk was observed in both men and women; however, the association was stronger in women. Although women had a lower rate of smoking than men, tobacco use was accounted for in the analyses. Women also tend to have lower occupational exposures to workplace carcinogens,²⁸ and this may help to explain the much lower BC incidence in women than men. At this juncture, there is no simple and satisfactory explanation for the stronger effect of the DII seen in women.

One of the possible mechanisms through which the observed positive association between DII and BC could occur is through oxidative and nitrative DNA damage in stem cells caused by chronic inflammation.²⁹ Diet-related chronic inflammation plays a role in the upregulation of various

cytokines such as tumor growth factor-beta and IL-6 that, in turn, promote cell transformation, survival, proliferation of tumor cells, and metastasis.³⁰

This hospital-based case-control study shares some of the limitations of such designs. Potential recall bias is possible; however, the comparability of recall between cases and controls was improved by interviewing all subjects in a hospital setting. As for selection bias, the study was hospital based, but cases and controls were interviewed in the same hospital setting, came from comparable catchment areas, and their participation was comparable and almost complete. In addition, awareness about any particular dietary hypothesis in BC etiology was limited in the Italian population. Among the strengths of this study are its large size and the use of a reproducible²³ and valid²² FFQ. Also, the DII score, which takes into account both pro- and anti-inflammatory food parameters that characterize human diet, more accurately reflects the relationship of the inflammatory potential of diet to affect cancer risk than would single nutrients or diet components considered individually.

Table 4. Odds ratio (OR) of bladder cancer and corresponding 95% CI according to quartiles of dietary inflammatory index (DII) among 690 cases and 665 controls, in strata of selected covariates (Italy, 2003-2014)

	Cases/ Controls	DII Quartiles, OR (95% CI)*				P_{trend}	$P_{\text{interaction}}$
		-5.94, -2.41	-2.40, -0.92	-0.91, 0.41	0.42, 4.58		
Sex							.002
Male	595/561	1 [†]	1.39 (0.95, 2.03)	1.12 (0.74, 1.70)	1.83 (1.14, 2.91)	.03	
Female	95/104	1 [†]	2.56 (0.83, 7.89)	11.27 (3.23, 39.31)	5.73 (1.46, 22.44)	.004	
Age (years)							.59
<65	255/297	1 [†]	1.25 (0.70, 2.22)	1.30 (0.71, 2.38)	1.53 (0.79, 2.96)	.21	
≥65	435/368	1 [†]	1.63 (1.04, 2.56)	1.53 (0.92, 2.52)	2.45 (1.38, 4.34)	.004	
Education (years)							.88
<7	292/273	1 [†]	1.46 (0.84, 2.53)	1.42 (0.77, 2.61)	1.56 (0.77, 3.14)	.23	
≥7	397/392	1 [†]	1.52 (0.95, 2.44)	1.41 (0.85, 2.33)	2.22 (1.27, 3.88)	.009	
Tobacco smoking							.36
Never smokers	96/237	1 [†]	1.42 (0.64, 3.14)	3.06 (1.26, 7.44)	4.04 (1.51, 10.80)	.003	
Ever smokers	585/428	1 [†]	1.58 (1.06, 2.34)	1.41 (0.92, 2.15)	1.96 (1.22, 3.15)	.01	

* Adjusted for age, sex, year of interview, study center, education (<7; 7-11; ≥12 years), tobacco smoking (never; former; current: <15, 15-24, ≥25 cigarettes/day), and total energy intake, when appropriate.

[†] Reference category.

In conclusion, this study indicates a detrimental role of a pro-inflammatory diet, as measured by higher DII scores, on BC risk through a process of inflammation. These results have to be established in further studies to confirm this association.

References

- Ferlay JSI, Ervik M, Dikshit R, et al. GLOBOCAN 2012 v1.1, cancer incidence and mortality worldwide: IARC Cancer Base No. 11 [Internet]. In: Cancer IAFRo, ed. Lyon, France; 2014. Available at: <http://globocan.iarc.fr>. Accessed May 20, 2016.
- Burger M, Catto JW, Dalbagni G, et al. Epidemiology and risk factors of urothelial bladder cancer. *Eur Urol*. 2013;63:234-241.
- Nelson D, Ganss R. Tumor growth or regression: powered by inflammation. *J Leukoc Biol*. 2006;80:685-690.
- Balkwill F, Charles KA, Mantovani A. Smoldering and polarized inflammation in the initiation and promotion of malignant disease. *Cancer Cell*. 2005;7:211-217.
- Nesi G, Nobili S, Cai T, Caini S, Santi R. Chronic inflammation in urothelial bladder cancer. *Virchows Arch*. 2015;467:623-633.
- Yao B, Yan Y, Ye X, et al. Intake of fruit and vegetables and risk of bladder cancer: a dose-response meta-analysis of observational studies. *Cancer Causes Control*. 2014;25:1645-1658.
- Wu W, Tong Y, Zhao Q, Yu G, Wei X, Lu Q. Coffee consumption and bladder cancer: a meta-analysis of observational studies. *Sci Rep*. 2015;5:9051.
- Johansson-Persson A, Ulmius M, Cloetens L, Karhu T, Herzig KH, Onning G. A high intake of dietary fiber influences C-reactive protein and fibrinogen, but not glucose and lipid metabolism, in mildly hypercholesterolemic subjects. *Eur J Nutr*. 2014;53:39-48.
- Estruch R, Martinez-Gonzalez MA, Corella D, et al. Effects of a Mediterranean-style diet on cardiovascular risk factors: a randomized trial. *Ann Intern Med*. 2006;145:1-11.
- Shivappa N, Steck SE, Hurley TG, Hussey JR, Hebert JR. Designing and developing a literature-derived, population-based dietary inflammatory index. *Public Health Nutr*. 2014;17:1689-1696.
- Shivappa N, Steck SE, Hurley TG, et al. A population-based dietary inflammatory index predicts levels of C-reactive protein in the Seasonal Variation of Blood Cholesterol Study (SEASONS). *Public Health Nutr*. 2014;17:1825-1833.
- Tabung FK, Steck SE, Zhang J, et al. Construct validation of the dietary inflammatory index among postmenopausal women. *Ann Epidemiol*. 2015;25:398-405.
- Shivappa N, Bosetti C, Zucchetto A, et al. Association between dietary inflammatory index and prostate cancer among Italian men. *Br J Nutr*. 2015;113:278-283.
- Shivappa N, Jackson MD, Bennett F, Hebert JR. Increased Dietary Inflammatory Index (DII) is associated with increased risk of prostate cancer in Jamaican men. *Nutr Cancer*. 2015;67:941-948.
- Tabung FK, Steck SE, Ma Y, et al. The association between dietary inflammatory index and risk of colorectal cancer among postmenopausal women: results from the Women's Health Initiative. *Cancer Causes Control*. 2015;26:399-408.
- Shivappa N, Zucchetto A, Serraino D, Rossi M, La Vecchia C, Hebert JR. Dietary inflammatory index and risk of esophageal squamous cell cancer in a case-control study from Italy. *Cancer Causes Control*. 2015;26:1439-1447.
- Shivappa N, Sandin S, Lof M, Hebert JR, Adami HO, Weiderpass E. Prospective study of dietary inflammatory index and risk of breast cancer in Swedish women. *Br J Cancer*. 2015;113:1099-1103.
- Dugue PA, Hodge AM, Brinkman MT, et al. Association between selected dietary scores and the risk of urothelial cell carcinoma: a prospective cohort study. *Int J Cancer*. 2016;139:1251-1260.
- Polesel J, Bosetti C, di Maso M, et al. Duration and intensity of tobacco smoking and the risk of papillary and non-papillary transitional cell carcinoma of the bladder. *Cancer Causes Control*. 2014;25:1151-1158.
- Turati F, Bosetti C, Polesel J, et al. Coffee, tea, cola, and bladder cancer risk: dose and time relationships. *Urology*. 2015;86:1179-1184.
- Salvini S, Parpinel M, Gnagnarella P, Maisonneuve P, Turrini A. *Banca Dati di Composizione degli Alimenti per Studi Epidemiologici in Italia*. Milan: Istituto Europeo di Oncologia; 1998.
- Decarli A, Franceschi S, Ferraroni M, et al. Validation of a food-frequency questionnaire to assess dietary intakes in cancer studies in Italy. Results for specific nutrients. *Ann Epidemiol*. 1996;6:110-118.
- Franceschi S, Barbone F, Negri E, et al. Reproducibility of an Italian food frequency questionnaire for cancer studies. Results for specific nutrients. *Ann Epidemiol*. 1995;5:69-75.
- Cavicchia PP, Steck SE, Hurley TG, et al. A new dietary inflammatory index predicts interval changes in serum high-sensitivity C-reactive protein. *J Nutr*. 2009;139:2365-2372.
- Ros MM, Bueno-de-Mesquita HB, Kampman E, et al. Plasma carotenoids and vitamin C concentrations and risk of urothelial cell carcinoma in the European Prospective Investigation into Cancer and Nutrition. *Am J Clin Nutr*. 2012;96:902-910.

26. Parkes GC, Whelan K, Lindsay JO. Smoking in inflammatory bowel disease: impact on disease course and insights into the aetiology of its effect. *J Crohns Colitis*. 2014;8:717-725.
27. Nanri A, Moore MA, Kono S. Impact of C-reactive protein on disease risk and its relation to dietary factors. *Asian Pac J Cancer Prev*. 2007;8:167-177.
28. Cumberbatch MG, Rota M, Catto JW, La Vecchia C. The role of tobacco smoke in bladder and kidney carcinogenesis: a comparison of exposures and meta-analysis of incidence and mortality risks. *Eur Urol*. 2016;70:458-466.
29. Ohnishi S, Ma N, Thanan R, et al. DNA damage in inflammation-related carcinogenesis and cancer stem cells. *Oxid Med Cell Longev*. 2013;2013:387014.
30. de Vivar Chevez AR, Finke J, Bukowski R. The role of inflammation in kidney cancer. *Adv Exp Med Biol*. 2014;816:197-234.