

# The nuclear matrix elements of $0\nu\beta\beta$ decay and the NUMEN project at INFN-LNS

F. CAPPUZZELLO<sup>1,2</sup>, C. AGODI<sup>1</sup>, F. BALESTRA<sup>3,4</sup>, R. BIJKER<sup>5</sup>,  
D. BONANNO<sup>6</sup>, D. BONGIOVANNI<sup>6</sup>, V. BRANCHINA<sup>2,6</sup>, S. CALABRESE<sup>1,2</sup>,  
L. CALABRETTA<sup>1</sup>, A. CALANNA<sup>1</sup>, D. CALVO<sup>3</sup>, D. CARBONE<sup>1</sup>,  
M. CAVALLARO<sup>1</sup>, M. COLONNA<sup>1</sup>, S. FERRERO<sup>3,4</sup>, A. FOTI<sup>2,6</sup>,  
P. FINOCCHIARO<sup>1</sup>, G. GIRAUDDO<sup>3,4</sup>, V. GRECO<sup>1,2</sup>, F. IAZZI<sup>3,4</sup>,  
R. INTROZZI<sup>3,4</sup>, G. LANZALONE<sup>1</sup>, A. LAVAGNO<sup>3,4</sup>, D. LO PRESTI<sup>2,6</sup>,  
F. LONGHITANO<sup>6</sup>, A. MUOIO<sup>1</sup>, L. PANDOLA<sup>1</sup>, D. RIFUGGIATO<sup>1</sup>,  
M.V. RUSLAN<sup>5</sup>, E. SANTOPINTO<sup>5</sup>, L. SCALTRITO<sup>3,4</sup>, S. TUDISCO<sup>1</sup>,  
V. ZAGATTO<sup>1</sup>

<sup>1</sup>INFN - Laboratori Nazionali del Sud, Catania, Italy

<sup>2</sup>Dipartimento di Fisica e Astronomia, Università di Catania, Italy

<sup>3</sup>INFN, Sezione di Torino, Torino, Italy

<sup>4</sup>Politecnico di Torino, Torino, Italy

<sup>5</sup>INFN, Sezione di Genova, Genova, Italy

<sup>6</sup>INFN, Sezione di Catania, I-95123 Catania, Italy

## Abstract

An innovative technique to access the nuclear matrix elements entering the expression of the life time of the double beta decay by relevant cross sections measurements of double charge exchange reactions is proposed. A key aspect of the project is the use of the MAGNEX large acceptance magnetic spectrometer, for the detection of the ejectiles, and of the LNS K800 Superconducting Cyclotron (CS), for the acceleration of the required high resolution and low emittance heavy-ion beams, already in operation at INFN Laboratori Nazionali del Sud in Catania (Italy).

# 1 Introduction

Neutrinoless double beta decay,  $0\nu\beta\beta$ , is at the present time strongly pursued both experimentally and theoretically [1]. Its observation will determine whether the neutrino is a Dirac or Majorana particle and will provide a measurement of the average neutrino mass, which is nowadays one of the most fundamental problems in physics. An innovative technique to access the Nuclear Matrix Elements (NME) entering the expression of the life time of the  $0\nu\beta\beta$  by relevant cross sections of double charge exchange (DCE) reactions is proposed. The basic point is the coincidence of the initial and final state wave-functions in the two classes of processes and the similarity of the transition operators, which in both cases present a superposition of Fermi, Gamow-Teller and rank-two tensor components with a relevant implicit momentum transfer. First pioneering experimental results obtained at the INFN-LNS laboratory for the  $^{40}\text{Ca}(^{18}\text{O}, ^{18}\text{Ne})^{40}\text{Ar}$  reaction at 270 MeV, give encouraging indication on the capability of the proposed technique to access relevant quantitative information.

First experimental results, obtained at the INFN-LNS laboratory in Catania, for the  $^{40}\text{Ca}(^{18}\text{O}, ^{18}\text{Ne})^{40}\text{Ar}$  reaction at 270 MeV, in a wide range of transferred momenta, give encouraging indication on the capability to access quantitative information towards the determination of the NME for  $0\nu\beta\beta$  decay [2]. On the basis of the above mentined ground-breaking achievement, we propose an ambitious project, NUMEN, with the aim to go deep insight in the HI-DCE studies on nuclei of interest in  $0\nu\beta\beta$  decay[3]. There are a number of important similarities among DCE and  $0\nu\beta\beta$  decay processes, despite they are mediated by different interactions, also the description of NMEs extracted from DCE and  $0\nu\beta\beta$  presents the same degree of complexity, with the advantage for DCE to be accessible in laboratory. However a simple relation between DCE cross sections and  $\beta\beta$ -decay half-lives is not trivial and needs to be explored.

## 2 The project

The availability of the MAGNEX spectrometer [4] for high resolution measurements of very suppressed reaction channels was essential for the first pilot experiment. Moreover the measurement of DCE high resolution energy spectra and accurate cross sections at very forward angles are key points to identify the transitions of interest [5]. The concurrent measurement of the other relevant reaction channels allows to isolate the direct DCE mechanism



Figure 1: A view of MAGNEX spectrometer at Laboratori Nazionali del Sud in Catania.



Figure 2: A view of K800 Superconducting Cyclotron at Laboratori Nazionali del Sud in Catania.

from the competing transfer processes. These are at least of 4th-order and can be effectively minimized by the choice of the proper projectile-target system and incident energy [6].

The present limit of low beam current we have experienced both for the CS accelerator and for the MAGNEX focal plane detector must be sensibly overcome. For a systematic study of the many hot cases of  $\beta\beta$  decays an upgraded set-up, able to work with two orders of magnitude more current than the present, is thus necessary. This goal can be achieved by a substantial change in the technologies used in the beam extraction and in the detection of the ejectiles. For the accelerator the use of a stripper induced extraction is an adequate choice. However with the present set-up it is difficult to suitably extend this research to the hot cases, where  $\beta\beta$  decay studies are and will be concentrated. For the spectrometer the main foreseen upgrades are:

1. The substitution of the present Focal Plane Detector (FPD) [7-8]

technology with tracker based on a micro-pattern electron amplifier;

2. The substitution of the wall of silicon pad stopping detectors with a wall of telescopes based on SiC-CsI detectors;

3. The enhancement of the maximum magnetic rigidity;

4. The introduction of an array of detectors for measuring the coincident  $\gamma$ -rays.

In this framework we propose four phases in the NUMEN project, looking forward to do, in the same time, both the experimental and the up-grade activity, as indicated in the following Phases of the project.

### 3 The facility upgrade

Once all the building block for the upgrade of the accelerator and spectrometer facility will be ready at the LNS a Phase3, connected to the disassembling of the old set-up and re-assembling of the new will start. An estimate of about 18-24 months is considered. Phase4: the experimental campaign The Phase4 will consist of a series of experimental campaigns at high beam intensities (some  $p\mu A$ ) and long experimental runs in order to reach in each experiment integrated charge of hundreds of mC up to C, for the experiments in coincidences, spanning all the variety of candidate isotopes for  $0\nu\beta\beta$  decay, like:  $^{48}\text{Ca}$ ,  $^{82}\text{Se}$ ,  $^{96}\text{Zr}$ ,  $^{100}\text{Mo}$ ,  $^{110}\text{Pd}$ ,  $^{124}\text{Sn}$ ,  $^{128}\text{Te}$ ,  $^{130}\text{Te}$ ,  $^{136}\text{Xe}$ ,  $^{148}\text{Nd}$ ,  $^{150}\text{Nd}$ ,  $^{154}\text{Sm}$ ,  $^{160}\text{Gd}$ ,  $^{198}\text{Pt}$ .

### 4 Perspectives

Once selected the optimal experimental condition for the different cases in the Phase2, with the upgrades, both of CS and the MAGNEX array, the Phase4 will be devoted to collect data addressed to give, with an accurate analysis, a rigorous determination of the absolute cross sections values and their uncertainties for all the system of interest, to the challenging determination of the  $0\nu\beta\beta$  decay nuclear matrix elements, that is the ambitious goal of NUMEN Project.

## References

[1] Report to the Nuclear Science Advisory Committee, Neutrinoless Double Beta Decay, 2014.

[2] F. Cappuzzello, et al., Heavy-ion double charge-exchange reactions: a tool towards  $0\nu\beta\beta$  nuclear matrix elements, Eur. Phys. J. A (2015) accepted.

[3] C. Agodi et al., Nuclear and Particle Physics Proceedings Vol 265-266, pag.28-30.

[4] F. Cappuzzello et al.,MAGNEX: an innovative large acceptance spectrometer for nuclear reaction studies in: Magnets: Types, Uses and Safety, Nova Publisher Inc., New York, 2011, pp 1-63.

[5] H. Matsubara, et al., Few-Body Systems 54 (2013) 1433.

[6] D. M. Brink, Phys. Lett. B 40 (1972) 37.

[7] M. Cavallaro, et al., Eur. Phys. J. A (2012) 48:59.

[8] D. Carbone, et al., Eur. Phys. J. A (2012) 48:60.