

Picomolar Amyloid- β Positively Modulates Synaptic Plasticity and Memory in Hippocampus

Daniela Puzzo,^{1,2} Lucia Privitera,^{1,2} Elena Leznik,¹ Mauro Fà,¹ Agnieszka Staniszewski,¹ Agostino Palmeri,² and Ottavio Arancio¹

¹Department of Pathology, Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Columbia University, New York, New York 10032, and

²Department of Physiological Sciences, University of Catania, Catania 95125, Italy

Amyloid- β (A β) peptides are produced in high amounts during Alzheimer's disease, causing synaptic and memory dysfunction. However, they are also released in lower amounts in normal brains throughout life during synaptic activity. Here we show that low picomolar concentrations of a preparation containing both A β ₄₂ monomers and oligomers cause a marked increase of hippocampal long-term potentiation, whereas high nanomolar concentrations lead to the well established reduction of potentiation. Picomolar levels of A β ₄₂ also produce a pronounced enhancement of both reference and contextual fear memory. The mechanism of action of picomolar A β ₄₂ on both synaptic plasticity and memory involves α 7-containing nicotinic acetylcholine receptors. These findings strongly support a model for A β effects in which low concentrations play a novel positive, modulatory role on neurotransmission and memory, whereas high concentrations play the well known detrimental effect culminating in dementia.

Key words: amyloid- β ; synaptic plasticity; memory; hippocampus; α -7 nicotinic receptors; Alzheimer's disease

Introduction

Amyloid- β (A β) peptides constitute the major component of amyloid plaques in Alzheimer's disease (AD). A β peptides derive from cleavage of the amyloid- β precursor protein (APP) (Kametsani, 2008). APP is initially cleaved by α - or β -secretases, generating large, soluble, secreted fragments (sAPP α and sAPP β) and membrane-associated Carboxy-terminal fragments (CTFs). A β peptides of various lengths (e.g., A β ₄₂ and A β ₄₀) are produced after β -secretase cleavage, followed by γ -secretase cleavage. Because of the relevance of the amyloid hypothesis in AD, toxic effects of high A β levels have been widely investigated during the last 20 years. Abnormally high amounts of A β have been shown to cause synaptic and memory dysfunction (Haass and Selkoe, 2007). However, A β is normally produced in the brain, where the *in vivo* concentration in the rodent brain has been estimated to be in the picomolar range (Cirrito et al., 2003). Nonetheless, most researchers have held the view that it is just a "garbage" product of APP metabolism generated during the production of other, biologically important APP fragments, with the exception of two studies suggesting that picomolar levels of A β ₄₀ play a neurotrophic role in cell cultures (Yankner et al., 1990; Plant et al., 2003),

and another work in which A β ₄₂ increased the number of newborn neurons in cultured neural stem cells (López-Toledano and Shelanski, 2004).

A positive role of A β in synaptic plasticity and memory in normal brain is supported by the observation that APP knock-out (KO) mice show long-term potentiation (LTP) and memory impairment (Dawson et al., 1999; Phinney et al., 1999; Seabrook et al., 1999). The knock-out approach, however, has precluded a clear assessment of the physiological role of A β because of the possibility that other APP fragments and APP itself might also be biologically important. For instance, studies on APP fragment function have demonstrated that the sAPP fragments may have neurotrophic properties and enhance synaptic plasticity and memory (Araki et al., 1991; Mattson, 1994; Mucke et al., 1994; Smith-Swintosky et al., 1994; Furukawa et al., 1996; Ishida et al., 1997; Meziane et al., 1998), and the intracellular CTF may regulate gene transcription, calcium signaling, synaptic plasticity, and memory (Cao and Südhof, 2001; Gao and Pimplikar, 2001; Kimberly et al., 2001; Leissring et al., 2002; Ma et al., 2007). Another important link between A β , synaptic plasticity, and memory has been suggested by studies in which the loss of presenilin function, the enzymatic subunit of the multicomponent γ -secretase protein complex, has been found to impair LTP and memory (Saura et al., 2004; Dewachter et al., 2006). Likewise, suppression of β -secretase function in BACE1 knock-out mice also impaired synaptic plasticity and memory (Laird et al., 2005). However, because of the diverse substrates and pathways activated by the secretases in addition to APP, it remains to be determined through what mechanism(s), A β or otherwise, loss of secretase function causes these effects. Because of these findings, we set out to investigate whether low amounts of A β ₄₂, in the picomolar

Received June 12, 2008; revised Oct. 23, 2008; accepted Nov. 18, 2008.

This work was supported by National Institutes of Health Grant NS049442 (to O.A.) and Alzheimer's Association Grant NIRG-07-59597 (to D.P.). We thank Cristina Alberini, Rusiko Bourtschouladze, Moses V. Chao, Gilbert Di Paolo, Ana Garcia-Osta, Paul M. Mathews, Ipe Ninan, Filippo Palermo, Marina Picciotto, Lorna Role, and David Talmage for helpful comments and discussion, Benjamin Sherman for the genotyping of α 7-nAChR KO mice, Gakuji Hashimoto and Fahad Aziz for the A β measurements, and the rest of the Arancio laboratory for numerous discussions.

Correspondence should be addressed to Dr. Ottavio Arancio, Columbia University Medical Center, Department of Pathology and Cell Biology, Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Irving Cancer Research Center 1130 St. Nicholas Avenue, Room 603, New York, NY 10032. E-mail: oa1@columbia.edu.

DOI:10.1523/JNEUROSCI.2692-08.2008

Copyright © 2008 Society for Neuroscience 0270-6474/08/2814537-09\$15.00/0

range, as in the normal brain, enhance synaptic plasticity and memory.

Materials and Methods

Animals. Mice were maintained on a 12 h light/dark cycle in temperature- and humidity-controlled rooms of the animal facility of Columbia University. Animals were killed by cervical dislocation followed by decapitation. Three- to 4-month-old male wild-type (WT) mice (C57BL/6) were obtained from a breeding colony kept in the animal facility of Columbia University. $\alpha 7$ -KO mice and their WT littermates were obtained by crossing heterozygous animals purchased from Jackson Laboratories (#003232, B6.129S7-Chrna7^{tm1Bay}/J). Mice from the 7-null mutation line were genotyped as follows: 2 mm tails from the heterozygous breedings were digested and the DNA was extracted using lysis buffer containing 1 M Tris-HCl, 0.5 M EDTA, 10% SDS, 5 M NaCl, proteinase K in dH₂O. Jackson Laboratories supplied the sequence of primers used to identify either the neo-cassette of the null mutation or the wild-type allele, for use with the PCR: forward, 5'-CCTGGTCCT-GCTGTGTTAACTGCTTC-3'; reverse WT($\alpha 7^+$), 5'-CTGCTGG-GAAATCCTAGGCACACTTGAG-3'; reverse Neo($\alpha 7^-$), 5'-GACAAGACCGGCTTCCATCC-3'. Thermocycling conditions were as follows: 95°C for 4 min; 35 cycles of 5°C for 30 s, 56°C for 60 s, 72°C for 90 s; 72°C for 10 min; store at 4°C. PCR products were run on a 2% agarose gel, using ethidium bromide UV detection of bands at 440 bp ($\alpha 7^+$) or 750 bp ($\alpha 7^-$).

A β preparation. A β_{42} was prepared as previously described (Puzzo et al., 2005). Briefly, the lyophilized peptide (American Peptide) was suspended in 100% 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP) (Sigma-Aldrich) to 1 mM. HFIP was allowed to evaporate, and the resulting clear peptide film was stored at 20°C. Twenty-four hours before use, the film was added to DMSO (Sigma-Aldrich) and sonicated for 10 min. A β_{42} -DMSO was diluted into the bath solution, vortexed for 30 s, and incubated at 4°C for 24 h. The biochemistry of this aged synthetic A β was routinely characterized using Western blot analysis in which A β samples were resolved by Tris-tricine PAGE under nondenaturing/nonreducing conditions, and then transferred onto a nitrocellulose membrane. After membrane incubation with the anti-human A β monoclonal antibody 6E10 (Covance Research Products), horseradish peroxidase chemiluminescence revealed the presence of both monomers and oligomers (supplemental Fig. 1, available at www.jneurosci.org as supplemental material). Scramble A β_{42} was purchased from AnaSpec Inc. Concentrations of A β_{42} were calculated based on the molecular weight of its monomeric peptide.

Electrophysiological measurements. Brain slices (400 μ m) from C57BL/6 mice were cut and maintained in an interface chamber at 29°C for 90 min before recording, as previously reported (Puzzo et al., 2005). The bath solution consisted of the following (in mM): 124.0 NaCl, 4.4 KCl, 1.0 Na₂HPO₄, 25.0 NaHCO₃, 2.0 CaCl₂, 2.0 MgSO₄, and 10.0 glucose. The flow rate of the perfusion was 1 ml/min. The stimulating electrode, a bipolar tungsten electrode, was placed at the level of the Schaeffer collateral fibers, whereas the recording electrode, a glass electrode filled with bath solution, was placed at the level of CA1 stratum radiatum. Basal synaptic transmission (BST) was assayed by plotting the stimulus voltages against slopes of field EPSP. For LTP experiments, a 15 min baseline was recorded every minute at an intensity that evokes a response \sim 35% of the maximum evoked response. LTP was induced using θ -burst stimulation (4 pulses at 100 Hz, with the bursts repeated at 5 Hz and each tetanus including three 10-burst trains separated by 15 s). For posttetanic potentiation (PTP) measurements, three 10-burst trains similar to those used to produce LTP were applied in the presence of 50 μ M D-APV. Mecamylamine (MCL) and α -bungarotoxin (α -BgTx) were purchased from Sigma-Aldrich.

For patch-clamp experiments, 350 μ m hippocampal slices were cut with a vibratome and maintained in a submerged chamber at 29°C, perfused with artificial cerebrospinal fluid containing (in mM): 125 NaCl, 2.5 KCl, 1.25 Na₂HPO₄, 25 NaHCO₃, 2 CaCl₂, 1.4 MgCl₂, 25 glucose, pH 7.4 (95% O₂, 5% CO₂). Slices were permitted to recover from cutting for at least 90 min before recordings. For recordings, neurons were voltage-clamped throughout the experiment. Patch pipettes (4–6 M Ω) were

pulled from thick-walled borosilicate glass tubing and filled with a solution containing (in mM): 117.5 Cs-methylsulfonate, 17.5 CsCl₂, 4 NaCl, 0.1 EGTA, 10 HEPES, 5 QX-314·Cl, 4 MgATP, 0.3 Na₂GTP, 10 phosphocreatine-Tris, pH adjusted to 7.3 with CsOH, osmolarity adjusted to 290 mOsm with sucrose. Currents were recorded with a Warner amplifier (PC-501A) and filtered at 1 kHz (holding potential, -70 mV). The input resistance was determined from the current at the end of a 5 mV, 10 ms hyperpolarization voltage step. To eliminate artifacts resulting from variation of the seal properties, the access resistance was monitored for constancy throughout all experiments. Synaptic input was evoked by Schaeffer collateral pathway stimulation of 150 μ s pulses at a frequency of 0.1 Hz using concentric bipolar electrodes. Ten minutes of stable access resistance was required before beginning measurements of the EPSC amplitude. The amplitude was measured automatically by using the Clampfit program (version 10.1) from Molecular Devices. For the *I*-*V* experiments, 100 μ M picrotoxin was added to the bath. To isolate the synaptic currents, membrane currents recorded at each membrane potential in the absence of a Schaeffer collateral stimulation were subtracted from the evoked synaptic responses. The AMPAR/NMDAR receptor ratio was calculated by dividing the amplitude of the AMPAR current measured at the peak response at -70 mV by the NMDAR current measured 30 ms after the peak at $+50$ mV. Miniature EPSCs were recorded at -60 mV in the presence of 1 μ M tetrodotoxin. PTP was induced using the same stimulation protocol as in the extracellular recordings. All of the electrophysiological experiments were performed using polypropylene tubing, which has been shown not to alter the ratio of different A β fragments (Lewczuk et al., 2006).

A β measurements. Brain regions were dissected, flash-frozen, homogenized, and extracted, as previously described (Citron et al., 1997), before sandwich ELISA using the mouse β -amyloid 1-42 kit from Invitrogen to quantitate A β_{42} .

Infusion technique. After anesthesia with 20 mg/kg Avertin, mice were implanted with a 26-gauge guide cannula into the dorsal part of the hippocampi (coordinates: posterior = 2.46 mm, lateral = 1.50 mm to a depth of 1.30 mm) (Paxinos, 1998). The cannulas were fixed to the skull with acrylic dental cement (Paladur). After 6–8 d, we bilaterally injected 200 pM A β_{42} , or 200 pM scramble A β_{42} , or vehicle in a final volume of 1 μ l over 1 min (0.903 pg) through infusion cannulas that were connected to a microsyringe by a polyethylene tube. For the Morris water maze, mice were injected 20 min before performing each session and the probe trial, whereas for fear conditioning, mice received a single injection 20 min before the training. Mice were handled once a day for 3 d before behavioral experiments. During infusion, animals were handled gently to minimize stress. After infusion, the needle was left in place for another minute to allow diffusion. After behavioral testing, a solution of 4% methylene blue was infused into the cannulas. Animals were killed and their brains were removed, frozen, and then cut at -20° with a cryostat for histological localization of infusion cannulas.

Behavioral studies. For the Morris water maze, mice were trained in two daily sessions (4 h apart), each consisting of three trials (1 min each), for 3 d. Time required to reach the hidden platform (latency) was recorded. During this time, the location of the platform was kept constant. The training was followed by four probe trials with the platform moved to test the retention of the spatial memory. The maze was divided into four quadrants. The percentage of time spent in the quadrant that previously contained the platform was recorded and analyzed with a video tracking system (HVS 2020; HVS Image). After the probe trials, visual, motor, and motivation skills were also tested using a visible platform to measure the time and the speed needed to reach a visible platform placed within the pool by means of the video tracking system. No difference in the time and the speed needed to reach the platform was observed among the different groups of mice.

For fear conditioning, mice were placed in a conditioning chamber for 2 min before the onset of a tone [conditioned stimulus (CS)] (a 30 s, 85 dB sound at 2800 Hz). In the last 2 s of the CS, mice were given a 2 s, 0.45 mA foot shock (unconditioned stimulus) through the bars of the floor. Then, the mice were left in the conditioning chamber for another 30 s. Freezing behavior, defined as the absence of movement except for that needed for breathing, was scored using Freezeview software. Contextual

fear learning was evaluated 24 h after training by measuring freezing for 5 min in the chamber in which the mice were trained. Cued fear learning was evaluated 24 h after contextual testing by placing mice in a novel context for 2 min (pre-CS test), after which they were exposed to the CS for 3 min (CS test), and freezing was measured. Sensory perception of the shock was determined through threshold assessment. Briefly, the electric current (0.1 mA for 1 s) was increased at 30 s intervals by 0.1–0.7 mA. Threshold to flinching (first visible response to shock), jumping (first extreme motor response), and screaming (first vocalized distress) was quantified for each animal by averaging of the shock intensity at which each animal manifested a behavioral response of that type to the foot shock. No difference in the sensory threshold assessment was observed among different groups of mice in experiments in which fear conditioning was tested. Moreover, different groups of mice had similar exploratory behavior, as demonstrated by a similar percentage of time spent in the center compartment and the number of entries into the center compartment in the open-field test.

Statistics. For all experiments, mice were coded by “blind” investigators with respect to treatment and genotype. Data are expressed as mean \pm SEM. Statistical analysis was performed with two-way ANOVA (with multiple comparisons or with repeated measures for LTP), Student’s *t* test (pairwise comparisons). The level of significance was set for $p < 0.05$.

Results

Picomolar concentrations of $A\beta_{42}$ enhance hippocampal long-term potentiation

Given that extracellular $A\beta$ levels are likely to be regulated by synaptic activity (Kamenetz et al., 2003; Cirrito et al., 2005), we speculated that $A\beta$ is normally involved in hippocampal LTP, the dominant model of activity-dependent synaptic plasticity. However, lack of information regarding range of $A\beta$ assemblies present in human brain places a challenge in examining the role of $A\beta$ in normal brain. As a first step toward the understanding of the putative physiological role of $A\beta$, we reasoned that a straightforward strategy was to perform a dose–response curve for the effect of $A\beta$ on LTP. As previously demonstrated (Haass and Selkoe, 2007), when hippocampal slices were perfused with an aged $A\beta_{42}$ preparation at 200 nM for 20 min, we observed an impairment of LTP at the synapses between Schaeffer collateral fibers and CA1 neurons ($n = 7$ slices from 7 mice, two-way ANOVA $F_{(1,17)} = 27.56$, $p < 0.0001$ compared with 12 vehicle-treated slices from 12 mice) (Fig. 1A). Surprisingly, when we perfused slices with lower concentrations of $A\beta_{42}$, the first portion of the concentration–response curve showed an enhancement of LTP with a maximal effect around 200 pM ($n = 6–10$ slices from 6–10 mice per $A\beta_{42}$ concentration) (Fig. 1A). Application of 200 pM human $A\beta_{42}$ for 20 min before tetanization increased CA1-LTP ($n = 10$ slices from 10 mice, $F_{(1,20)} = 7.20$, $p = 0.014$ compared with 12 vehicle-treated slices from 12 mice) (Fig. 1B). $A\beta_{42}$ alone did not affect basal transmission ($n = 6$ slices from 6 mice, $F_{(1,12)} = 0.05$, $p = 0.814$ compared with 8 vehicle-treated slices from 8 mice) (Fig. 1B), nor did application of $A\beta_{42}$ for 20 min immediately after the tetanus ($n = 5$ slices from 5 mice, $F_{(1,8)} = 0.34$, $p = 0.573$ compared with 5 vehicle-treated slices from 5 mice) (supplemental Fig. 2, available at www.jneurosci.org as supplemental material). As a control for nonspecific effects of $A\beta_{42}$, a peptide consisting of scramble $A\beta_{42}$ sequence failed to enhance LTP ($n = 6$ slices from 6 mice, $F_{(1,14)} = 0.44$, $p = 0.514$) or baseline transmission ($n = 4$ slices from 4 mice, $F_{(1,10)} = 0.02$, $p = 0.918$) (Fig. 1B). Interestingly, previous studies have reported endogenous brain concentrations of $A\beta_{42}$ in the picomolar range (Rozmahel et al., 2002; Phinney et al., 2003; Pawlik et al., 2004; Yao et al., 2004; Mastrangelo et al., 2005; Schmidt et al., 2005). When we measured the levels of $A\beta_{42}$ in




Figure 1. $A\beta_{42}$ has two opposite effects on LTP in hippocampus. **A**, Concentration–response curve for the effect of $A\beta_{42}$ on CA1-LTP indicating that the peptide has an enhancing effect with a peak around 200 pM, whereas it has an opposite detrimental effect above 20 nM. The dotted line and the shaded area around it correspond to the amount of potentiation and the SE range in vehicle-treated slices. The residual potentiation was calculated by averaging the last 5 min of LTP. **B**, Perfusion of hippocampal slices with a preparation containing human $A\beta_{42}$ (200 pM), but not scramble $A\beta_{42}$ (200 pM), for 20 min before a theta-burst stimulation increases LTP without affecting baseline transmission. The horizontal bar indicates the period during which $A\beta$ was added to the bath solution. Each bar denotes the mean \pm SEM in this and the following figures.

normal mouse brain using sandwich ELISA, cerebellum, cortex, and hippocampus also showed values in the picomolar range ($n = 4$ mice per group) (supplemental Fig. 3, available at www.jneurosci.org as supplemental material). Thus, the concentration of 200 pM suggested by our dose–response curve is likely to approximate the endogenous levels of $A\beta_{42}$.

Picomolar concentrations of $A\beta_{42}$ enhance hippocampal-dependent memory

Given that LTP is thought to be associated with learning and memory, we next assessed the effects of low doses of $A\beta_{42}$ on memory. Cannulas were implanted bilaterally into the mouse

dorsal hippocampi (Fig. 2A). After 6–8 d, animals were infused with 200 pM $A\beta_{42}$, 200 pM scramble $A\beta_{42}$, or vehicle, and after 20 min tested for reference and contextual fear memory.

Reference memory can be studied with the Morris water maze, a widely used spatial learning test known to require hippocampal function (Schenk and Morris, 1985). Mice were trained to find a hidden platform beneath the surface of the water. Mice that had received $A\beta_{42}$ needed less time to find the platform after six sessions (Fig. 2B) compared with vehicle-infused mice. Statistical analysis revealed a significant difference in the overall performance of $A\beta$ -treated mice compared with that of vehicle-treated animals ($n = 15/14$ mice; $F_{(1,27)} = 6.66$, $p = 0.016$) (Fig. 2B). Planned comparisons of latency on each individual session revealed a significant difference between $A\beta$ -treated mice and vehicle-treated mice in the sixth session ($t_{(27)} = 2.3$, $p = 0.029$). Thus, we also assessed reference memory with the probe trial, another test of spatial long-term memory (Schenk and Morris, 1985). This task is performed after the sixth hidden-platform session. The platform is removed from the water and the animals are allowed to search for 60 s. The amount of time spent in each quadrant of the maze can be used to evaluate the spatial bias of an animal's search pattern. We found that $A\beta$ -treated mice spent more time than vehicle-treated mice in the target quadrant (TQ), where the platform had been located during training ($t_{(27)} = 2.22$, $p = 0.035$) (Fig. 2C). The time spent in the TQ by the vehicle-treated mice was also verified by the single proportion test to ensure that WT mice had learning ability. These mice had a significant probability to spend >25% of the given time in the TQ (64%, $p = 0.002$). Moreover, they spent significantly more time in the TQ compared with other quadrants ($p = 0.022$). In interleaved experiments, infusion with scramble $A\beta_{42}$ did not affect mouse performance during both the six sessions with the hidden platform and the probe trials ($n = 12$ mice; hidden platform: $t_{(24)} = 0.42$, $p = 0.676$; probe trials: $t_{(24)} = 0.33$, $p = 0.743$; probability to spend >25% of the given time in the TQ: 83%, $p = 0.0001$; comparison TQ vs other quadrants: $p = 0.026$) (Fig. 2B, C). A visible platform trial performed after the probe trials did not reveal any significant difference in the time to reach the platform among the three groups during the four sessions of the task (time at the fourth session: 11.21 ± 1.79 s for $A\beta_{42}$ -infused mice; 12.45 ± 1.75 s for vehicle-treated mice; 12.74 ± 2.14 s for scramble-treated mice; $F_{(2,37)} = 0.19$, $p = 0.825$) (data not shown).

Next, we studied contextual fear memory (Phillips and LeDoux, 1992), a form of associative learning for which hippocampal function is indispensable. Mice were trained to associate neutral stimuli with an aversive one. We found no difference in the




Figure 2. Picomolar concentrations of $A\beta_{42}$ enhance hippocampal-dependent memory. **A**, Schematic representation showing cannulas implanted bilaterally into the dorsal hippocampi. **B**, Bilateral injections of human $A\beta_{42}$ (200 pM), but not scramble $A\beta_{42}$ (200 pM), into dorsal hippocampi, 20 min before the session improve the performance with the Morris water maze task both as the mice search for the hidden platform (**C**) and for the probe test. **D**, Bilateral injections of human $A\beta_{42}$ (200 pM), but not scramble $A\beta_{42}$ (200 pM), into dorsal hippocampi, 20 min before training, enhance contextual conditioning performance as the mice are exposed to the context after 24 h. The asterisks indicate statistical significance in this and the following figures.

freezing behavior among the three groups of mice during the training phase of the fear conditioning ($F_{(2,48)} = 0.04$, $p = 0.961$) (Fig. 2D). However, when fear learning was assessed 24 h later by measuring freezing behavior, the absence of all movement except for that necessitated by breathing, in response to representation of the context, $A\beta_{42}$ -treated mice showed an enhancement of freezing ($n = 19$ mice, $t_{(34)} = 2.36$, $p = 0.024$ compared with 17 vehicle-treated mice) (Fig. 2D). In contrast, scramble $A\beta_{42}$ did not affect freezing ($n = 15$ mice; $t_{(30)} = 0.33$, $p = 0.743$ compared with vehicle-injected mice) (Fig. 2D). Finally, when cued fear learning, a type of learning that depends on amygdala function (Phillips and LeDoux, 1992), was assessed on the same animals 24 h after contextual learning by measuring freezing in response to representation of the auditory cue within a completely different context, we found no difference between $A\beta_{42}$ -infused mice and vehicle-infused mice ($t_{(34)} = 0.20$, $p = 0.838$) (supplemental Fig. 4, available at www.jneurosci.org as supplemental material), indicating that behavioral changes produced by $A\beta_{42}$ were the




Figure 3. Picomolar concentrations of $A\beta_{42}$ enhance PTP without affecting NMDA and AMPA receptor currents. **A**, Example of EPSCs evoked at holding potentials ranging from -70 mV to $+50$ mV with a step of 20 mV. Dotted lines represent time points at which AMPA receptor (AMPA) and NMDA receptor (NMDAR) currents were measured. **B**, **C**, Perfusion of hippocampal slices with human $A\beta_{42}$ (200 pM) does not affect current–voltage relationship for NMDAR current (I-NMDAR) (**B**) and AMPAR current (I-AMPA) (**C**). **D**, Perfusion of hippocampal slices with human $A\beta_{42}$ (200 pM) for 20 min does not affect EPSC amplitude. Each point of the trace is the average of six consecutive recordings. The horizontal bar indicates the period of perfusion with $A\beta$. **E**, Perfusion of hippocampal slices with human $A\beta_{42}$ (200 pM) does not affect EPSC amplitude distributions. Data from one cell before and after perfusion with $A\beta_{42}$ are shown. **F**, **G**, Perfusion of hippocampal slices with human $A\beta_{42}$ (200 pM) for 20 min does not affect mEPSC frequency (**F**) and amplitude (**G**). Each point represents the average of a 2 min period. Frequency and amplitude were normalized to the average value during the 6 min before $A\beta$ application. The horizontal bar indicates the period of perfusion with $A\beta$. **H**, Perfusion of hippocampal slices with human $A\beta_{42}$ (200 pM) in D -APV (50 μ M) enhances PTP. The horizontal bars indicate the period during which $A\beta$ and/or APV were added to the bath solution.

result of a selective hippocampus-dependent effect on fear learning. Together, these experiments indicate that low doses of $A\beta_{42}$ cause a long-lasting increase in synaptic strength and enhance memory.

Picomolar concentrations of $A\beta_{42}$ enhance posttetanic potentiation

Our next goal was to identify the mechanism of the $A\beta$ -induced enhancement in LTP and memory. Given that both NMDA and AMPA receptors are known to play a key role in CA1-LTP (Lisman and Raghavachari, 2006), we assessed the current–voltage (I – V) relationships for NMDA and AMPA receptor currents before and after perfusion of the slices with 200 pM $A\beta_{42}$ for 20 min. $A\beta$ did not affect either I – V relationships for the two currents or AMPA/NMDA receptor current ratios (1.06 ± 0.14 in $A\beta$ -treated slices vs 1.02 ± 0.09 in vehicle-treated slices, $n = 3$ slices from 3 mice per group) (Fig. 3A–C). Moreover, $A\beta$ did not alter amplitude ($n = 6$ slices from 6 mice, $F_{(1,7)} = 0.41$, $p = 0.542$ compared with 3 vehicle-treated slices from 3 mice) (Fig. 3D) or amplitude distribution (Fig. 3E) of AMPA receptor-mediated EPSCs. These findings suggest that the enhanced LTP was not caused by postsynaptic changes in NMDA and AMPA receptor currents.

To further investigate the mechanism of the $A\beta$ -induced enhancement in LTP and memory, we recorded spontaneous miniature EPSCs (mEPSCs) in the presence of tetrodotoxin (1 μ M) to block sodium channels. Neither the mean frequency nor the mean amplitude of mEPSCs was affected by the treatment with 200 pM human $A\beta_{42}$ for 20 min ($n = 3$ slices from 3 mice both for the $A\beta_{42}$ -treated group and the vehicle-treated group; frequency: $F_{(1,4)} = 0.40$, $p = 0.557$; amplitude: $F_{(1,4)} = 0.44$, $p = 0.541$) (Fig. 3F, G), suggesting that $A\beta$ does not affect spontaneous release of neurotransmitter.

An alternative mechanism for the long-lasting enhancement in synaptic strength might be an increase of transmitter release during the tetanus. To test this possibility, we examined PTP in the presence of the NMDA antagonist D -APV (50 μ M) to block LTP inductive mechanisms. PTP corresponds to a transient increase of glutamate release from the presynaptic terminal that results from brief periods of high-frequency stimulation with Ca^{2+} buildup within the terminal that triggers mechanisms of short-term synaptic plasticity (Zucker and Regehr, 2002). Perfusion of the slices with 200 pM human $A\beta_{42}$ for 20 min increased PTP ($n = 11$ slices from 11 mice; $t_{(20)} = 2.18$, $p = 0.041$) (Fig. 3H).

We obtained similar results when we used a patch-clamp technique to examine PTP with and without $A\beta$ and found no additional synaptic currents ($n = 3$ slices from 3 mice) (data not shown), suggesting that $A\beta$ enhances transmitter release during the tetanus.

The enhancement of synaptic plasticity and memory induced by picomolar concentrations of $A\beta_{42}$ involves $\alpha 7$ -nicotinic ACh receptors.

$A\beta$ is known to act through multiple targets (Small et al., 2001). Given that $A\beta_{42}$ has been shown to activate $\alpha 7$ -nicotinic ACh receptors (nAChRs) at presynaptic nerve endings of synaptosomes when administered in the low picomolar range (Dougherty et al., 2003), we tested whether $\alpha 7$ -nAChRs are involved in the increase in PTP by picomolar levels of $A\beta_{42}$. When hippocampal slices were perfused with the nonselective nAChR blocker MCL ($3 \mu\text{M}$ for 20 min), the $A\beta$ -induced PTP increase was blocked in slices that had previously shown an enhancement of PTP when perfused with $A\beta$ alone ($n = 11$ slices from 11 mice; $t_{(20)} = 2.24$, $p = 0.036$ comparing $A\beta$ to $A\beta + \text{MCL}$) (Fig. 4A). In interleaved experiments, MCL alone did not affect PTP ($n = 9$ slices from 9 mice; $t_{(16)} = 0.55$, $p = 0.589$) (supplemental Fig. 5, available at www.jneurosci.org as supplemental material). Moreover, the selective $\alpha 7$ -nAChR blocker, $\alpha\text{-BgTx}$ ($0.1 \mu\text{M}$ for 20 min) also blocked the $A\beta$ -induced PTP increase ($n = 10$ slices from 10 mice; $t_{(18)} = 5.59$, $p < 0.0001$ comparing $A\beta$ to $A\beta + \alpha\text{-BgTx}$) (Fig. 4B). The slices were still capable of displaying the $A\beta$ -induced PTP enhancement if perfused again with $A\beta$ alone after washout of both $\alpha\text{-BgTx}$ and $A\beta$ ($t_{(18)} = 3.99$, $p = 0.001$) (Fig. 4B). Furthermore, both the PTP increase by $A\beta$ alone and its block by $\alpha\text{-BgTx}$ were still present during perfusion with the GABA receptor antagonist picrotoxin ($50 \mu\text{M}$) ($A\beta$, $407.56 \pm 26.99\%$; $A\beta + \alpha\text{-BgTx}$, $233.57 \pm 15.72\%$; $n = 7$ slices from 7 mice, $t_{(12)} = 5.29$, $p < 0.0001$) (data not shown), suggesting that $\alpha 7$ -nAChRs located in inhibitory interneurons (Freedman et al., 1993) are not involved in the $A\beta$ -induced enhancement of transmitter release occurring during tetanic stimulation. Finally, in interleaved experiments, $\alpha\text{-BgTx}$ alone did not affect PTP ($n = 7$ slices from 7 mice; $t_{(12)} = 2.26$, $p = 0.410$) (supplemental Fig. 5b, available at www.jneurosci.org as supplemental material).

To provide genetic evidence for the involvement of $\alpha 7$ -nAChRs in $A\beta_{42}$ -induced synaptic plasticity increase, we performed additional experiments using $\alpha 7$ -nAChR KO mice. When hippocampal slices from these animals were perfused with 200 pM $A\beta_{42}$ for 20 min before inducing LTP, the peptide failed to enhance LTP ($n = 8$ slices from 8 mice, $F_{(1,15)} = 0.001$, $p = 0.975$ compared with 9 vehicle-treated slices from 9 KO mice) (Fig. 4C). In contrast, $A\beta_{42}$ was still capable of enhancing LTP in slices from wild-type littermates ($n = 7$ slices from 7 mice, $F_{(1,12)} = 11.42$, $p =$




Figure 4. The enhancement of synaptic plasticity and memory by picomolar concentrations of $A\beta_{42}$ involves $\alpha 7$ -nAChRs. **A**, Hippocampal slices perfused with MCL ($3 \mu\text{M}$) concurrent with human $A\beta_{42}$ (200 pM) and D-APV ($50 \mu\text{M}$) no longer display the $A\beta$ -induced PTP enhancement. The horizontal bars indicate the period of perfusion with MCL, $A\beta$, and APV. **B**, Hippocampal slices perfused with $\alpha\text{-BgTx}$ ($0.1 \mu\text{M}$) concurrent with human $A\beta_{42}$ (200 pM) and D-APV ($50 \mu\text{M}$) no longer display the $A\beta$ -induced PTP enhancement. The enhancement is still present after washout of $\alpha\text{-BgTx}$ if slices are perfused again with $A\beta_{42}$ (200 pM). The horizontal bars indicate the period of perfusion with $\alpha\text{-BgTx}$, $A\beta$, and APV. **C**, Perfusion of hippocampal slices with human $A\beta_{42}$ (200 pM) for 20 min before tetanus does not increase LTP in slices from $\alpha 7$ -KO mice, whereas it still enhances potentiation in slices from WT littermates. BST was normal in the $\alpha 7$ -KO mice (data not shown). The horizontal bar indicates the period of perfusion with $A\beta$. **D**, **E**, Bilateral injections of human $A\beta_{42}$ (200 pM) into dorsal hippocampi, 15 min before training, do not enhance reference or contextual memory in $\alpha 7$ -KO mice, whereas they still enhance memory in WT littermates.

0.005 compared with 7 vehicle-treated slices from 7 wild-type mice) (Fig. 4C). $A\beta_{42}$ alone without tetanus did not affect baseline transmission either in $\alpha 7$ -KO or WT slices ($106.85 \pm 4.05\%$ and $98.53 \pm 4.49\%$, $n = 4$ slices from 4 mice for both groups;

$F_{(1,6)} = 0.10$, $p = 0.757$ and $F_{(1,6)} = 0.70$, $p = 0.433$) (data not shown).

To determine whether the effect of $A\beta_{42}$ on memory shares the same molecular mechanism as that on synaptic plasticity, we also sought to demonstrate that the enhancement of memory by picomolar levels of $A\beta_{42}$ involves $\alpha 7$ -nAChRs. Hippocampal infusion of 200 pM $A\beta_{42}$ failed to enhance reference memory in $\alpha 7$ -KO mice ($n = 10$ mice vs 9 vehicle-treated KO mice; $t_{(17)} = 0.15$, $p = 0.881$ at the six hidden sessions) (Fig. 4D), whereas it still increased memory in WT littermates ($n = 10$ mice vs 9 vehicle-treated WT mice, $t_{(17)} = 2.40$, $p = 0.028$) (Fig. 4D). Probe trials also did not show an increase in memory in $\alpha 7$ -KO mice, whereas WT-littermates still showed an increase in the time spent in the TQ (data not shown). Similarly, contextual fear memory in $\alpha 7$ -KO mice was not enhanced ($n = 7$ mice; $t_{(12)} = 0.54$, $p = 0.599$ compared with 7 vehicle-treated KO mice) (Fig. 4E), whereas WT-littermates still showed the enhancement ($n = 9$ mice; $t_{(16)} = 2.42$, $p = 0.028$ compared with 9 vehicle-treated WT mice) (Fig. 4E). Cued fear learning showed no difference between the four groups of mice (WT-vehicle, $52.45 \pm 5.04\%$; WT- $A\beta$, $51.83 \pm 5.44\%$; $\alpha 7$ -KO-vehicle, $49.54 \pm 5.86\%$; $\alpha 7$ -KO- $A\beta$, $50.24 \pm 8.18\%$; $t_{(12)} = 0.07$, $p = 0.945$ comparing $\alpha 7$ -KO-vehicle to $\alpha 7$ -KO- $A\beta$) (data not shown). Together, these findings indicate that both synaptic plasticity and memory rely on $\alpha 7$ -nAChR involvement for the enhancing effect of $A\beta$.

Discussion

Growing evidence suggests that APP and its derivatives (i.e., the products of the α -, β -, γ -, and ϵ -secretase actions) display regulatory effects on synaptic transmission before the onset of neurodegeneration (Arancio and Chao, 2007). Our data provide robust evidence that an aged preparation of human $A\beta_{42}$, when used at low concentrations, presumably neighboring those found in the normal brain (Rozmahel et al., 2002; Phinney et al., 2003; Pawlik et al., 2004; Yao et al., 2004; Mastrangelo et al., 2005; Schmidt et al., 2005), enhances LTP and memory. These effects are opposite to those (well established) caused by high levels of $A\beta_{42}$, namely an impairment of LTP and memory.

We found that picomolar concentrations of an aged $A\beta_{42}$ preparation containing both monomers and oligomers enhance synaptic plasticity and memory. However, it is impossible to know the molarities of different oligomers and monomers to which slices and hippocampi were actually exposed because $A\beta$ can easily change its conformation by the time it reaches the brain tissue after its initial preparation. Moreover, scramble $A\beta$ does not share the same capability of forming oligomers as normal $A\beta$. Furthermore, injected $A\beta$ used in our behavioral experiments would add to the baseline levels of endogenous murine $A\beta$, which might increase concentrations beyond those identified as LTP-enhancing in the acute slice studies. Finally, there is uncertainty on the molarities of monomers and different oligomeric forms of $A\beta$ existing at synapses. Thus, to address the issue of the concentration of $A\beta$ responsible for its effects, we expressed the molarity of $A\beta$ based on the molecular weight of its monomeric peptide. In addition, careful monitoring of our preparation among different experiments using PAGE analysis did not show a large variability from batch to batch in the relative percentage of different $A\beta$ species. We kept constant the temperature of the bath solution in our electrophysiological experiments because temperature is known to affect $A\beta$ conformation. We also used polypropylene tubing, which has been shown not to alter the ratio of different $A\beta$ fragments (Lewczuk et al., 2006). Finally, as previously demonstrated (for instance, Puzzo et al., 2005), our preparation was

capable of impairing LTP at a concentration of 200 nM. Nonetheless, we still cannot rigorously define the concentration and form(s) of $A\beta_{42}$ responsible for the enhancing effects on synaptic plasticity and memory and whether scramble $A\beta$ is present in the same species and concentration as normal $A\beta$. Nevertheless, the main observation of this manuscript, showing that $A\beta_{42}$ enhances synaptic plasticity and memory, is clear and has important implications for mechanisms regulating the strength of synaptic connections and memory in hippocampus. It should also be noted that our results are consistent with loss of function studies in mice lacking APP, presenilin, or BACE1 (Dawson et al., 1999; Phinney et al., 1999; Seabrook et al., 1999; Saura et al., 2004; Laird et al., 2005; Dewachter et al., 2006). The enhancement of synaptic plasticity and memory with our gain of function approach specific to $A\beta$ is likely, therefore, to be relevant to normal physiology.

Previous studies on a putative physiological function of $A\beta$ have suggested that it selectively depresses excitatory synaptic transmission (Kamenetz et al., 2003). However, they relied on preparations exposed to high amounts of $A\beta$ derived from overexpression of APP. Given that high levels of $A\beta$ are known to impair synaptic function and memory (Haass and Selkoe, 2007), the discrepancy with our findings may be reconciled by the high amount of $A\beta$ used in these previous studies, which might have produced a toxic effect on synaptic function. In contrast, we have carefully monitored the total concentration of $A\beta$ by exogenously applying the peptide, and demonstrating that we could produce both depression and enhancement of synaptic plasticity according to the concentration used in the experiment.

Analysis of baseline transmission and AMPA receptor currents revealed no changes in these parameters during perfusion with $A\beta$. Thus, we also investigated NMDA receptor currents, because they are known to play a key role in synaptic plasticity. However, similar to AMPA receptor currents, exposure to $A\beta$ did not affect NMDA receptor currents, and therefore, we had to exclude this possibility. As a next step, we investigated the miniature release of neurotransmitter before and after perfusion with $A\beta_{42}$ and found no change in mEPSC frequency and amplitude, suggesting that $A\beta$ alone does not affect these parameters. Finally, we investigated PTP, a phenomenon dependent on Ca^{2+} buildup within the presynaptic terminal, and found that picomolar concentrations of $A\beta_{42}$ enhance neurotransmitter release during high-frequency stimulation. Thus, we believe that the most likely explanation for the enhancing effect of $A\beta$ is that the increase in Ca^{2+} levels produced by $A\beta$ paired with a single stimulus is not sufficient to overcome a threshold above which neurotransmitter release is enhanced. $A\beta$, in turn, would need to be associated with high-frequency stimulation for Ca^{2+} buildup within the presynaptic terminal above a given threshold and activation of mechanisms of synaptic plasticity. Because activation of $\alpha 7$ -nAChRs is necessary for the $A\beta$ -induced increase of synaptic plasticity and memory, we propose a model whereby $A\beta$ released by neuronal activity during vesicle exocytosis (Cirrito et al., 2005) under normal conditions may modify glutamate release with a mechanism dependent upon activation of $\alpha 7$ -nAChRs. This results in an increase of synaptic plasticity and memory (supplemental Fig. 6, available at www.jneurosci.org as supplemental material). This model is consistent with the observation that $A\beta_{42}$ might bind selectively and with picomolar affinity to $\alpha 7$ -nAChRs (Wang et al., 2000), or it might be responsible for regulation of nAChR function through strong binding with membrane lipids (Small et al., 2007). Moreover, it is also consistent with the observation that $A\beta_{42}$ activates $\alpha 7$ -nAChRs at presynaptic nerve endings of synaptosomes when administered in

the low picomolar range (Dougherty et al., 2003) [whereas it would act on non- $\alpha 7$ -nAChRs at nanomolar concentrations (Dougherty et al., 2003)]. In addition, this model is consistent with the finding that activation of $\alpha 7$ -nAChRs is involved in diverse brain functions including synaptic plasticity and memory (Levin and Simon, 1998; Jones et al., 1999) and enhances transmitter release in several brain regions including the hippocampus (Gray et al., 1996; Radcliffe and Dani, 1998), the spinal cord dorsal horn (Genzen and McGehee, 2003), the olfactory bulb, and the amygdala (Girod et al., 2000). Finally, this model is consistent with the observation that nicotinic activity at pyramidal neurons boosts LTP induction (Ji et al., 2001). Clearly, although we find that $\alpha 7$ -nAChRs are involved in the $A\beta$ -induced enhancement of synaptic plasticity and memory, it remains to be determined whether these effects are mediated by a direct physical interaction of the peptide with the $\alpha 7$ -nAChR. Indeed, the effects of $A\beta_{42}$ might be more complex than those accounted for by bona fide $\alpha 7$ -nAChR agonists, such as acetylcholine and nicotine. Notwithstanding these findings, the involvement of $\alpha 7$ -nAChRs we report here is likely to provide an important contribution to the enhancing effect of $A\beta_{42}$ at picomolar concentrations.

Major drug discovery efforts are ongoing to develop strategies to decrease $A\beta$ load (Haass and Selkoe, 2007). Our work does not challenge the amyloid hypothesis. However, our lack of understanding of the physiological role of $A\beta$ may present important issues when designing effective and safe approaches to AD therapy. Our findings strongly support the possibility that $A\beta_{42}$ itself may be an important modulator of synaptic plasticity and memory in the normal brain. Indeed, and paradoxically, the use of drugs that mimic $A\beta$ structure or are targeted to the receptor(s) on which $A\beta$ acts under normal physiological conditions, or even of $A\beta$ itself or $A\beta$ derivatives, might serve to enhance memory at appropriate concentrations and conditions.


References

- Araki W, Kitaguchi N, Tokushima Y, Ishii K, Aratake H, Shimohama S, Nakamura S, Kimura J (1991) Trophic effect of beta-amyloid precursor protein on cerebral cortical neurons in culture. *Biochem Biophys Res Commun* 181:265–271.
- Arancio O, Chao MV (2007) Neurotrophins, synaptic plasticity and dementia. *Curr Opin Neurobiol* 17:325–330.
- Cao X, Südhof TC (2001) A transcriptionally [correction of transcriptively] active complex of APP with Fe65 and histone acetyltransferase Tip60. *Science* 293:115–120.
- Cirrito JR, May PC, O'Dell MA, Taylor JW, Parsadanian M, Cramer JW, Audia JE, Nissen JS, Bales KR, Paul SM, DeMattos RB, Holtzman DM (2003) *In vivo* assessment of brain interstitial fluid with microdialysis reveals plaque-associated changes in amyloid- β metabolism and half-life. *J Neurosci* 23:8844–8853.
- Cirrito JR, Yamada KA, Finn MB, Sloviter RS, Bales KR, May PC, Schoepp DD, Paul SM, Mennerick S, Holtzman DM (2005) Synaptic activity regulates interstitial fluid amyloid-beta levels in vivo. *Neuron* 48:913–922.
- Citron M, Westaway D, Xia W, Carlson G, Diehl T, Levesque G, Johnson-Wood K, Lee M, Seubert P, Davis A, Kholodenko D, Motter R, Sherrington R, Perry B, Yao H, Strome R, Lieberburg I, Rommens J, Kim S, Schenk D, et al. (1997) Mutant presenilins of Alzheimer's disease increase production of 42-residue amyloid-beta-protein in both transfected cells and transgenic mice. *Nat Med* 3:67–72.
- Dawson GR, Seabrook GR, Zheng H, Smith DW, Graham S, O'Dowd G, Bowery BJ, Boyce S, Trumbauer ME, Chen HY, Van der Ploeg LH, Srinathsinghi DJ (1999) Age-related cognitive deficits, impaired long-term potentiation and reduction in synaptic marker density in mice lacking the beta-amyloid precursor protein. *Neuroscience* 90:1–13.
- Dewachter I, Ris L, Croes S, Borghgraef P, Voets T, Nilius B, Godaux E, Van Leuven F (2006) Differential effects of presenilin-1 deficiency and mutant presenilin-1 on synaptic activity. In: *Alzheimer's Association 10th International Conference on Alzheimer's Disease and Related Disorders*, p 558, O53-03-08. Madrid. Chicago: Alzheimer's Association.
- Dougherty JJ, Wu J, Nichols RA (2003) β -Amyloid regulation of presynaptic nicotinic receptors in rat hippocampus and neocortex. *J Neurosci* 23:6740–6747.
- Freedman R, Wetmore C, Strömberg I, Leonard S, Olson L (1993) α -Bungarotoxin binding to hippocampal interneurons: immunocytochemical characterization and effects on growth factor expression. *J Neurosci* 13:1965–1975.
- Furukawa K, Sopher BL, Rydel RE, Begley JG, Pham DG, Martin GM, Fox M, Mattson MP (1996) Increased activity-regulating and neuroprotective efficacy of alpha-secretase-derived secreted amyloid precursor protein conferred by a C-terminal heparin-binding domain. *J Neurochem* 67:1882–1896.
- Gao Y, Pimplikar SW (2001) The gamma-secretase-cleaved C-terminal fragment of amyloid precursor protein mediates signaling to the nucleus. *Proc Natl Acad Sci U S A* 98:14979–14984.
- Genzen JR, McGehee DS (2003) Short- and long-term enhancement of excitatory transmission in the spinal cord dorsal horn by nicotinic acetylcholine receptors. *Proc Natl Acad Sci U S A* 100:6807–6812.
- Girod R, Barazangi N, McGehee D, Role LW (2000) Facilitation of glutamatergic neurotransmission by presynaptic nicotinic acetylcholine receptors. *Neuropharmacology* 39:2715–2725.
- Gray R, Rajan AS, Radcliffe KA, Yakehiro M, Dani JA (1996) Hippocampal synaptic transmission enhanced by low concentrations of nicotine. *Nature* 383:713–716.
- Haass C, Selkoe DJ (2007) Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid beta-peptide. *Nat Rev Mol Cell Biol* 8:101–112.
- Ishida A, Furukawa K, Keller JN, Mattson MP (1997) Secreted form of beta-amyloid precursor protein shifts the frequency dependency for induction of LTD, and enhances LTP in hippocampal slices. *Neuroreport* 8:2133–2137.
- Ji D, Lape R, Dani JA (2001) Timing and location of nicotinic activity enhances or depresses hippocampal synaptic plasticity. *Neuron* 31:131–141.
- Jones S, Sudweeks S, Yakel JL (1999) Nicotinic receptors in the brain: correlating physiology with function. *Trends Neurosci* 22:555–561.
- Kamenetz F, Tomita T, Hsieh H, Seabrook G, Borchelt D, Iwatsubo T, Sisodia S, Malinow R (2003) APP processing and synaptic function. *Neuron* 37:925–937.
- Kametani F (2008) Epsilon-secretase: reduction of amyloid precursor protein epsilon-site cleavage in Alzheimer's disease. *Curr Alzheimer Res* 5:165–171.
- Kimberly WT, Zheng JB, Guénette SY, Selkoe DJ (2001) The intracellular domain of the beta-amyloid precursor protein is stabilized by Fe65 and translocates to the nucleus in a notch-like manner. *J Biol Chem* 276:40288–40292.
- Laird FM, Cai H, Savonenko AV, Farah MH, He K, Melnikova T, Wen H, Chiang HC, Xu G, Koliatsos VE, Borchelt DR, Price DL, Lee HK, Wong PC (2005) BACE1, a major determinant of selective vulnerability of the brain to amyloid- β amyloidogenesis, is essential for cognitive, emotional, and synaptic functions. *J Neurosci* 25:11693–11709.
- Leissring MA, Murphy MP, Mead TR, Akbari Y, Sugarman MC, Jannatipour M, Anliker B, Müller U, Saftig P, De Strooper B, Wolfe MS, Golde TE, LaFerla FM (2002) A physiologic signaling role for the gamma-secretase-derived intracellular fragment of APP. *Proc Natl Acad Sci U S A* 99:4697–4702.
- Levin ED, Simon BB (1998) Nicotinic acetylcholine involvement in cognitive function in animals. *Psychopharmacology (Berl)* 138:217–230.
- Lewczuk P, Beck G, Esselmann H, Bruckmoser R, Zimmermann R, Fiszler M, Bibl M, Maler JM, Kornhuber J, Wiltfang J (2006) Effect of sample collection tubes on cerebrospinal fluid concentrations of tau proteins and amyloid beta peptides. *Clin Chem* 52:332–334.
- Lisman J, Raghavachari S (2006) A unified model of the presynaptic and postsynaptic changes during LTP at CA1 synapses. *Sci STKE* 2006:re11.
- López-Toledano MA, Shelanski ML (2004) Neurogenic effect of β -amyloid peptide in the development of neural stem cells. *J Neurosci* 24:5439–5444.
- Ma H, Lesné S, Kotilinek L, Steidl-Nichols JV, Sherman M, Younkin L, Younkin S, Forster C, Sergeant N, Delacourte A, Vassar R, Citron M, Kofuji P, Boland LM, Ashe KH (2007) Involvement of beta-site APP cleaving enzyme 1 (BACE1) in amyloid precursor protein-mediated en-


- hancement of memory and activity-dependent synaptic plasticity. *Proc Natl Acad Sci U S A* 104:8167–8172.
- Mastrangelo P, Mathews PM, Chishti MA, Schmidt SD, Gu Y, Yang J, Mazzella MJ, Coomaraswamy J, Horne P, Strome B, Pelly H, Levesque G, Ebeling C, Jiang Y, Nixon RA, Rozmahel R, Fraser PE, St George-Hyslop P, Carlson GA, Westaway D (2005) Dissociated phenotypes in presenilin transgenic mice define functionally distinct gamma-secretases. *Proc Natl Acad Sci U S A* 102:8972–8977.
- Mattson MP (1994) Secreted forms of beta-amyloid precursor protein modulate dendrite outgrowth and calcium responses to glutamate in cultured embryonic hippocampal neurons. *J Neurobiol* 25:439–450.
- Meziane H, Dodart JC, Mathis C, Little S, Clemens J, Paul SM, Ungerer A (1998) Memory-enhancing effects of secreted forms of the beta-amyloid precursor protein in normal and amnesic mice. *Proc Natl Acad Sci U S A* 95:12683–12688.
- Mucke L, Masliah E, Johnson WB, Ruppe MD, Alford M, Rockenstein EM, Forss-Petter S, Pietropaolo M, Mallory M, Abraham CR (1994) Synaptotrophic effects of human amyloid beta protein precursors in the cortex of transgenic mice. *Brain Res* 666:151–167.
- Pawlik M, Sastre M, Calero M, Mathews PM, Schmidt SD, Nixon RA, Levy E (2004) Overexpression of human cystatin C in transgenic mice does not affect levels of endogenous brain amyloid beta peptide. *J Mol Neurosci* 22:13–18.
- Paxinos G (1998) Mouse brain in stereotaxic coordinates, Ed 2. New York: Academic.
- Phillips RG, LeDoux JE (1992) Differential contribution of amygdala and hippocampus to cued and contextual fear conditioning. *Behav Neurosci* 106:274–285.
- Phinney AL, Calhoun ME, Wolfer DP, Lipp HP, Zheng H, Jucker M (1999) No hippocampal neuron or synaptic bouton loss in learning-impaired aged beta-amyloid precursor protein-null mice. *Neuroscience* 90:1207–1216.
- Phinney AL, Drisaldi B, Schmidt SD, Lugowski S, Coronado V, Liang Y, Horne P, Yang J, Sekoulidis J, Coomaraswamy J, Chishti MA, Cox DW, Mathews PM, Nixon RA, Carlson GA, St George-Hyslop P, Westaway D (2003) In vivo reduction of amyloid-beta by a mutant copper transporter. *Proc Natl Acad Sci U S A* 100:14193–14198.
- Plant LD, Boyle JP, Smith IF, Peers C, Pearson HA (2003) The production of amyloid β peptide is a critical requirement for the viability of central neurons. *J Neurosci* 23:5531–5535.
- Puzzo D, Vitolo O, Trinchese F, Jacob JP, Palmeri A, Arancio O (2005) Amyloid- β peptide inhibits activation of the nitric oxide/cGMP/cAMP-responsive element-binding protein pathway during hippocampal synaptic plasticity. *J Neurosci* 25:6887–6897.
- Radcliffe KA, Dani JA (1998) Nicotinic stimulation produces multiple forms of increased glutamatergic synaptic transmission. *J Neurosci* 18:7075–7083.
- Rozmahel R, Huang J, Chen F, Liang Y, Nguyen V, Ikeda M, Levesque G, Yu G, Nishimura M, Mathews P, Schmidt SD, Mercken M, Bergeron C, Westaway D, St George-Hyslop P (2002) Normal brain development in PS1 hypomorphic mice with markedly reduced gamma-secretase cleavage of betaAPP. *Neurobiol Aging* 23:187–194.
- Saura CA, Choi SY, Beglopoulos V, Malkani S, Zhang D, Shankaranarayana Rao BS, Chattarji S, Kelleher RJ 3rd, Kandel ER, Duff K, Kirkwood A, Shen J (2004) Loss of presenilin function causes impairments of memory and synaptic plasticity followed by age-dependent neurodegeneration. *Neuron* 42:23–36.
- Schenk F, Morris RG (1985) Dissociation between components of spatial memory in rats after recovery from the effects of retrohippocampal lesions. *Exp Brain Res* 58:11–28.
- Schmidt SD, Nixon RA, Mathews PM (2005) ELISA method for measurement of amyloid-beta levels. *Methods Mol Biol* 299:279–297.
- Seabrook GR, Smith DW, Bowery BJ, Easter A, Reynolds T, Fitzjohn SM, Morton RA, Zheng H, Dawson GR, Sirinathsinghji DJ, Davies CH, Collingridge GL, Hill RG (1999) Mechanisms contributing to the deficits in hippocampal synaptic plasticity in mice lacking amyloid precursor protein. *Neuropharmacology* 38:349–359.
- Small DH, Mok SS, Bornstein JC (2001) Alzheimer's disease and Abeta toxicity: from top to bottom. *Nature reviews* 2:595–598.
- Small DH, Maksud D, Kerr ML, Ng J, Hou X, Chu C, Mehrani H, Unabia S, Azari MF, Loiacono R, Aguilar MI, Chebib M (2007) The beta-amyloid protein of Alzheimer's disease binds to membrane lipids but does not bind to the alpha7 nicotinic acetylcholine receptor. *J Neurochem* 101:1527–1538.
- Smith-Swintosky VL, Pettigrew LC, Craddock SD, Culwell AR, Rydel RE, Mattson MP (1994) Secreted forms of beta-amyloid precursor protein protect against ischemic brain injury. *J Neurochem* 63:781–784.
- Wang HY, Lee DH, D'Andrea MR, Peterson PA, Shank RP, Reitz AB (2000) beta-Amyloid(1–42) binds to alpha7 nicotinic acetylcholine receptor with high affinity. Implications for Alzheimer's disease pathology. *J Biol Chem* 275:5626–5632.
- Yankner BA, Duffy LK, Kirschner DA (1990) Neurotrophic and neurotoxic effects of amyloid beta protein: reversal by tachykinin neuropeptides. *Science* 250:279–282.
- Yao J, Petanceska SS, Montine TJ, Holtzman DM, Schmidt SD, Parker CA, Callahan MJ, Lipinski WJ, Bisgaier CL, Turner BA, Nixon RA, Martins RN, Ouimet C, Smith JD, Davies P, Laska E, Ehrlich ME, Walker LC, Mathews PM, Gandy S (2004) Aging, gender and APOE isotype modulate metabolism of Alzheimer's Abeta peptides and F-isoprostanines in the absence of detectable amyloid deposits. *J Neurochem* 90:1011–1018.
- Zucker RS, Regehr WG (2002) Short-term synaptic plasticity. *Annu Rev Physiol* 64:355–405.




Supplemental Fig. 1. Tris-Tricine PAGE Western blot analysis of A β preparation in non-denaturing/non-reducing conditions showing different bands corresponding to monomers (4.5 kDa), trimers (13.5 kDa) and tetramers (18 kDa).




Supplemental Fig. 2. Post-tetanic application of picomolar concentrations of Aβ₄₂ does not vary LTP. Perfusion of hippocampal slices with human Aβ₄₂ (200 pM) for 20 min after a theta-burst stimulation does not affect LTP, nor baseline transmission. The horizontal bar indicates the period during which Aβ₄₂ was added to the bath solution.




Supplemental Fig. 3. Levels of endogenous Aβ₄₂ are in the picomolar range. Aβ₄₂ was measured by sandwich ELISA in mouse cerebellum, cortex and hippocampus (n = 4 mice).



Supplemental Fig. 4. Cued conditioning is not affected by human A β_{42} . Bilateral injections of human A β_{42} (200 pM) or scramble A β_{42} (200 pM) into dorsal hippocampi, 15 min prior to training, do not affect freezing during cued conditioning.



Supplemental Fig. 5. Block of $\alpha 7$ -nAChRs does not affect PTP. **a**, Perfusion of hippocampal slices with MCL (3 μ M) in D-APV (50 μ M) does not affect PTP. The horizontal bars indicate the period during which MCL and APV were added to the bath solution. **b**, Perfusion of hippocampal slices with α -BgTx (0.1 μ M) in D-APV (50 μ M) does not affect PTP. The horizontal bars indicate the period during which α -BgTx and APV were added to the bath solution.



Supplemental Fig. 6. Schematic representation of our model indicating that neuronal activity (1) releases A β during exocytosis (2). A mechanism dependent upon activation of presynaptic α 7-nAChRs (3) would cause an increase in neurotransmitter release (4), boosting synaptic plasticity and memory (5).