



Varieties with quadratic entry locus, II

Paltin Ionescu and Francesco Russo

ABSTRACT

We continue the study, begun in [Rus07], of secant defective manifolds having ‘simple entry loci’. We prove that such manifolds are rational and describe them in terms of tangential projections. Using also the work of [IR07], their classification is reduced to the case of Fano manifolds of high index, whose Picard group is generated by the hyperplane section class. Conjecturally, the former should be linear sections of rational homogeneous manifolds. We also provide evidence that the classification of linearly normal dual defective manifolds with Picard group generated by the hyperplane section should follow along the same lines.

Introduction

An n -dimensional closed submanifold $X \subset \mathbb{P}^N$ is *secant defective* if its secant variety $SX \subseteq \mathbb{P}^N$ has dimension less than $2n + 1$, the expected dimension. The secant defect of X is then $\delta = \delta(X) := 2n + 1 - \dim(SX)$. Secant defective manifolds naturally fall into two categories. Manifolds admitting ‘non-trivial projections’ are those for which $SX \neq \mathbb{P}^N$, see [Sev01, Zak93], while manifolds ‘of small codimension’ correspond to the other case, $SX = \mathbb{P}^N$, see [BL72]. Let $x, y \in X$ be two general points and let p be a general point on the line $\langle x, y \rangle$. Consider the closure of the locus of secants to X passing through p . Its trace on X , denoted as Σ_p , is called the *entry locus* (with respect to p) and has dimension δ . Our aim is to find classification results for secant defective manifolds whose entry loci are simple enough. Consider embedded manifolds $X \subset \mathbb{P}^N$ as above, such that through two general points $x, y \in X$ there passes an r -dimensional quadric hypersurface, say Q^r , contained in X . Observe that $Q^r \subseteq \Sigma_p$ if $p \in \langle x, y \rangle$; in particular $r \leq \delta$, see [KS02]. When $r = 1$, we call such manifolds *conic-connected* (CCMs for short). The extremal case $r = \delta$ was called ‘manifolds with *local quadratic entry locus*’ (LQELMs), while the special case when $Q^r = \Sigma_p$ was named ‘manifolds with *quadratic entry locus*’ (QELMs); see [KS02, Rus07, IR07]. Being rationally connected, these special classes of secant defective manifolds may be studied in the context of Mori theory; see [Mor79, Mor82, KMM92] and also [Deb01, Kol96, Hwa01]. The interested reader can find further motivation and various examples in the introduction to [Rus07]. The present paper continues the line of investigation started in [Rus07]; see also [IR07]. We acknowledge once more our intellectual debt to the classical work by Scorza; see [Sco08, Sco09]. The content of the paper is described below.

Tangential projections turned out to be a basic tool in the classical works (by Severi, Scorza, Terracini, and others) on secant defective manifolds. In what follows, by the δ -partial tangential projection we mean projection from a δ -codimensional linear subspace (passing through x) of the projective tangent space at $x \in X$. In Theorem 2.1 we show that the generic δ -partial tangential

Received 23 March 2007, accepted in final form 11 January 2008.

2000 Mathematics Subject Classification 14N05, 14M99, 14J45.

Keywords: secant defective, quadratic entry locus manifold, tangential projection, conic-connected, dual defective.

The first author is partially supported by the Italian Programme ‘Incentivazione alla mobilità di studiosi stranieri e italiani residenti all’estero’. The second author is partially supported by CNPq (Centro Nacional de Pesquisa), Grants 300961/2003-0, 308745/2006-0 and 474475/2006-9, and by PRONEX/FAPERJ–Algebra Comutativa e Geometria Algebrica.

This journal is © [Foundation Compositio Mathematica](#) 2008.

projection of a QELM is birational. The proof is based on a degeneration technique introduced in [CMR04] (see also [CR06]). The key point is that the general fibre of the (full) tangential projection is a degeneration of the general entry locus. Conversely, if the generic δ -partial tangential projection of X is birational, then X is a LQELM. Moreover, we show that defective LQELMs are rational, extending the theorem on the rationality of ‘manifolds with one apparent double point’, proved in [CMR04]. In § 3 we compare various properties of LQELMs and CCMs; see Proposition 3.2. Examples show that the former class is much larger than the first. In particular, using results by Hwang and Kebekus [HK05] (see also [IR07]), we see that complete intersections (of high dimension with respect to the multidegree) are CCM, while the only LQELM which are complete intersections are the hyperquadrics; see Proposition 3.4. We also characterize LQELMs (when $\delta \geq 3$) in terms of the projective geometry of the variety of lines passing through a general point and contained in X ; see Proposition 3.5 and [Rus07, Theorem 2.3]. The main properties of LQELMs are collected in Theorem 3.7. In particular, their classification is reduced to the case when the Picard group is generated by the class of the hyperplane section. Conjecturally, QELMs with $\text{Pic}(X) \cong \mathbb{Z}\langle \mathcal{O}_X(1) \rangle$ are linear sections of rational homogeneous manifolds (which are completely classified); see Remark 3.8. The last section contains two applications. The first one, due to Fu [Fu08], gives a substantial improvement of one of the main results from [KS02] and is based on ideas developed in [Rus07, IR07] and the present paper. The other application is a new proof of the classification, due to Ein [Ein86], of manifolds with small dual. To the best of our knowledge, this is the first time that the classification of dual defective manifolds has been connected to (and, conjecturally, even reduced to, cf. Remark 4.5) that of secant defective manifolds, via the key concept of (L)QELM.

1. Preliminaries

We work over the field of complex numbers. Notation and terminology are the same as in [Rus07]; we recall below some of the relevant facts.

Let $X \subset \mathbb{P}^N$ be an irreducible non-degenerate projective variety of dimension n . Let

$$SX = \overline{\bigcup_{\substack{x \neq y \\ x, y \in X}} \langle x, y \rangle} \subseteq \mathbb{P}^N$$

be the secant variety to X ; see also the construction in (3).

Clearly $\dim(SX) \leq \min\{N, 2n + 1\}$. If $\dim(SX) < 2n + 1$, then $X \subset \mathbb{P}^N$ is said to be *secant defective*. The *secant defect* of $X \subset \mathbb{P}^N$ is equal to $\delta(X) := 2n + 1 - \dim(SX)$.

For $p \in SX \setminus X$, the closure of the locus of couples of distinct points of X spanning secant lines passing through p is called *the entry locus of X with respect to $p \in SX$* and it will be denoted by $\Sigma_p(X)$. The closure of the locus of secant lines to X passing through p is a cone over $\Sigma_p(X)$; let us call it $C_p(X)$. If $X \subset \mathbb{P}^N$ is smooth, then $\Sigma_p(X) = C_p(X) \cap X$ as schemes for general $p \in SX$; see, for example, [FR81, Lemma 4.5]. Moreover, it is easy to see that for general $p \in SX$, $\Sigma_p(X)$ is equidimensional of dimension equal to $\delta(X)$. Thus $\dim(C_p(X)) = \delta(X) + 1$. In general, $\Sigma_p(X)$ may be reducible.

Let $X \subset \mathbb{P}^N$ be an irreducible non-degenerate projective manifold of dimension n and secant defect $\delta \geq 0$. (1)

In the following definition, we consider varieties having the *simplest entry locus*.

DEFINITION 1.1 (cf. also [KS02, Rus07, IR07]). Let X be as in (1).

- (i) X is said to be a *quadratic entry locus manifold of type $\delta \geq 0$* , briefly a *QELM of type δ* , if for general $p \in SX$ the entry locus $\Sigma_p(X)$ is a quadric hypersurface of dimension δ .

- (ii) X is said to be a *local quadratic entry locus manifold of type* $\delta \geq 0$, briefly a *LQELM of type* δ , if through two general points there passes a quadric hypersurface of dimension δ contained in X .
- (iii) X is said to be a *conic-connected manifold*, briefly a *CCM*, if through two general points of X there passes an irreducible conic contained in X .

Note that, for $\delta = 0$, being a LQELM imposes no restriction on X . Clearly, any QELM is LQEL and any defective LQELM is CC.

LEMMA 1.2 (cf. Lemma 1.2 in [Rus07]). *Let X be a LQELM with $\delta > 0$ and let $x, y \in X$ be general points. There is a unique quadric hypersurface of dimension δ , say $Q_{x,y}$, passing through x, y and contained in X . Moreover, $Q_{x,y}$ is irreducible.*

The following proposition is easy to prove.

PROPOSITION 1.3. *Let $X \subset \mathbb{P}^N$ be an irreducible non-degenerate smooth projective variety.*

- (i) *If $X' \subset \mathbb{P}^M$, $M \leq N - 1$, is an isomorphic projection of X , then X' is a LQELM if and only if X is a LQELM.*
- (ii) *If X is a (L)QELM of type $\delta \geq 1$, then a general hyperplane section is a (L)QELM of type $\delta - 1$.*
- (iii) *If X is a QELM and $SX = \mathbb{P}^N$, then X is linearly normal.*

Proof. Parts (i) and (ii) are standard and left to the reader, so that we prove only part (iii). Suppose that $X \subset \mathbb{P}^N$ were the isomorphic projection of $\overline{X} \subset \mathbb{P}^{N+1}$ from a point $q \in \mathbb{P}^{N+1} \setminus S\overline{X}$. Since $N = \dim(SX) = \dim(S\overline{X})$, $S\overline{X}$ would be a hypersurface of degree at least two. A general $p \in SX$ would be the projection from q of at least two different points p_1, p_2 belonging to $\langle q, p \rangle \cap S\overline{X}$. Then the projections of the entry loci $\Sigma_{p_1}(\overline{X})$ and $\Sigma_{p_2}(\overline{X})$ yield two irreducible components of $\Sigma_p(X)$. This is a contradiction, since $\Sigma_p(X)$ is a smooth, hence irreducible, δ -dimensional quadric hypersurface; see [FR81, pp. 964–966]. \square

If $x \in X \subset \mathbb{P}^N$ is a smooth point, we denote by $\mathbf{T}_x X$ the affine Zariski tangent space at x and by $T_x X$ its projective closure in \mathbb{P}^N .

The dimension of the image of the projection of $X \subset \mathbb{P}^N$ from a general tangent space to X , called the *tangential projection* of $X \subset \mathbb{P}^N$, is easily computed via the Terracini lemma (see, for example, [Rus07, § 1]). Let $x \in X \subset \mathbb{P}^N$ be a general point and let

$$\pi_x : X \dashrightarrow W_x \subseteq \mathbb{P}^{N-n-1} \tag{2}$$

be the projection of X from $T_x X$. We have $\dim(W_x) = n - \delta$, so that a general fiber of π_x is of pure dimension δ .

LEMMA 1.4 (cf. Lemma 1.6 in [Rus07]). *Let $X \subset \mathbb{P}^N$ be a smooth irreducible non-degenerate variety, and assume that $\delta > 0$. The irreducible components of the closure of a general fibre of π_x are not linear.*

2. Tangential projections and the geometry of LQELMs

The main result of this section is the following theorem, generalizing [CMR04, Corollary 4.2], where the case of QELMs of type $\delta = 0$ in \mathbb{P}^{2n+1} , i.e. of *varieties with one apparent double point*, was considered.

THEOREM 2.1. *Let $X \subset \mathbb{P}^N$ be as in (1) and let $x \in X$ be a general point. Then the following hold.*

- (i) If X is a QELM of type $\delta \geq 0$, the projection from a general δ -codimensional subspace of $T_x X$ passing through x is birational onto its image. In particular, if $SX = \mathbb{P}^N$, then X is rational.
- (ii) Conversely, if the projection from a general δ -codimensional subspace of $T_x X$ passing through x is birational, then X is a LQELM of type δ .
- (iii) If $X \subset \mathbb{P}^N$ is a LQELM of type $\delta > 0$, then X is rational.

We recall some results from [CMR04] (see also [CR06]) in order to point out a relation between the entry loci of a variety and a general fiber of the tangential projection. We believe this relation, contained in Proposition 2.2 below, is interesting in itself.

Let $X \subset \mathbb{P}^N$ be an irreducible non-degenerate projective variety and let

$$S_X := \overline{\{(x, y, z) \mid x \neq y, z \in \langle x, y \rangle\}} \subset X \times X \times \mathbb{P}^N$$

be the *abstract secant variety* of $X \subset \mathbb{P}^N$, which is an irreducible projective variety of dimension $2n + 1$. Let us consider the projections of S_X onto the factors $X \times X$ and \mathbb{P}^N .

$$\begin{array}{ccc}
 & S_X & \\
 p_1 \swarrow & & \searrow p_2 \\
 X \times X & & \mathbb{P}^N
 \end{array} \tag{3}$$

With this notation we get

$$p_2(S_X) = \overline{\bigcup_{\substack{x \neq y \\ x, y \in X}} \langle x, y \rangle} = SX \subseteq \mathbb{P}^N.$$

Let $L = \langle x, y \rangle$ with $x \in X$ and $y \in X$ general points, i.e. L is a general secant line to X , and let $p \in \langle x, y \rangle \subseteq SX \subseteq \mathbb{P}^N$ be a general point. We fix coordinates on L so that the coordinate of x is 0; let U be an open subset of $\mathbb{A}_{\mathbb{C}}^1 \subset L$ containing $0 = x$. Let $p_2 : S_X \rightarrow SX \subseteq \mathbb{P}^N$ be as above and let

$$Z_U = p_2^{-1}(U) \subset S_X.$$

By changing U , if necessary, we can suppose that $p_2 : Z_U \rightarrow U$ is flat over $U \setminus \{0\}$ and that $\dim(Z_U)_t = \delta(X)$ for every $t \neq 0$. The projection of $p_1((Z_U)_t)$ onto one of the factors is Σ_t , the entry locus of X with respect to t for every $t \neq 0$.

Moreover, by definition, a point $(r, s) \in X \times X$, $r \neq s$, belongs to $(Z_U)_t$, $t \neq 0$, if and only if $t \in \langle r, s \rangle$, that is if and only if $(r, s) \in p_2^{-1}(t)$. Thus, if $\psi_t : X \dashrightarrow \mathbb{P}^{N-1}$ is the projection from t onto a disjoint \mathbb{P}^{N-1} , we can also suppose that ψ_t is a morphism for every $t \neq 0$ and a rational map not defined at $x = 0$ for $t = 0$. The above analysis says that the *abstract entry locus* $(Z_U)_t$, $t \neq 0$, can be considered as the closure in $X \times X$ of the double point locus scheme of ψ_t , minus the diagonal $\Delta_X \subset X \times X$.

Let $T = \langle T_x X, y \rangle$, so that T is a general \mathbb{P}^{n+1} containing $T_x X$ and a general point $y \in X$. By definition $\pi_x^{-1}(\pi_x(y)) = T \cap X \setminus (T_x X \cap X)$, with the notation as in (2). Let

$$F_y = \overline{\pi_x^{-1}(\pi_x(y))}$$

be the closure of the fiber of π_x through y . Every irreducible component of F_y has dimension $\delta(X)$ by the Terracini lemma and by the generality of y ; see the discussion after (2). Generic smoothness ensures also that there exists only one irreducible component of F_y through y .

By using the same ideas as in [CMR04], we have the following result, not explicitly stated in *loc. cit.*, because a slightly different degeneration was considered. For more details about the construction recalled above and below, we refer to [CMR04, §§ 3 and 4] and [CR06, § 2].

PROPOSITION 2.2. *Let the notation be as above. The closure of the fiber of π_x through a general point $y \in X$ is contained in the flat limit of the family $\{(Z_U)_t\}_{t \neq 0}$. In other words, the closure of a general fiber of the tangential projection is a degeneration of the general entry locus of X .*

Proof. We shall look at ψ_t as a family of morphisms and study the limit of the *double point scheme* $(Z_U)_t$.

Consider the products $\mathcal{X} = X \times U$ and $\mathbb{P}_U = \mathbb{P}^{N-1} \times U$. The projections ψ_t , for $t \in U$, fit together to give a rational map $\psi : \mathcal{X} \dashrightarrow \mathbb{P}_U$, which is defined everywhere except at the pair $(x, 0)$. In order to extend the projection not defined at $x \in X$, we have to blow up \mathcal{X} at $(x, 0)$. Let $\sigma : \tilde{\mathcal{X}} \rightarrow \mathcal{X}$ be this blowing-up and let $Z \simeq \mathbb{P}^n$ be the exceptional divisor. Looking at the obvious morphism $\varphi : \tilde{\mathcal{X}} \rightarrow U$, we see that this is a flat family of varieties over U . The fiber $\tilde{\mathcal{X}}_t$ over a point $t \in U \setminus \{0\}$ is isomorphic to X , whereas the fiber $\tilde{\mathcal{X}}_0$ over $t = 0$ is of the form $\tilde{\mathcal{X}}_0 = \tilde{X} \cup Z$, where $\tilde{X} \rightarrow X$ is the blowing-up of X at x , and $\tilde{X} \cap Z = E$ is the exceptional divisor of this blowing-up, the intersection being transverse. Reasoning as in [CMR04, Lemma 3.1], it is easy to see that ψ_0 acts on \tilde{X} as the projection from the point $0 = x$, while it maps Z isomorphically onto the linear space $\psi_0(T) = \mathbb{P}^n$. This immediately implies that every point of $T \cap X$, different from x , appears in the ‘double point scheme’ of $\psi_0 : \tilde{X} \cup Z \rightarrow \mathbb{P}^{N-1}$. Therefore F_y , being of dimension $\delta(X)$, is contained in the flat limit of $\{(Z_U)_t\}_{t \neq 0}$, proving the assertion. \square

For an irreducible variety $X \subset \mathbb{P}^N$ we denote by $\mu(X)$ the number of secant lines passing through a general point of SX . If $\delta(X) > 0$, then $\mu(X)$ is infinite, while for $\delta(X) = 0$ the above number is finite and in this case

$$\nu(X) = \mu(X) \cdot \deg(SX)$$

is called *the number of apparent double points* of $X \subset \mathbb{P}^N$. With these definitions we obtain the following generalization of [CMR04, Theorem 4.1] (see also [CR06, Theorem 2.7]).

THEOREM 2.3. *Let $X \subset \mathbb{P}^N$ be as in (1). If $\delta(X) = 0$, then*

$$0 < \deg(\pi_x) \leq \mu(X).$$

In particular, for a QELM of type $\delta = 0$, the general tangential projection is birational.

If $X \subset \mathbb{P}^N$ is a QELM of type $\delta > 0$, then the general fiber of π_x is irreducible. More precisely the closure of the fiber of π_x passing through a general point $y \in X$ is the entry locus of a general point $p \in \langle x, y \rangle$, i.e. a smooth quadric hypersurface.

Proof. If $\delta(X) = 0$, then for $t \in U \setminus \{0\}$ the zero-dimensional scheme $(Z_U)_t$ has length equal to $2\mu(X)$. The zero-dimensional scheme F_y contains $\deg(\pi_x)$ isolated points, yielding $2 \deg(\pi_x)$ points in the flat limit of $\{(Z_U)_t\}_{t \neq 0}$ by Proposition 2.2 and proving the first part.

Suppose X is a QELM of type $\delta > 0$. Then for every $t \neq 0$ the δ -dimensional scheme $(Z_U)_t$ is a smooth quadric hypersurface by definition of QELMs. The fiber F_y contains the entry locus Σ_p of a general point $p \in \langle x, y \rangle$, which is a smooth quadric hypersurface of dimension δ passing through x and y . By Proposition 2.2 the variety F_y is also contained in the flat limit of $\{(Z_U)_t\}_{t \neq 0}$. Therefore, F_y coincides with Σ_p . In fact, in this case the family $\{(Z_U)_t\}_{t \neq 0}$ is constant. \square

Proof of Theorem 2.1. Suppose that $X \subset \mathbb{P}^N$ is a QELM of type $\delta \geq 0$. If $\delta = 0$ then the first part of Theorem 2.3 yields that π_x is birational onto its image (see also [CMR04, Corollary 4.2]).

Suppose from now on that $\delta > 0$. The projection from a general codimension δ linear subspace $L \subseteq T_x X$ passing through x is a rational map $\pi_L : X \dashrightarrow \mathbb{P}^{N-n+\delta-1}$. For general $y \in X$, the linear space $\langle L, y \rangle$ is obtained by cutting $\langle T_x X, y \rangle$ with δ general hyperplanes H_1^y, \dots, H_δ^y passing through x and y . From the second part of Theorem 2.3, it follows that, for general $y \in X$, $\overline{\pi_x^{-1}(\pi_x(y))}$ is a

δ -dimensional quadric $Q_{x,y}$. Then

$$\pi_L^{-1}(\pi_L(y)) \subseteq Q_{x,y} \cap H_1^y \cap \cdots \cap H_\delta^y = \{x, y\},$$

yielding the birationality of π_L onto its image. If $SX = \mathbb{P}^N$, then $N - n + \delta - 1 = n$. Part (i) is now completely proved.

Suppose we are in the hypothesis of part (ii). Let $L = \mathbb{P}^{n-\delta} \subset T_x X$ be a general linear subspace passing through x . Then L is the tangent space of a general codimension δ linear section of $X \subset \mathbb{P}^N$ passing through x , let us say Z . Thus the restriction of π_L to Z , $\pi_{L|Z} : Z \dashrightarrow \mathbb{P}^{N-\delta-\dim(Z)-1} = \mathbb{P}^{N-n-1}$, is the projection from $L = T_x Z$. Since π_L restricted to X is birational onto its image, then $\pi_{L|Z}$ is easily seen to also be birational onto its image. Moreover, looking at $\pi_{L|Z}$ as the projection from $T_x Z$, we get $\pi_{L|Z}(Z) = \pi_x(X) = W_x \subseteq \mathbb{P}^{N-n-1}$. Each irreducible component of a general fiber of π_x produces at least one point in a general fiber of $\pi_{L|Z}$. Hence $\pi_x : X \dashrightarrow W_x \subseteq \mathbb{P}^{N-n-1}$ has irreducible general fibers of dimension δ by the birationality of $\pi_{L|Z}$.

We shall prove inductively that $F_y = \overline{\pi_x^{-1}(\pi_x(y))}$ is a δ -dimensional quadric for general $y \in X$. So there is no loss of generality in supposing that $\delta = 1$, by passing to a general linear section; see Proposition 1.3. We claim that set theoretically $L \cap F_y = \{x\}$. We have $F_y \subset \langle T_x X, y \rangle$ so that $T_x X \cap F_y$ consists of a finite number of points. By the generality of L we get $L \cap F_y \subseteq \{x\}$. Let $t = \pi_x(y)$. Since $\langle L, t \rangle$ is a hyperplane in $\langle T_x X, y \rangle = \langle T_x X, t \rangle$, intersecting $\pi_x^{-1}(\pi_x(y))$ transversally at a unique point, we get that either we are in the case of the claim or F_y is a line. This last case is excluded by Lemma 1.4.

Let $q = \pi_{L|Z}^{-1}(t) = \langle L, t \rangle \cap Z \setminus (L \cap Z)$. By definition,

$$\langle L, t \rangle = \langle L, q \rangle. \tag{4}$$

Consider the projection from t onto $T_x X$, let us say $\psi_t : \langle T_x X, t \rangle \dashrightarrow T_x X$. Let $\tilde{F}_y = \psi_t(F_y)$. By definition $x \in \tilde{F}_y$ because $x \in F_y$. Moreover, we claim that $L \cap \tilde{F}_y$ is supported at x , so that \tilde{F}_y is a line through x . Indeed, if $z \in L \cap \tilde{F}_y$, then there exists $w \in \langle z, t \rangle \cap F_y \subset \langle L, t \rangle \cap F_y = \langle L, q \rangle \cap F_y$, where the last equality follows from (4). Thus either $w = x$ or $w = q$ and in any case $x = \psi_t(w) = z$. Therefore, $F_y \subset \langle \tilde{F}_y, t \rangle \simeq \mathbb{P}^2$; moreover, the Trisecant lemma implies that the line $\langle x, y \rangle$ cuts transversally X , hence also *a fortiori* F_y , at x and at y . The line $\langle x, y \rangle$ is contained in the plane $\langle \tilde{F}_y, t \rangle$, so that $\deg(F_y) = 2$ and F_y is a smooth conic passing through x and y , concluding the proof.

Let us prove part (iii). Fix a general point $x \in X$ and denote by \mathcal{Q}_x (an irreducible component of) the family of δ -dimensional quadric hypersurfaces contained in X and passing through x . Let $\pi : \mathcal{F}_x \rightarrow \mathcal{Q}_x$ be the universal family and let $\varphi : \mathcal{F}_x \rightarrow X$ be the tautological morphism.

Assume first that $\delta = 1$. Then φ is surjective by definition of a LQELM and birational by Lemma 1.2. Note that π has a section, corresponding to the point x . Denote by $\mathcal{E} \subset \mathcal{F}_x$ the image of this section. Consider the blowing-up $\sigma : X' \rightarrow X$ of X at x . Since both \mathcal{Q}_x and π are generically smooth, the birational map $\psi = \sigma^{-1} \circ \varphi$ is defined at a general point of \mathcal{E} . Moreover, as \mathcal{E} is contracted by φ to the point x , then ψ sends \mathcal{E} to E , the exceptional divisor of σ . So we have the following diagram.

$$\begin{array}{ccc} \mathcal{F}_x & \xrightarrow{\psi} & X' \\ \pi \downarrow & \searrow \varphi & \downarrow \sigma \\ \mathcal{Q}_x & & X \end{array}$$

By the Zariski main theorem the map ψ^{-1} is defined at a general point of E , and hence ψ induces a birational map $\psi_0 : \mathcal{E} \dashrightarrow E = \mathbb{P}^{n-1}$. Thus \mathcal{F}_x is birational to \mathbb{P}^n , being birational to a family of conics with a section over a rational base, and X is a rational variety, as claimed. A general version of the above argument appears in [IN03, Proposition 3.1].

Suppose now that $\delta \geq 2$ and fix H a general hyperplane section of X through x . Using Lemma 1.2, we see that sending a quadric hypersurface through x to its trace on H yields a birational map between the families $\mathcal{Q}_x(X)$ and $\mathcal{Q}_x(H)$. So, we see inductively that $\mathcal{Q}_x(X)$ is a rational variety of dimension $n - \delta$. Therefore, \mathcal{F}_x is rational, as the family $\mathcal{F}_x \rightarrow \mathcal{Q}_x$ has a section. Being birational to \mathcal{F}_x by Lemma 1.2, X is rational too. \square

Remark 2.4. Bronowski claims in [Bro32] that $X \subset \mathbb{P}^{2n+1}$ is a variety with one apparent double point if and only if the projection of X to \mathbb{P}^n from a general tangent space to X is a birational map; he also formulates a generalization to arbitrary secant k -planes; see [CR06] for more details. Unfortunately, Bronowski's argument is unclear and we do not know of any convincing proof for this statement. Ciliberto *et al.* [CMR04, Corollary 4.2] (see also Theorem 2.1) prove one implication showing the rationality of varieties with one apparent double point. The open implication would be a very useful tool for constructing examples.

We can generalize the *Bronowski conjecture* to the following: a smooth irreducible n -dimensional variety $X \subset \mathbb{P}^{2n+1-\delta(X)}$ is a QELM if and only if the projection from a general codimension $\delta(X)$ linear subspace of $T_x X$ passing through x is birational. Theorem 2.1 proves one implication, yielding the rationality of QELMs and extending [CMR04, Corollary 4.2]. One may consult [CR06] for other generalizations of the above conjecture to higher secant varieties.

It is worth mentioning that the above results reveal the following interesting picture for the tangential projections of QELMs of type $\delta \geq 0$ with $SX = \mathbb{P}^N$: for $\delta = 0$ we project from the whole space and we have varieties with one apparent double point; at the other extreme we found the stereographic projection of quadric hypersurfaces, the only QELMs of type equal to their dimension.

3. LQEL versus CC

We recall the following definition from [BBI00].

DEFINITION 3.1 [BBI00]. A smooth rational curve $C \subset X$, where X is a projective manifold of dimension n , is a *quasi-line* if $N_{C|X} \simeq \bigoplus_1^{n-1} \mathcal{O}_{\mathbb{P}^1}(1)$.

The relation between the notions LQELM and CCM is clarified in the following.

PROPOSITION 3.2. Let $X \subset \mathbb{P}^N$ be a CCM of secant defect δ . Let $C = C_{x,y}$ be a general conic through the general points $x, y \in X$ and let c be the point representing C in the Hilbert scheme of X . Let \mathcal{C}_x be the unique irreducible component of the Hilbert scheme of conics passing through x which contains the point c .

- (i) We have $n + \delta \geq -K_X \cdot C = \dim(\mathcal{C}_x) + 2 \geq n + 1$.
- (ii) The equality $-K_X \cdot C = n + \delta$ holds if and only if $X \subset \mathbb{P}^N$ is a LQELM.
- (iii) The following conditions are equivalent:
 - (a) $\dim(\mathcal{C}_x) = n - 1$;
 - (b) C is a quasi-line;
 - (c) all conics through x, y are non-degenerate.
- (iv) If $\delta \geq 3$, then $X \subset \mathbb{P}^N$ is a Fano manifold with $\text{Pic}(X) \simeq \mathbb{Z}\langle \mathcal{O}_X(1) \rangle$ and index $i(X) = (\dim(\mathcal{C}_x)/2) + 1$.

Proof. We have the universal family $g : \mathcal{F}_x \rightarrow \mathcal{C}_x$ and the tautological morphism $f : \mathcal{F}_x \rightarrow X$, which is surjective. Since $C \in \mathcal{C}_x$ is a general conic and since $x \in X$ is general, we get $\dim(\mathcal{C}_x) = -K_X \cdot C - 2$.

Take a general point $y \in X$ and a general $p \in \langle x, y \rangle$. The conics passing through x and y are parameterized by $g(f^{-1}(y))$, which has pure dimension

$$\dim(\mathcal{F}_x) - n = \dim(\mathcal{C}_x) + 1 - n = -K_X \cdot C - 1 - n.$$

We claim that the locus of conics through x and y , denoted by $\mathcal{L}_{x,y}$, has dimension $-K_X \cdot C - n$ and is clearly contained in the irreducible component of the entry locus (with respect to p) through x and y . Indeed, conics through x, y and another general point $z \in \mathcal{L}_{x,y}$ have to be finitely many. Otherwise, their locus would fill up the plane $\langle x, y, z \rangle$ and this would imply that the line $\langle x, y \rangle$ is contained in X . However, we have excluded linear spaces from the definition of CCMs and LQELMs. Therefore, $\delta \geq -K_X \cdot C - n$, that is, $-K_X \cdot C \leq n + \delta$.

The locus of conics through x and y is contained in $\langle T_x X, y \rangle \cap \langle x, T_y X \rangle$, which is a linear space of dimension $\delta + 1$. Indeed, by the Terracini lemma,

$$\dim(T_x X \cap T_y X) = 2n - \dim(SX) = \delta - 1.$$

By the Trisecant lemma, $x \notin T_y X$ and $y \notin T_x X$, so that

$$\dim(\langle T_x X, y \rangle \cap \langle x, T_y X \rangle) = \delta + 1.$$

If $-K_X \cdot C = n + \delta$, then, for $p \in \langle x, y \rangle$ general, the irreducible component $\Sigma_{x,y}^p$ of the entry locus passing through x and y coincides with the locus of conics through x and y . Thus $\Sigma_{x,y}^p$ is a quadric hypersurface by the Trisecant lemma and by the generality of x and y (if $\delta = n$, $X \subset \mathbb{P}^{n+1}$ is a quadric hypersurface). So, part (ii) is proved.

Next we see part (iii). Conditions (a) and (b) are equivalent, since the normal bundle of C in X is ample, of degree $\dim(\mathcal{C}_x)$.

Assume that condition (a) holds. The dimension of the subfamily consisting of reducible conics from \mathcal{C}_x has dimension at most $n - 2$. Hence their locus is of dimension at most $n - 1$ and does not contain the general point y . So we have condition (c).

Assume that condition (c) holds. Then, by bend and break there are finitely many conics through x and y , giving condition (a).

Finally, part (iv) follows from the Barth–Larsen theorem, the fact that X contains moving conics and part (i). \square

Examples 3.3. (i) For $n \geq 3$, let X be a smooth cubic hypersurface in \mathbb{P}^{n+1} or the smooth complete intersection of two hyperquadrics in \mathbb{P}^{n+2} . Use, for example, [BBI00, Theorem 3.2] and induction on n to see that X is CC. In the first case, X is a hypersurface of degree three so it cannot be a LQELM. In the second case, $\delta(X) = n - 1$; as the Picard group is generated by the hyperplane section, X cannot contain hyperquadrics of dimension $n - 1$.

(ii) CCMs $X \subset \mathbb{P}^{N+1}$ of secant defect $\delta(X) = \delta - 1 \geq 2$, constructed from QELMs $Z \subset \mathbb{P}^N$ of type $\delta \geq 3$.

Let $Z \subset \mathbb{P}^N$ be a QELM of type $\delta \geq 3$. Consider \mathbb{P}^N as a hyperplane in \mathbb{P}^{N+1} , take $q \in \mathbb{P}^{N+1} \setminus \mathbb{P}^N$ and let $W = C_q(Z) \subset \mathbb{P}^{N+1}$ be the cone over Z of vertex q . Let $X = C_q(Z) \cap Q \subset \mathbb{P}^{N+1}$, where $Q \subset \mathbb{P}^{N+1}$ is a general quadric hypersurface. Then $SX = C_q(SZ)$, yielding $\delta(X) = \delta(Z) - 1 \geq 2$. Moreover, $\text{Pic}(X) \simeq \mathbb{Z}\langle \mathcal{O}_X(1) \rangle$ and $X \subset \mathbb{P}^{N+1}$ is a CCM of index $i(X) = (n + \delta(X) - 1)/2$.

Indeed, let $\pi : T = \mathbb{P}(\mathcal{O}_Z \oplus \mathcal{O}_Z(1)) \rightarrow Z$ and let $E \subset T$ be the section at infinity of π . If $\varphi : T = \mathbb{P}(\mathcal{O}_Z \oplus \mathcal{O}_Z(1)) \rightarrow \mathbb{P}^{N+1}$ is the tautological morphism given by (a sublinear system of) $|\mathcal{O}_T(1)|$, then $\varphi(T) = W$ and $X \subset \mathbb{P}^{N+1}$ can be naturally thought of as an element of $|\mathcal{O}_T(2)|$. Recall that φ restricts to an isomorphism between $T \setminus E$ and $W \setminus \{q\}$. By adjunction we get that

$\omega_X = \mathcal{O}_X(1 - i(Z))$, that is,

$$i(X) = i(Z) - 1 = \frac{n + \delta}{2} - 1 = \frac{n + \delta(X) - 1}{2}.$$

In particular, $X \subset \mathbb{P}^{N+1}$ is a Fano manifold. Adjunction formula also says that the double cover $\pi_q : X \rightarrow Z$, induced by the projection from q , $\pi_q : \mathbb{P}^{N+1} \setminus \{q\} \dashrightarrow \mathbb{P}^N$, is ramified along a hyperplane section of $Z \subset \mathbb{P}^N$.

Take two general points $x, y \in X$ and let $x' = \pi_q(x)$ and $y' = \pi_q(y)$. Through x' and y' there passes a smooth quadric hypersurface $Q' \subset \mathbb{P}^{\delta+1}$ of dimension $\delta \geq 3$. Let $\tilde{Q} = C_q(Q') \subset \mathbb{P}^{\delta+2}$ and let $\tilde{X} = C_q(Q') \cap Q \subset \mathbb{P}^{\delta+2}$. Then $\tilde{X} \subset \mathbb{P}^{\delta+2}$ is a (smooth) complete intersection of two quadric hypersurfaces passing through x and y , so that $X \subset \mathbb{P}^{N+1}$ is a CCM since $\tilde{X} \subset \mathbb{P}^{\delta+2}$ is CC by part (i) above.

(iii) When $\delta = 1$, CCMs and LQELMs are the same. In this case a general conic through two general points is a quasi-line. Examples are easy to construct, e.g. $X = G \cap H_1 \cap H_2 \cap H_3$, where G is the Grassmannian of lines in \mathbb{P}^m , $m \geq 4$, and H_i are general hyperplane sections of its Plücker embedding.

(iv) Assume that $\delta = 2$ and X is CC. Let C be a general conic passing through two general points. Then X is a LQELM if and only if C is not a quasi-line; see Proposition 3.2. Examples of the LQEL case are given by taking $X = G \cap H_1 \cap H_2$, where G and H_i are as above. Examples of the case where X contains quasi-lines are got by applying the construction in part (ii) above starting with $Z = G \cap H$, a hyperplane section of the same Grassmannian.

Many examples of CCMs which are not LQEL come from the following.

PROPOSITION 3.4 (cf. Corollary 2.5 in [IR07]).

- (i) If $X \subset \mathbb{P}^{n+r}$ is a smooth non-degenerate complete intersection of multi-degree (d_1, d_2, \dots, d_r) with $n > 3(\sum_1^r d_i - r - 1)$, then X is a CCM.
- (ii) If X is a secant defective LQELM and a complete intersection, then X is a hyperquadric.

Proof. Part (i) is exactly [IR07, Corollary 2.5].

We show part (ii). If $n = 2$ use [Rus07, Proposition 3.3] to conclude. If $n \geq 3$, by the Lefschetz theorem, we have $\text{Pic}(X) \cong \mathbb{Z}\langle \mathcal{O}_X(1) \rangle$ and Proposition 3.2 gives $i(X) = (n + \delta)/2$. Let $r = N - n$. We have $2n + 1 - \delta = \dim(SX) \leq N = n + r$, so $\delta \geq n + 1 - r$. Assuming X to be a complete intersection of type (d_1, \dots, d_r) , with $d_i \geq 2$ for all i , we get $i(X) = n + r + 1 - \sum_1^r d_i$; hence

$$n + 2r + 2 - 2 \sum_1^r d_i = 2i(X) - n = \delta \geq n + 1 - r \quad \text{and} \quad 4r \leq 2 \sum_1^r d_i \leq 3r + 1,$$

so $r = 1$. □

The proof of the following criterion for recognizing LQELMs illustrates the role of CCMs. Recall from [Rus07] that, if $x \in X$ is a general point, we denote by $Y_x \subset \mathbb{P}(\mathbf{T}_x^* X)$ the variety of lines through x , contained in X .

PROPOSITION 3.5. Let $X \subset \mathbb{P}^N$ be as in (1) and assume that $\delta \geq 3$. The following assertions are equivalent:

- (i) X is a LQELM;
- (ii) if $x \in X$ is a general point, then $\dim(Y_x) \geq (n + \delta)/2 - 2$ and $SY_x = \mathbb{P}^{n-1}$.

Proof. The fact that part (i) implies (ii) was shown in [Rus07, Theorem 2.3] and, in fact, equality holds in part (ii).

Assume part (ii). $\delta \geq 3$ implies via the Barth–Larsen theorem that $\text{Pic}(X) \cong \mathbb{Z}\langle \mathcal{O}_X(1) \rangle$. Moreover,

$$i(X) = \dim(Y_x) + 2 \geq \frac{n + \delta}{2} \geq \frac{n + 3}{2}.$$

So we may apply [HK05, Theorem 3.14] to deduce that X is a CCM. In the notation of Proposition 3.2, we have $2i(X) = -K_X \cdot C \geq n + \delta$. Therefore, X is a LQELM by combining parts (i) and (ii) of Proposition 3.2. \square

The following proposition is an application of [Rus07, Theorem 2.3].

PROPOSITION 3.6. *Assume that X is a LQELM of type $\delta < n$.*

- (i) *If $\delta > n/2$ then $\delta \leq 6$ and $n \leq 10$. These cases are classified in [Rus07, Corollary 3.1].*
- (ii) *If $\delta > n/3$ then $\delta \leq 10$ and $n \leq 26$. These cases may also be classified by the same method.*
- (iii) *If $k \geq 4$, $\delta > n/k$, then $\delta \leq 2(k + 2)$.*

Proof. See [Rus07, Corollary 3.1] for part (i).

Assume $n/3 < \delta \leq n/2$. Using [Rus07, Theorem 2.8], we find $\delta \leq 10$ and the following possibilities for the pairs (n, δ) : $(2, 1), (4, 2), (5, 2), (7, 3), (8, 4), (10, 4), (13, 5), (14, 6), (15, 7), (16, 8), (25, 9)$ and $(26, 10)$.

To prove part (iii) we proceed by induction on $k \geq 3$. We may suppose that $\delta \geq 10$. Assume that we have $n/k \geq \delta > n/(k + 1)$ and $k \geq 4$. In the notation of [Rus07, Theorem 2.4], we let $X_1 = Y_x$, $\delta_1 = \delta(X_1) = \delta - 2$, $n_1 = \dim(X_1) = (n + \delta)/2 - 2$. If $\delta_1 > n_1/k$ we get $\delta \leq 2(k + 3)$, completing the induction. If $\delta_1 \leq n_1/k$ it follows that

$$\delta \leq \frac{n + 4(k - 1)}{2k - 1}, \quad \text{so} \quad \frac{n}{k + 1} < \frac{n + 4(k - 1)}{2k - 1}.$$

This gives

$$10k \leq \delta k \leq n < \frac{4(k^2 - 1)}{k - 2},$$

so $3k^2 - 10k + 2 < 0$. Therefore, $k = 3$; a contradiction. \square

Consider the following list of examples of QELMs.

- (i) $\nu_2(\mathbb{P}^n) \subset \mathbb{P}^{n(n+3)/2}$.
- (ii) The projection of $\nu_2(\mathbb{P}^n)$ from the linear space $\langle \nu_2(\mathbb{P}^s) \rangle$, where $\mathbb{P}^s \subset \mathbb{P}^n$ is a linear subspace; equivalently $X \simeq \text{Bl}_{\mathbb{P}^s}(\mathbb{P}^n)$ embedded in \mathbb{P}^N by the linear system of quadric hypersurfaces of \mathbb{P}^n passing through \mathbb{P}^s ; alternatively $X \simeq \mathbb{P}_{\mathbb{P}^r}(\mathcal{E})$ with $\mathcal{E} \simeq \mathcal{O}_{\mathbb{P}^r}(1)^{\oplus n-r} \oplus \mathcal{O}_{\mathbb{P}^r}(2)$, $r = 1, 2, \dots, n-1$, embedded by $|\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)|$. Here $N = n(n+3)/2 - \binom{s+2}{2}$ and s is an integer such that $0 \leq s \leq n-2$.
- (iii) A hyperplane section of the Segre embedding $\mathbb{P}^a \times \mathbb{P}^b \subset \mathbb{P}^{N+1}$. Here $n \geq 3$ and $N = ab + a + b - 1$, where $a \geq 2$ and $b \geq 2$ are such that $a + b = n + 1$.
- (iv) $\mathbb{P}^a \times \mathbb{P}^b \subset \mathbb{P}^{ab+a+b}$ Segre embedded, where a, b are positive integers such that $a + b = n$.

The essential properties of LQELMs are collected in the next theorem. It follows by putting together Theorem 2.1(iii), [IR07, Theorem 2.2] and [Rus07, Theorem 2.3(4d)].

THEOREM 3.7. *Let $X \subset \mathbb{P}^N$ be a defective LQELM. Then X is Fano and rational. Moreover, either X is an isomorphic projection of one of the manifolds listed immediately above or $\text{Pic}(X) \cong \mathbb{Z}\langle \mathcal{O}_X(1) \rangle$ and $i(X) = (n + \delta)/2$. If $\delta \geq 3$, then $Y_x \subset \mathbb{P}^{n-1}$ is a QELM of type $\delta - 2$, dimension $(n + \delta)/2 - 2$ and such that $SY_x = \mathbb{P}^{n-1}$.*

Remark 3.8. Via the above theorem, the classification of secant defective QELMs is reduced to the case where the Picard group is \mathbb{Z} . Let us say that $X \subset \mathbb{P}^N$ is *maximal* if X is not a hyperplane section of some (non-degenerate) manifold $X' \subset \mathbb{P}^{N+1}$. The following tempting conjecture would, if true, lead to a complete classification of defective QELMs.

CONJECTURE. Any maximal defective QELM with Picard group \mathbb{Z} is homogeneous.

Note that homogeneous manifolds arising from irreducible representations of (semi)simple complex algebraic groups are QELMs (see [Zak93, ch. III] and [Kaj99]), and the secant defective ones are classified completely (see *loc. cit.*). Moreover, the results in [Rus07, §3] confirm the conjecture for $\delta > n/2$. The next two finiteness results for QELMs with $\delta < n$ would follow from the above conjecture:

- (i) $\delta \leq 8$;
- (ii) if $\delta \geq 5$, then $n \leq 16$.

Proposition 3.6 may be seen as supporting these finiteness expectations. Also, in [Fu08, Corollary 2], it is proved that $\delta \leq (n + 8)/3$ holds for any LQELM with $\delta < n$. In particular, result (i) follows from result (ii).

4. Two applications

The first application is due to Fu [Fu08], who found it by relying on the ideas and techniques from [Rus07], [IR07] and Proposition 3.2. We mention his result in order to illustrate the usefulness of our point of view.

THEOREM 4.1 (cf. Theorem 2 in [Fu08]). *Let X be as in (1) and assume that X is swept out by hyperquadrics of dimension greater than $[n/2] + 1$, all passing through a fixed point $x \in X$. Then X is a hyperquadric.*

This result substantially improves the main application in [KS02], with a much shorter proof.

Our second application concerns the classification of manifolds with small duals.

For an irreducible variety $Z \subset \mathbb{P}^N$, we define $\text{def}(Z) = N - 1 - \dim(Z^*)$ as the dual defect of $Z \subset \mathbb{P}^N$, where $Z^* \subset \mathbb{P}^{N*}$ is the dual variety of $Z \subset \mathbb{P}^N$. In [Ein86, Theorem 2.4] it is proved that if $\text{def}(X) > 0$, then $\text{def}(X) \equiv n \pmod{2}$, a result usually attributed to Landman; see the Introduction of [Ein86]. Moreover, the Zak theorem on tangencies implies that $\dim(X^*) \geq \dim(X)$ for a smooth non-degenerate variety $X \subset \mathbb{P}^N$; see [Zak93, I.2.5].

We combine the geometry of CCMs and LQELMs to give a new proof of [Ein86, Theorem 4.5]. Our approach avoids the use of Beilinson spectral sequences and more sophisticated computations as in [Ein86, Lemmas 4.2, 4.3 and Theorem 4.4].

We begin by recalling some basic facts from [Ein86].

PROPOSITION 4.2. *Let $X \subset \mathbb{P}^N$ be as in (1) and assume that $\text{def}(X) > 0$. Then:*

- (i) [Ein86, Theorem 2.4] *through a general point $x \in X$ there passes a line $L_x \subset X$ such that $-K_X \cdot L_x = (n + \text{def}(X) + 2)/2$, so that $\text{def}(X) \equiv n \pmod{2}$;*
- (ii) [Ein86, Theorem 3.2] *$\text{def}(X) = n - 2$ if and only if $X \subset \mathbb{P}^N$ is a scroll over a smooth curve, i.e. it is a \mathbb{P}^{n-1} -bundle over a smooth curve, whose fibers are linearly embedded.*

The following proposition reinterprets the result of Hwang and Kebekus [HK05, Theorem 3.14] on Fano manifolds with large index.

PROPOSITION 4.3 (cf. [HK05]; see also Proposition 2.4 in [IR07]). *Let $X \subset \mathbb{P}^N$ be as in (1). Assume that X is a Fano manifold with $\text{Pic}(X) \cong \mathbb{Z}\langle \mathcal{O}_X(1) \rangle$ and let $x \in X$ be a general point.*

- (i) *If $i(X) > 2n/3$, then X is a CCM with $\delta > n/3$.*
- (ii) *If $\text{def}(X) > 0$ and $\text{def}(X) > (n - 6)/3$, then X is a CCM with $\delta \geq \text{def}(X) + 2$. Moreover, if $\delta = \text{def}(X) + 2$, then X is a LQELM of type $\delta = \text{def}(X) + 2$.*
- (iii) *If X is a LQELM of type δ and $\text{def}(X) > 0$, then $\delta = \text{def}(X) + 2$.*

Proof. See [IR07, Proposition 2.4] for the proof of part (i).

In the hypothesis of part (ii), [Ein86, Theorem 2.4] yields $i(X) = (n + \text{def}(X) + 2)/2 > 2n/3$ so that X is a CCM by part (i). Proposition 3.2 yields $\delta \geq \text{def}(X) + 2$ and also the remaining assertions of parts (ii) and (iii). \square

We recall that according to the Hartshorne conjecture, if $n > \frac{2}{3}N$, then $X \subset \mathbb{P}^N$ should be a complete intersection and that complete intersections have no dual defect. Thus, assuming the Hartshorne conjecture, the following result yields the complete list of manifolds $X \subset \mathbb{P}^N$ such that $\dim(X^*) = \dim(X)$. The second part says that under the LQEL hypothesis the same results hold without any restriction (see also Remark 4.5 below).

THEOREM 4.4. *Let $X \subset \mathbb{P}^N$ be as in (1) and assume that $\dim(X) = \dim(X^*)$.*

- (i) [Ein86, Theorem 4.5] *If $N \geq 3n/2$, then X is projectively equivalent to one of the following:*
 - (a) *a smooth hypersurface $X \subset \mathbb{P}^{n+1}$, $n = 1, 2$;*
 - (b) *a Segre variety $\mathbb{P}^1 \times \mathbb{P}^{n-1} \subset \mathbb{P}^{2n-1}$;*
 - (c) *the Plücker embedding $\mathbb{G}(1, 4) \subset \mathbb{P}^9$;*
 - (d) *the 10-dimensional spinor variety $S^{10} \subset \mathbb{P}^{15}$.*
- (ii) *If X is a LQELM, then it is projectively equivalent either to a smooth quadric hypersurface $Q \subset \mathbb{P}^{n+1}$ or to a variety as in (b), (c), (d) above.*

Proof. Clearly $\text{def}(X) = 0$ if and only if $X \subset \mathbb{P}^{n+1}$ is a hypersurface, giving respectively case (a) or that of a smooth quadric hypersurface. From now on we suppose that $\text{def}(X) > 0$ and hence $n \geq 3$. By parts (i) and (ii) of Proposition 4.2, $\text{def}(X) = n - 2$ and $N = 2n - 1$ if and only if we are in case (b); see also [Ein86, Theorem 3.3, part c].

Thus, we may assume that $0 < \text{def}(X) \leq n - 4$, that is, $N \leq 2n - 3$. Therefore, $\delta \geq 4$ and X is a Fano manifold with $\text{Pic}(X) \cong \mathbb{Z}\langle \mathcal{O}_X(1) \rangle$. Moreover, in case (i), $\text{def}(X) = N - n - 1 > (n - 6)/3$ by hypothesis. Thus Proposition 4.3 yields that X is also a CCM with $\delta \geq \text{def}(X) + 2$. Also taking into account the last part of Proposition 4.3, from now on we can suppose that X is a CCM with $\delta \geq \text{def}(X) + 2 \geq 3$.

We have $n - \delta \leq N - 1 - n = \text{def}(X) \leq \delta - 2$, that is, $\delta \geq n/2 + 1$. The Zak linear normality theorem implies $SX = \mathbb{P}^N$, so that

$$N = \dim(SX) = 2n + 1 - \delta \leq \frac{3n}{2}.$$

Since $N \geq 3n/2$, we get $N = 3n/2$, $\delta = n/2 + 1 = \text{def}(X) + 2$ and n even. Therefore, X is a LQELM of type $\delta = n/2 + 1$ by Proposition 4.3. Using [Rus07, Corollary 3.1] concludes the proof, yielding cases (c) and (d). \square

Remark 4.5. Let $X \subset \mathbb{P}^N$ be as in (1). Assume that $\text{def}(X) > 0$ and $\text{Pic}(X) \cong \mathbb{Z}\langle \mathcal{O}_X(1) \rangle$. We conjecture that X is a LQELM and even a QELM if, moreover, it is assumed to be linearly normal. Combined with Remark 3.8, this would imply that maximal dual defective manifolds with $\text{Pic}(X) \cong \mathbb{Z}\langle \mathcal{O}_X(1) \rangle$ are homogeneous, as already conjectured in [BS95].

Our last application is the following.

PROPOSITION 4.6. *Let $X \subset \mathbb{P}^N$ be as in (1), with $\text{Pic}(X) \cong \mathbb{Z}\langle \mathcal{O}_X(1) \rangle$ and $\text{def}(X) > 0$. Assume, moreover, that $\text{def}(X) > (n - 6)/3$ and $N \geq (5n + 2)/3$. Then $SX \neq \mathbb{P}^N$.*

Proof. By Proposition 4.3, X is a CCM. Assume that $SX = \mathbb{P}^N$. We get

$$N = \dim(SX) = 2n + 1 - \delta \geq \frac{5n + 2}{3},$$

so $\delta \leq (n + 1)/3$. It follows that $3 \text{def}(X) \geq n - 5 \geq 3\delta - 6$. By Proposition 4.3, X is a LQELM of type $\delta = (n + 1)/3$ and $\text{def}(X) = \delta - 2$. From the list in the proof of Proposition 3.6(ii) it follows that $\delta \leq 2$, so $\text{def}(X) = 0$. This is a contradiction. \square

ACKNOWLEDGEMENTS

Both authors are grateful to the organizers of the Conference ‘Projective varieties with unexpected properties’, that took place in Siena, 8–13 June 2004. Our collaboration started while taking part in this very pleasant and fruitful mathematical event.

REFERENCES

- BBI00 L. Bădescu, M. C. Beltrametti and P. Ionescu, *Almost-lines and quasi-lines on projective manifolds*, in *Complex Analysis and Algebraic Geometry (in memory of Michael Schneider)* (de Gruyter, Berlin, 2000), 1–27.
- BL72 W. Barth and M. E. Larsen, *On the homotopy groups of complex projective algebraic manifolds*, *Math. Scand.* **30** (1972), 88–94.
- BS95 M. C. Beltrametti and A. J. Sommese, *The adjunction theory of complex projective varieties*, de Gruyter Exp. Math., vol. 16 (de Gruyter, Berlin, 1995).
- Bro32 J. Bronowski, *The sum of powers as canonical expressions*, *Proc. Cambridge Philos. Soc.* **29** (1932), 69–82.
- CMR04 C. Ciliberto, M. Mella and F. Russo, *Varieties with one apparent double point*, *J. Algebraic Geom.* **13** (2004), 475–512.
- CR06 C. Ciliberto and F. Russo, *Varieties with minimal secant degree and linear systems of maximal dimension on surfaces*, *Adv. Math.* **200** (2006), 1–50.
- Deb01 O. Debarre, *Higher-Dimensional Algebraic Geometry*, Universitext (Springer, Berlin, 2001).
- Ein86 L. Ein, *Varieties with small dual variety. I*, *Invent. Math.* **86** (1986), 63–74.
- Fu08 B. Fu, *Inductive characterizations of hyperquadrics*, *Math. Ann.* **340** (2008), 185–194.
- FR81 T. Fujita and J. Roberts, *Varieties with small secant varieties: the extremal case*, *Amer. J. Math.* **103** (1981), 953–976.
- Hwa01 J. M. Hwang, *Geometry of minimal rational curves on Fano manifolds*, in *School on Vanishing Theorems and Effective Results in Algebraic Geometry*, Trieste, 2000, ICTP Lecture Notes, vol. 6 (Abdus Salam Int. Cent. Theoret. Phys., 2001), 335–393.
- HK05 J. M. Hwang and S. Kebekus, *Geometry of chains of minimal rational curves*, *J. reine angew. Math.* **584** (2005), 173–194.
- IN03 P. Ionescu and D. Naie, *Rationality properties of manifolds containing quasi-lines*, *Internat. J. Math.* **14** (2003), 1053–1080.
- IR07 P. Ionescu and F. Russo, *Conic-connected manifolds*, Preprint (2006), math. AG/0701885.
- Kaj99 H. Kaji, *Homogeneous projective varieties with degenerate secants*, *Trans. Amer. Math. Soc.* **351** (1999), 533–545.
- Kol96 J. Kollár, *Rational Curves on Algebraic Varieties*, *Ergeb. Math. Grenzgeb. (3)*, vol. 32 (Springer, Berlin, 1996).

- KMM92 J. Kollár, Y. Miyaoka and S. Mori, *Rationally connected varieties*, J. Algebraic Geom. **1** (1992), 429–448.
- KS02 Y. Kachi and E. Sato, *Segre’s reflexivity and an inductive characterization of hyperquadrics*, Mem. Amer. Math. Soc. **160** (2002).
- Mor79 S. Mori, *Projective manifolds with ample tangent bundle*, Ann. of Math. (2) **110** (1979), 593–606.
- Mor82 S. Mori, *Threefolds whose canonical bundles are not numerically effective*, Ann. of Math. (2) **116** (1982), 133–176.
- Rus07 F. Russo, *Varieties with quadratic entry locus. I*, Preprint (2006), math. AG/0701889.
- Sco08 G. Scorza, *Sulla determinazione delle varietà a tre dimensioni di S_r ($r \geq 7$) i cui S_3 tangenti si tagliano a due a due*, Rend. Circ. Mat. Palermo **25** (1908), 193–204; contained in *Opere Scelte*, vol. I (Edizioni Cremonese, 1960), 167–182.
- Sco09 G. Scorza, *Sulle varietà a quattro dimensioni di S_r ($r \geq 9$) i cui S_4 tangenti si tagliano a due a due*, Rend. Circ. Mat. Palermo **27** (1909), 148–178; contained in *Opere Scelte*, vol. I (Ed. Cremonese, 1960), 252–292.
- Sev01 F. Severi, *Intorno ai punti doppi impropri di una superficie generale dello spazio a quattro dimensioni e ai suoi punti tripli apparenti*, Rend. Circ. Mat. Palermo **15** (1901), 33–51.
- Zak93 F. L. Zak, *Tangents and secants of algebraic varieties*, Translations in Mathematical Monographs, vol. 127 (American Mathematical Society, Providence, RI, 1993).

Paltin Ionescu ionescu@dim.unige.it

University of Bucharest and Institute of Mathematics of the Romanian Academy

Current address: Dipartimento di Matematica, Università degli Studi di Genova, Via Dodecaneso, 35, 16146 Genova, Italy

Francesco Russo frusso@dmi.unict.it

Dipartimento di Matematica e Informatica, Università degli Studi di Catania, Viale A. Doria 6, 95125 Catania, Italy