

# Vitamin E is the major contributor to the antioxidant capacity in lambs fed whole dried citrus pulp

G. Luciano<sup>1†</sup>, V. Roscini<sup>1</sup>, S. Mattioli<sup>1</sup>, S. Ruggeri<sup>1</sup>, R. S. Gravador<sup>2</sup>, A. Natalello<sup>3</sup>, M. Lanza<sup>3</sup>, A. De Angelis<sup>3</sup> and A. Priolo<sup>3</sup>

<sup>1</sup>Dipartimento di Scienze Agrarie Alimentari e Ambientali, Università degli Studi di Perugia, Borgo XX Giugno 74, 06121, Perugia, Italy; <sup>2</sup>School of Agriculture and Food Science, University College Dublin, Belfield, Dublin 4, Dublin, Ireland; <sup>3</sup>Dipartimento di Agricoltura, Alimentazione e Ambiente, Università degli Studi di Catania, Via Valdisavoia 5, 95123, Catania, Italy

(Received 13 November 2015; Accepted 4 July 2016; First published online 11 August 2016)

*The aim of this study was to investigate the effect of dietary whole dried citrus pulp (DCP) on the antioxidant status of lamb tissues. In total, 17 lambs were divided into two groups and fed for 56 days: a barley-based concentrate diet (CON – eight animals), or a concentrate-based diet including 35% DCP to partially replace barley (CIT – nine animals). The CIT diet contained a double concentration of phenolic compounds than the CON diet (7.9 v. 4.0 g/kg dry matter (DM), respectively), but had no effect ( $P > 0.05$ ) on the overall antioxidant capacity of the hydrophilic fraction of blood plasma, liver and muscle. The CIT diet contained clearly more  $\alpha$ -tocopherol than the CON diet (45.7 v. 10.3 mg/kg DM), which could explain the higher concentration of  $\alpha$ -tocopherol in liver, plasma and muscle ( $P < 0.05$ ). The dietary treatment had no effect on the extent of lipid peroxidation, measured as thiobarbituric acid and reactive substances assay (TBARS values) in the faeces, small intestine, liver, plasma and muscle. Nevertheless, when muscle homogenates were incubated in the presence of  $Fe^{3+}$ /ascorbate to induce lipid peroxidation, the muscle from lambs fed DCP displayed lower TBARS values ( $P < 0.01$ ), which negatively correlated with the concentration of  $\alpha$ -tocopherol in muscle. These results showed that feeding whole DCP to ruminants increases the antioxidant status of muscle through an increase in the deposition of  $\alpha$ -tocopherol.*

**Keywords:** citrus pulp, phenolic compounds,  $\alpha$ -tocopherol, ruminants, antioxidant

## Implications

In several areas, dried citrus pulp (DCP) may conveniently replace conventional cereals in diets for ruminants. Feeding DCP to small ruminants could improve important quality traits of meat, such as the resistance to oxidative deterioration. Citrus pulp (CP) notoriously contains phenolic compounds with antioxidant properties. Moreover, high levels of lipophilic antioxidants, such as vitamin E, may occur in whole DCP. This study demonstrates, for the first time, that vitamin E contributes most to the improvement of the antioxidant capacity of tissues from lambs fed whole DCP and highlights that the standardization of CP is needed to optimize its positive effects.

## Introduction

Citrus fruits represent one of the most important agricultural productions in several areas worldwide where they are destined to human consumption as fresh fruits or processed juice.

In the latter case, a substantial amount of waste biomass is produced, known as whole CP, composed of peels, inside portions of the fruit and seeds. The CP can be processed by pressing and drying the biomass to obtain the dried citrus pulp (DCP), which can be used in the formulation of diets for ruminants to replace high levels (30% or higher) of cereals (Bampidis and Robinson, 2006). Feeding CP to ruminants can result in positive effects related to the presence of residual bioactive molecules. Citrus fruits have received special attention for their content of phenolic compounds and it has been shown that citrus by-products are richer in polyphenols than the portions consumed by humans (Balasundram *et al.*, 2006; Nogata *et al.*, 2006). Recently, we have demonstrated that feeding lambs with concentrate-based diets including up to 35% DCP increased the resistance of muscle lipids and proteins to oxidative deterioration (Gravador *et al.*, 2014; Inserra *et al.*, 2014). The results of these studies were discussed in the light of the possible antioxidant effects of the polyphenols present in CP, in agreement with results provided by studies in which sheep were fed with either citrus extracts or with purified citrus flavonoids (Gladine *et al.*, 2007c; Bodas *et al.*, 2012).

<sup>†</sup> E-mail: giuseppe.luciano@unipg.it

However, although citrus flavonoids clearly possess antioxidant properties, research has not yet clarified their bioavailability and their mechanisms of action when ingested as components of the diet of animals, including humans. Indeed, it is known that dietary flavonoids are poorly absorbed in the gastrointestinal tract and, if absorbed, can be actively metabolized in the animal organism (Khan *et al.*, 2014). Therefore, possible indirect effects have been proposed to explain their potential antioxidant activities *in vivo*, such as the interaction with other antioxidants, or possible antioxidant activities exerted in the gastrointestinal tract (Halliwell *et al.*, 2005). This is, for example, the case of dietary high molecular weight condensed tannins, which have been shown to increase the antioxidant capacity of sheep liver and muscle through unknown mechanisms that do not apparently involve their absorption and deposition in the tissues (Luciano *et al.*, 2011; López-Andrés *et al.*, 2013). Moreover, when CP is fed to ruminants, it should be stressed that compounds other than polyphenols can greatly contribute to the total intake of bioactive molecules. Indeed, if whole CP does not undergo further processing before drying, it may contain remarkable levels of essential oils and tocopherols occurring in peels and seeds (Fontanel, 2013).

In the light of the above, the aim of this study was to understand the antioxidant effects of dietary whole DCP in lamb tissues. We hypothesize (i) that citrus polyphenols could increase the overall antioxidant capacity of animal tissues and (ii) that residual lipophilic antioxidants, such as vitamin E, in whole DCP might contribute to increasing the antioxidant status of the animal tissues. Moreover, possible effects of antioxidants present in the DCP might be exerted in the gastrointestinal tract of the animals, leading to a reduction of oxidative damages at the level of the intestine (Lindley *et al.*, 1994). To test these hypotheses, we used the same lambs from the experiment of Inserra *et al.* (2014) and we measured the antioxidant status of the lipophilic and hydrophilic fractions, as well as the extent of lipid oxidation, in the liver, blood plasma and muscle. Furthermore, lipid oxidation was measured in the small intestine and in the faeces.

## Material and methods

### Animals and diets

The experiment was conducted from February to May 2012 and the experimental protocol was approved by the European Community under the Seventh Framework Programme for Research, Technological Development and Demonstration Activities (Integrated Project LOW-INPUTBREEDS FP7-CP-IP 222623). All the procedures involving animals were conducted by specialized personnel. In total, 17 male Comisana lambs, born within 10 days, were selected. At 90 days of age, lambs were weighed (mean BW 19.12 kg  $\pm$  SD 4.26 kg) and individually penned indoor. Animals were randomly assigned to two dietary treatments and, for 10 days were adapted to the experimental diets. Subsequently, for 56 days, lambs were fed a barley-based concentrate diet (CON – eight animals) or a concentrate-based

**Table 1** Ingredients and chemical composition of the experimental diets

Ingredients (g/100 g of fresh weight)	CON	CIT
Barley	60	23
Soybean meal	9	13
Dehydrated alfalfa	20	20
Wheat bran	11	9
Whole dried citrus pulp	0	35
Chemical composition		
DM (g/100 g of fresh weight)	88.9	90.6
CP (g/100 g DM)	18.0	17.8
NDF (g/100 g DM)	34.6	33.1
Crude fat (g/100 g DM)	1.5	2.3
Metabolizable energy (MJ/kg DM)	10.4	10.5
Total phenolic compounds (g/kg DM)	4.0	7.9
$\alpha$ -tocopherol (mg/kg DM)	10.3	45.7

CON = control barley-based concentrate diet; CIT = concentrate diet including 35% (as-fed) whole dried citrus pulp; DM = dry matter.

diet including 35% as-fed whole DCP (including CP, peel and seeds) in partial replacement of barley (CIT – nine animals). The diets were formulated to be isonitrogenous and isoenergetic and no vitamin E ( $\alpha$ -tocopheryl acetate or  $\alpha$ -tocopherol) was added. The ingredients and chemical composition of the diets are reported in Table 1. All the ingredients were ground (5-mm screen) and mixed to avoid selection. Each day, the diets were given to lambs at 0900 h and the feed was continuously available in the feeders until 1800 h, after which feeders were removed. For each animal, the amount of feed offered and refused was recorded daily. Water was continuously available. The weight of the lambs was recorded weekly at 0800 h before supplying fresh feed.

### Sampling procedures

Samples of the feedstuffs were collected four times during the trial, finely milled (0.5-mm screen), vacuum-packed and stored at  $-30^{\circ}\text{C}$ . A sample of each diet was finally prepared for analyses by mixing equal amounts of the subsamples collected during the trial. Before 3 days of the end of the trial, faeces (~15 g) were collected directly from the rectum of each lamb, vacuum-packaged and stored at  $-30^{\circ}\text{C}$ . Before 2 days of slaughtering, after 12 h of fasting, blood (~8 ml) was sampled from the jugular vein of each lamb and collected into a heparin tube. The tubes were kept refrigerated at  $4^{\circ}\text{C}$ , centrifuged for 10 min at  $1500 \times g$  at  $4^{\circ}\text{C}$  and the plasma was stored at  $-80^{\circ}\text{C}$ . At the end of the experiment, lambs were transported to a commercial abattoir and slaughtered. The small intestine (20 cm length) was sampled ~5 cm after the pylorus and stored at  $-80^{\circ}\text{C}$ . Samples of liver and of *longissimus thoracis et lumborum* muscle (~150 g) were collected from the carcasses within 20 min from slaughtering, vacuum-packed and stored at  $-80^{\circ}\text{C}$ .

### Analyses of feeds

Dry matter (DM) and CP of the experimental diets were determined according to the Association of Official Analytical Chemists (1990) procedures. The NDF was analysed according to Van Soest *et al.* (1991) and the fat was

extracted according to procedure of Folch *et al.* (1957). The metabolizable energy was calculated using the 'ASSIST.T' software, version 1.3.1 developed by CRPA spa, Italy ([www.crupa.it/assist](http://www.crupa.it/assist)). Total phenolic compounds were assessed according to Makkar *et al.* (1993) with an extraction from feedstuffs using aqueous acetone (70%, v/v) and a subsequent analysis by Folin–Ciocalteu reagent. Standard solutions of gallic acid (GA) were used to express the phenolic compounds as g of GA equivalents/kg of DM. The procedure described by Schüep and Rettenmaier (1994) was adapted to the measurement of vitamin E ( $\alpha$ -tocopherol) in feedstuffs. In brief, feed samples were homogenized with an aqueous butylated hydroxytoluene (BHT) solution (0.06%). Vitamin E was extracted three times using hexane/ethyl acetate, dried under  $N_2$  and dissolved with acetonitrile. The analysis was performed using a Perkin Elmer series 200 HPLC (Perkin Elmer, Italia, Milan, Italy), equipped with an autosampler (Jasco, model AS 950-10, Tokyo, Japan) and an Ultrasphere ODS column (250  $\times$  4.6 mm internal diameter, 5  $\mu$ m particles size; CPS Analytica, Milan, Italy). The flow rate was 1.5 ml/min and  $\alpha$ -tocopherol was identified using a FD detector (Jasco, model FP-1525) set at excitation and emission wavelengths of 295 and 328 nm, respectively). Quantification was based on external calibration curves prepared with varying amounts of pure  $\alpha$ -tocopherol (Sigma-Aldrich, Bornem, Belgium) in ethanol.

#### *Measurement of the antioxidant status of liver, plasma and muscle*

The radical scavenging and the reducing capacity was measured in plasma and in aqueous extracts from liver and muscle by means of the trolox equivalent antioxidant capacity (TEAC), the ferric reducing antioxidant power (FRAP) and the Folin–Ciocalteu assays in order to measure the contribution of hydrophilic antioxidants to the antioxidant status of the tissues. The liver (0.5 g) and muscle (1 g) were homogenized with 10 ml distilled water, centrifuged at 3000  $\times$  g for 20 min at 4°C and the supernatant was filtered using Whatman No. 541 filter paper. Plasma was appropriately diluted with distilled water for analyses. The TEAC, FRAP and Folin–Ciocalteu assays were performed using the methods described by Luciano *et al.* (2011) and López-Andrés *et al.* (2013).

Vitamin E ( $\alpha$ -tocopherol) was determined in the lipid fraction extracted from liver, plasma and muscle as a measurement of the antioxidant capacity of the lipophilic fraction. The  $\alpha$ -tocopherol in muscle and liver samples was extracted according to Schüep and Rettenmaier (1994) with some modifications, whereas  $\alpha$ -tocopherol was extracted from blood plasma according to Zaspel and Csallany (1983). In brief, liver and muscle (2 g) were homogenized with aqueous BHT (0.06%), saponified with ethanolic KOH (60%) at 70°C for 30 min and extracted with hexane/ethyl acetate (9:1, v/v). Plasma (0.5 ml) was saponified in 1 M ethanolic KOH at 50°C for 1 h, sonicated and extracted two times with 4 ml of *n*-hexane containing BHT (200 mg/l). The extracts were dried under  $N_2$  stream and resuspended with

acetonitrile. The HPLC analysis of  $\alpha$ -tocopherol was performed as described above for feedstuffs.

#### *Lipid oxidation measurements*

Lipid oxidation in faeces was measured using the TBARS assay described by Pierre *et al.* (2003) with modifications as follows. Faecal water was prepared by homogenizing 1.5 g of faeces with 10 ml of cold (2°C) distilled water for 60 s. The homogenates were incubated for 1 h at 37°C with thorough mixing every 15 min. Subsequently, the homogenates were centrifuged for 10 min at 2500  $\times$  g at 20°C and the supernatant (2 ml) was further centrifuged for 20 min at 20 000  $\times$  g at 20°C. In glass test tube, 100  $\mu$ l of supernatant was diluted with 100  $\mu$ l of distilled water. SDS (200  $\mu$ l, 8.1%, w/v) and 2 ml of thiobarbituric acid reagent (0.5%, w/v thiobarbituric acid in 10% v/v acetic acid) were added. For each sample, a blank was prepared by replacing the thiobarbituric acid reagent with 2 ml of 10% acetic acid. Samples were incubated at 95°C for 60 min, after which the absorbance was measured at 532 nm.

The TBARS assay described by Siu and Draper (1978) was adapted to the measurement of lipid oxidation in the small intestine and plasma as follows. The small intestine was dissected longitudinally, rinsed with distilled water and cut finely using a scalpel. The intestine (1 g), was homogenized with 10 ml of cold distilled water (2°C) for 60 s. Then, 10 ml trichloroacetic acid (10%, w/v) was added, samples were vortex-mixed and centrifuged at 3000  $\times$  g for 20 min at 4°C and the supernatants were filtered using Whatman No. 541 filter paper. In a glass test tube, 1 ml of 0.06 M aqueous thiobarbituric acid was added to 4 ml of filtrate. The samples were incubated at 80°C for 90 min, after which the absorbance was measured at 532 nm. Blood plasma (1 ml) was diluted with 5 ml of distilled water. Trichloroacetic acid (10% w/v, 1.5 ml) was added to 1.5 ml of diluted plasma, after which the samples were mixed and centrifuged at 2500  $\times$  g for 10 min. The supernatant (2 ml) was placed into a glass test tube and 500  $\mu$ l of 0.06 M aqueous thiobarbituric acid was added. The samples were mixed and incubated at 80°C for 90 min in a water bath, after which the absorbance was measured at 532 nm. The TBARS assay in liver and muscle was performed following the same procedure described by Inserra *et al.* (2014).

The resistance of muscle lipids to peroxidation was also assessed by incubating muscle homogenates in the presence of the ferric chloride/sodium ascorbate (Fe/Asc) system as catalyst of lipid peroxidation, following the procedure described by Monahan *et al.* (2005) with modifications. The muscle (12.5 g) was homogenized with 37.5 g cold phosphate buffer (4°C, 50 mM, pH 7.0) and a volume (3 ml) was collected immediately. Then, ferric chloride hexahydrate and L-sodium ascorbate were added at equimolar concentration to the remaining muscle homogenate (45  $\mu$ M final concentration), which was incubated in a temperature-controlled IKA KS-4000 orbital shaker (IKA-Werke GmbH & Co. KG, Staufen, Germany) set at 37°C and 190 r.p.m. The homogenate (3 ml) was collected after 30 and 60 min of

incubation. Each collected aliquot of homogenate was mixed with trichloroacetic acid (3 ml, 10%) and filtered. The clear filtrate was used for the TBARS assay following the same procedure as for liver and muscle. In all the TBARS assays described above, standard solutions of 1,1,3,3-tetraethoxypropane were used to express the results as  $\mu\text{g}$  of malonaldehyde (MDA)/g of faeces, intestine, liver and muscle, or as  $\mu\text{g}$  MDA/ml of plasma.

#### Statistical analyses

The data related to the antioxidant status parameters and lipid oxidation measured in faeces, intestine, liver, plasma and muscle were analysed using a GLM model to test the effect of the dietary treatment (diet: CON or CIT). The data related to the development of lipid oxidation in muscle homogenates incubated in the presence of Fe/Asc were analysed using a GLM mixed model. The fixed factors tested in the model were the diet, the incubation time (time: 0, 30 and 60 min) and their interaction (diet  $\times$  time), whereas the individual animal was considered a random factor nested with the diet. *Post hoc* comparisons of the least squares means were performed using the Tukey's adjustment for multiple comparisons. In all the statistical analyses, the individual lambs were considered as the experimental units and significance was declared for  $P \leq 0.05$ . The statistical analyses were performed using the software Minitab, version 16 (Minitab Inc., State College, PA, USA).

## Results and discussion

#### Antioxidant status parameters of liver, plasma and muscle

The concentration of phenolic compounds in the DCP used in the present study was 10.87 g GAE eq./kg DM, which is similar to the content of GAE eq./kg reported for whole lemon powder (García-Salas *et al.*, 2013). Therefore, it was expected that including 35% DCP in a diet for lambs to partially replace barley could increase the concentration of phenolic compounds in the diet, which was confirmed by the twofold greater concentration of phenolic compounds found in the CIT diet compared with the CON diet (Table 1). Citrus phenolic compounds possess antioxidant activities and their intake with the diet has been suggested to increase the antioxidant capacity of animal tissues. In the light of this, we recently demonstrated that the oxidative stability of the meat from the same lambs used in this study was increased by feeding the CIT diet and we attributed this effect to the presence of flavonoids in the DCP (Inserra *et al.*, 2014). However, it has been reported that the possible mechanisms explaining the antioxidant effects of dietary citrus flavonoids are far from being clear and several authors suggest that their antioxidant effects *in vivo* are likely to be overestimated (Khan *et al.*, 2014). Regarding the bioavailability of phenolic compounds, it was demonstrated that the muscle from sheep fed rosemary contained several phenolic compounds originally present in the diet and this was associated to a higher overall antioxidant capacity of muscle, measured by FRAP and TEAC assays (Moñino *et al.*, 2008). With respect to citrus

flavonoids, Bodas *et al.* (2012) demonstrated that naringenin could be detected in the liver from lambs receiving a diet supplemented with 1.5 g/kg naringin. Similarly, Gladine *et al.* (2007c) showed that the administration of citrus extract to sheep at 10% of the diet DM resulted in the appearance of naringenin in blood plasma, but had no effect on the radical scavenging activity of plasma, measured with the TEAC assay. The same authors reported that supplementing the diet of rats with either 5 or 0.5 g/kg of citrus extract did not affect the radical scavenging activity of blood plasma, liver and muscle (Gladine *et al.*, 2007a and 2007b). These results are in agreement with those found in the present study in which the assessment of the overall antioxidant capacity of the hydrophilic fraction of blood plasma, liver and muscle showed that feeding CP did not affect the radical scavenging activity and the reducing capacity of tissues, measured with the TEAC, FRAP and Folin–Ciocalteu assays ( $P > 0.05$ ; Table 2). The TEAC, FRAP and Folin–Ciocalteu assays used in the present experiment, and in the above cited studies, are not specific for phenolic compounds and they measure the overall contribution of antioxidant compounds bearing radical scavenging and reducing properties (Niki, 2010). In this sense, these assays could detect the presence of absorbed phenolic compounds and/or of their metabolites if these molecules are still able to exert antioxidant activities. Moreover, these tests could indicate if dietary phenolic

**Table 2** Effect of the dietary treatment on the antioxidant status of liver, plasma and muscle

	Group <sup>1</sup>		SEM	P-value
	CON	CIT		
<b>Liver</b>				
TEAC <sup>2</sup>	28.35	27.43	0.487	0.524
FRAP <sup>3</sup>	18.16	17.34	0.534	0.609
Folin–Ciocalteu <sup>4</sup>	2.62	2.68	0.045	0.303
$\alpha$ -tocopherol <sup>5</sup>	20.06	38.52	4.160	0.022
<b>Plasma</b>				
TEAC <sup>2</sup>	5.92	6.37	0.167	0.274
FRAP <sup>3</sup>	0.91	0.91	0.019	0.923
Folin–Ciocalteu <sup>4</sup>	0.60	0.55	0.016	0.112
$\alpha$ -tocopherol <sup>5</sup>	0.27	0.43	0.040	0.042
<b>Muscle</b>				
TEAC <sup>2</sup>	7.05	7.55	0.385	0.478
FRAP <sup>3</sup>	3.14	3.16	0.087	0.659
Folin–Ciocalteu <sup>4</sup>	0.46	0.45	0.011	0.726
$\alpha$ -tocopherol <sup>5</sup>	0.49	1.51	0.171	0.001

<sup>1</sup>Group CON was fed a control barley-based concentrate diet. Group CIT was fed a concentrate diet including 35% (as-fed) whole dried citrus pulp.

<sup>2</sup>Trolox equivalent antioxidant capacity (TEAC). Expressed as  $\mu\text{moles}$  of trolox equivalents/g for liver and muscle, or as  $\mu\text{moles}$  of trolox equivalents/ml for plasma.

<sup>3</sup>Ferric reducing antioxidant power (FRAP). Expressed as  $\mu\text{moles}$  of  $\text{Fe}^{+2}$  equivalents/g for liver and muscle, or as  $\mu\text{moles}$  of  $\text{Fe}^{+2}$  equivalents/ml for plasma.

<sup>4</sup>Expressed as mg of gallic acid equivalents (GAE)/g for liver and muscle, or as mg of GAE/ml for plasma.

<sup>5</sup>Expressed as  $\mu\text{g}$  of  $\alpha$ -tocopherol/g for liver and muscle, or as  $\mu\text{g}$  of  $\alpha$ -tocopherol/ml for plasma.

compounds are able to indirectly improve the overall antioxidant capacity of animal tissues through mechanisms that do not necessarily involve their detectable presence in the tissues. For example, recent studies demonstrated that feeding lambs with fresh herbage or with a plant extract rich in condensed tannins increased the overall antioxidant capacity of liver and plasma measured by the FRAP and the Folin–Ciocalteu assays (López-Andrés *et al.*, 2013 and 2014). However, the authors reported that no phenolic compounds, nor their metabolites, could be detected in the tissues and concluded that dietary phenolic compounds might be able to improve the antioxidant capacity *in vivo* through still unknown indirect antioxidant effects. The results obtained in the present study seem to suggest that dietary flavonoids might have not been effective in increasing the antioxidant capacity in the tissues tested. Certainly, it will be important in future studies to test the possible bioavailability of flavonoids when DCP is fed to animals instead of selected flavonoids or purified citrus extracts.

Among the exogenous antioxidants of dietary origin, vitamin E has been extensively proved to play a major role in protecting animal tissues against lipid peroxidation, with  $\alpha$ -tocopherol accounting alone for most of this activity (Burton and Traber, 1990). In the present experiment,  $\alpha$ -tocopherol was measured in the lipid fraction extracted from liver, blood plasma and muscle to assess the possible effect of the dietary treatment on the antioxidant status of the lipophilic fraction of the tissues. We found that feeding lambs with the diet containing 35% DCP (CIT) increased the concentration of  $\alpha$ -tocopherol in plasma ( $P < 0.05$ ) and resulted in a twofold and threefold greater concentration of  $\alpha$ -tocopherol in liver and in muscle, respectively ( $P < 0.05$  and  $P < 0.01$  for liver and muscle, respectively; Table 2). Mourão *et al.* (2008) found that the inclusion of 5% or 10% DCP in the diets of broilers did not affect the concentration of  $\alpha$ -tocopherol in muscle compared with a conventional cereal-based diet. To our knowledge, no information has been previously reported on the effect of feeding CP to ruminants on the vitamin E concentration in the tissues. In the present study, the experimental diets were not supplemented with vitamin E ( $\alpha$ -tocopheryl acetate or  $\alpha$ -tocopherol) in order to better appreciate the contribution of DCP to the vitamin E content of the formulated diet. The higher concentration of  $\alpha$ -tocopherol found in liver, plasma and muscle from lambs fed the CIT diet were at least partially due to the greater concentration of  $\alpha$ -tocopherol in the CIT diet compared with the CON diet, consequent to the partial replacement of barley with DCP (Table 1). Indeed, the  $\alpha$ -tocopherol content of the DCP included in the CIT diet was 80.7 mg/kg, whereas barley contained 12.2 mg/kg. Very scarce information is available on the content of fat-soluble compounds, such as vitamin E, in CP. However, it should be considered that seeds are removed during the production of citrus juice and, together with peel and pulp, are a component of DCP. It has been reported that citrus seeds can contain remarkable concentrations of essential fatty acids, such as linoleic and  $\alpha$ -linolenic acids, whereas noticeable concentrations of

$\alpha$ -tocopherol can be found in the unsaponifiable matter of the seed oil (Fontanel, 2013). For example, we recently demonstrated that replacing barley with 24% and 35% whole DCP in concentrate-based diets for lambs resulted in a higher intake of fat and fatty acids (Lanza *et al.*, 2015), which demonstrates that DCP may represent a dietary source of fat-soluble compounds.

#### Resistance to lipid oxidation

Studies on the antioxidant effects of dietary phenolic compounds suggest a possible direct antioxidant activity exerted in the gastrointestinal tract, such as the reduction of lipid peroxidation, which would result in an overall improvement of the animal antioxidant status (Kerem *et al.*, 2006). Moreover, antioxidant effects in the gastrointestinal tract might be obtained by increasing the intake of vitamin E (Halliwell *et al.*, 2005). In the present study, the inclusion of 35% DCP in the diet increased the concentrations of both phenolic compounds and  $\alpha$ -tocopherol of the diet and, therefore, the amounts of these compounds reaching the gastrointestinal tract of the lambs compared with the CON diet. We have measured the extent of lipid oxidation in the small intestine and in the faeces to assess possible antioxidant effects of DCP exerted in the gastrointestinal tract. The results obtained showed that lipid oxidation (TBARS values) in the small intestine and in the faeces was not affected by the dietary treatment ( $P > 0.05$ ; Table 3). These results can be partially explained considering that the diets supplemented to the lambs were not expected to promote lipid peroxidation, as we did not supplement the diets with readily oxidizable fatty acids or with pro-oxidant compounds. Indeed, most of the studies demonstrating effects of the diet on the lipid oxidation in the intestine or in the faeces have been conducted using monogastric animals, such as rats, which were generally subjected to factors inducing oxidative stress including, for example, feeding diets rich in haem iron or in fat and oxidized fatty acids, or diets simulating antioxidant deficiency (Lindley *et al.*, 1994; Pierre *et al.*, 2003).

In the present study, it was hypothesized that possible effects of citrus antioxidants in the gastrointestinal tract,

**Table 3** Effect of the dietary treatment on lipid oxidation (TBARS values) measured in small intestine, faeces, liver, plasma and muscle

	Group <sup>1</sup>		SEM	P-value
	CON	CIT		
TBARS values <sup>2</sup>				
Faeces	1.25	1.23	0.161	0.955
Small intestine	1.23	0.87	0.136	0.197
Liver	1.95	1.74	0.132	0.437
Plasma	0.19	0.17	0.011	0.390
Muscle	0.19	0.14	0.050	0.634

<sup>1</sup>Group CON was fed a control barley-based concentrate diet. Group CIT was fed a concentrate diet including 35% (as-fed) whole dried citrus pulp.

<sup>2</sup>Expressed as  $\mu$ g of malonaldehyde (MDA)/g for small intestine, faeces, liver and muscle, or as  $\mu$ g MDA/ml for plasma.


together with their absorption and delivery to the liver, plasma and muscle could have increased the resistance of these tissues to lipid oxidation. We found no effect of the dietary treatment on the extent of lipid oxidation (TBARS values) measured in liver, plasma and muscle ( $P > 0.05$ ; Table 3). In the case of citrus phenolic compounds, it has been demonstrated that adding a citrus extract to diets enriched in highly oxidizable fatty acids reduced the extent of lipid peroxidation in the liver and plasma of rats (Gladine *et al.*, 2007a and 2007b). The same authors included 10% of citrus extract in the diet of sheep which were made highly susceptible to lipid oxidation already by continuous infusion of linseed oil into the duodenum (Gladine *et al.*, 2007c). This experimental design allowed to generate an oxidative challenge already *in vivo* and the authors reported that citrus extract increased the resistance of plasma lipids to oxidation. Once again the fact that, in the present study, animals were not subjected to conditions promoting lipid peroxidation could partially explain why an effect of feeding CP on lipid oxidation was not observed in liver, plasma and muscle collected from the animals and immediately stored at  $-80^{\circ}\text{C}$  without being subjected to further oxidative stresses until analyses. Nevertheless, using the same animals of the present experiment, Inserra *et al.* (2014) demonstrated that the CIT diet reduced the susceptibility to lipid oxidation of muscle, which could be appreciated when meat was stored for 6 days at  $4^{\circ}\text{C}$ . Therefore, in the present study, we also assessed the resistance of muscle lipids to oxidation by incubating muscle homogenates in the presence of Fe/Asc as catalyst for inducing lipid peroxidation (Monahan *et al.*, 2005). We found that TBARS values dramatically increased with time of incubation in the presence of the catalyst ( $P < 0.001$ ; Figure 1). Moreover, the dietary treatment affected the TBARS values measured over the incubation time and a significant diet  $\times$  time interaction was found ( $P < 0.01$ ). Indeed, as shown in Figure 1, although the TBARS values did not differ between the CON and CIT treatments at

the beginning of the incubation, the muscle from lambs fed with the CIT diet exhibited lower TBARS values compared with the CON treatment after 30 and 60 min ( $P < 0.01$ ). These results confirmed those reported by Inserra *et al.* (2014) and demonstrate that the inclusion of CP in a concentrate-based diet for lambs is able to improve the resistance of muscle to lipid oxidation under pro-oxidant conditions. In the light of the results found for the antioxidant status parameters measured in muscle, it can be stated that, in the present study, the positive effect of DCP on the resistance of muscle lipids to oxidation could be explained by the higher deposition of  $\alpha$ -tocopherol in muscle from the animals fed the CIT diet compared with the CON treatment. Indeed, it has been widely demonstrated that the concentration of vitamin E in muscle plays a major role in protecting lipids from oxidation and some authors were able to develop models to predict the development of lipid oxidation in meat based on the levels of vitamin E in muscle (Sales and Koukolová, 2011). In the present study, we found no correlation between the  $\alpha$ -tocopherol content in the muscle and the TBARS value at the start of the incubation in the presence of Fe/Asc catalyst, whereas a negative correlation was evident with the TBARS values measured after 30 min ( $-0.61$ ;  $P < 0.01$ ) and 60 min ( $-0.55$ ;  $P = 0.01$ ).

## Conclusions

Despite the fact that flavonoids are generally considered the main antioxidants in citrus fruits, the results of this study lead to the conclusion that  $\alpha$ -tocopherol plays the major role in the improvement of the antioxidant capacity of muscle from lambs fed diets supplemented with DCP. The greater concentration of  $\alpha$ -tocopherol found in the liver, plasma and muscle from lambs fed CP compared with the control treatment could be explained in the light of the higher concentration of  $\alpha$ -tocopherol in the CIT diet. This result was rather unexpected, as there is no information on the effects of feeding CP to ruminants on the concentration of  $\alpha$ -tocopherol in the animal tissues.

Certainly, further research should clarify the antioxidant effect of dietary CP in ruminants. Indeed, the chemical composition of CP can greatly vary depending on several factors. For example, the whole CP can undergo further processing before drying to obtain essential oils and seed oil, thus reducing the content of potential antioxidants in the final product destined to animals. Therefore, it would be necessary to study different available CP in order to standardize its application in livestock feeding and to optimize the positive effects on meat quality.



**Figure 1** Interactive effect of the dietary treatment (CON and CIT) and of the time of incubation (0, 30 and 60 min) on the TBARS values in muscle homogenates incubated with the ferric chloride/sodium ascorbate system. Values presented are the least squares means with standard error bars. Diets were as follows: control barley-based concentrate diet (CON); concentrate diet including 35%, as-fed, dried citrus pulp (CIT). <sup>a,b,c,d</sup>Indicate differences between mean values ( $P < 0.05$ ). MDA = malonaldehyde.

## Acknowledgements

The authors gratefully acknowledge funding from the European Community financial participation under the Seventh Framework Programme for Research, Technological Development and Demonstration Activities, for the Integrated Project LOW-INPUTBREEDS FP7-CP-IP 222623. The views expressed in this

publication are the sole responsibility of the authors and do not necessarily reflect the views of the European Commission. Neither the European Commission nor any person acting on behalf of the Commission is responsible for the use which might be made of the information contained herein.

## References

- Association of Official Analytical Chemists 1990. Official method of analysis, vol. 2, 15th edition. AOAC, Arlington, VA, USA.
- Balasundram N, Sundram K and Samman S 2006. Phenolic compounds in plants and agri-industrial by-products: antioxidant activity, occurrence and potential uses. *Food Chemistry* 99, 191–203.
- Bampidis VA and Robinson PH 2006. Citrus by-products as ruminant feeds: a review. *Animal Feed Science and Technology* 128, 175–217.
- Bodas R, Prieto N, Jordán MJ, López-Campos O, Giráldez FJ, Morán L and Andrés S 2012. The liver antioxidant status of fattening lambs is improved by naringin dietary supplementation at 0.15% rates but not meat quality. *Animal* 6, 863–870.
- Burton GW and Traber MG 1990. Vitamin E: antioxidant activity, biochemistry and bioavailability. *Annual Review of Nutrition* 10, 357–382.
- Folch J, Lees M and Stanley GHS 1957. A simple method for the isolation and purification of lipids from animal tissue. *Journal of Biological Chemistry* 226, 497–509.
- Fontanel D 2013. Unsaponifiable matter in plant seed oils, 1st edition. Springer-Verlag, Berlin and Heidelberg, Germany.
- García-Salas P, Gómez-Caravaca AM, Arráez-Román D, Segura-Carretero A, Guerra-Hernández E, García-Villanova B and Fernández-Gutiérrez A 2013. Influence of technological processes on phenolic compounds, organic acids, furanic derivatives, and antioxidant activity of whole-lemon powder. *Food Chemistry* 141, 869–878.
- Gladine C, Morand C, Rock E, Bauchart D and Durand D 2007a. Plant extracts rich in polyphenols (PREP) are efficient antioxidants to prevent lipoperoxidation in plasma lipids from animals fed *n*-3 PUFA supplemented diets. *Animal Feed Science and Technology* 136, 281–296.
- Gladine C, Morand C, Rock E, Guffrat D, Bauchart D and Durand D 2007b. The antioxidative effect of plant extracts rich in polyphenols differs between liver and muscle tissues in rats fed *n*-3 PUFA rich diets. *Animal Feed Science and Technology* 139, 257–272.
- Gladine C, Rock E, Morand C, Bauchart D and Durand D 2007c. Bioavailability and antioxidant capacity of plant extracts rich in polyphenols, given as a single acute dose, in sheep made highly susceptible to lipoperoxidation. *British Journal of Nutrition* 98, 691–701.
- Gravador RS, Jongberg S, Andersen ML, Luciano G, Priolo A and Lund MN 2014. Dietary citrus pulp improves protein stability in lamb meat stored under aerobic conditions. *Meat Science* 97, 231–236.
- Halliwell B, Rafter J and Jenner A 2005. Health promotion by flavonoids, tocopherols, tocotrienols and other phenols: direct or indirect effects? Antioxidants or not? *American Journal of Clinical Nutrition* 81, 268–276.
- Inserra L, Priolo A, Biondi L, Lanza M, Bognanno M, Gravador R and Luciano G 2014. Dietary citrus pulp reduces lipid oxidation in lamb meat. *Meat Science* 96, 1489–1493.
- Kerem Z, Chertrit D, Shoseyov O and Regev-Shoshani G 2006. Protection of lipids from oxidation by epicatechin, trans-resveratrol, and gallic and caffeic acids in intestinal model systems. *Journal of Agricultural and Food Chemistry* 54, 10288–10293.
- Khan MK, Huma ZE and Dangles O 2014. A comprehensive review on flavanones, the major citrus polyphenols. *Journal of Food Composition and Analysis* 33, 85–104.
- Lanza M, Scerra M, Bognanno M, Bognanno M, Buccioni A, Cilione C, Biondi L, Priolo A and Luciano G 2015. Fatty acid metabolism in lambs fed citrus pulp. *Journal of Animal Science* 93, 3179–3188.
- Lindley KJ, Goss-Sampson MA, Muller DPR and Milla PJ 1994. Lipid peroxidation and electrogenic ion transport in the jejunum of the vitamin E deficient rat. *Gut* 35, 34–39.
- López-Andrés P, Luciano G, Vasta V, Gibson TM, Biondi L, Priolo A and Mueller-Harvey I 2013. Dietary quebracho tannins are not absorbed, but increase the antioxidant capacity of liver and plasma in sheep. *British Journal of Nutrition* 110, 632–639.
- López-Andrés P, Luciano G, Vasta V, Gibson TM, Scerra M, Biondi L, Priolo A and Mueller-Harvey I 2014. Antioxidant effect of ryegrass phenolics in lamb liver and plasma. *Animal* 8, 51–57.
- Luciano G, Vasta V, Monahan FJ, López-Andrés P, Biondi L, Lanza M and Priolo A 2011. Antioxidant status, colour stability and myoglobin resistance to oxidation of *longissimus dorsi* muscle from lambs fed a tannin-containing diet. *Food Chemistry* 124, 1036–1042.
- Makkar HPS, Blümmel M, Borowy NK and Becker K 1993. Gravimetric determination of tannins and their correlations with chemical and protein precipitation methods. *Journal of the Science of Food and Agriculture* 61, 161–165.
- Monahan FJ, Skibsted LH and Andersen ML 2005. Mechanism of oxymyoglobin oxidation in the presence of oxidizing lipids in bovine muscle. *Journal of Agricultural and Food Chemistry* 53, 5734–5738.
- Moñino I, Martínez C, Sotomayor JA, Lafuente A and Jordán MJ 2008. Polyphenolic transmission to Segureño lamb meat from ewes' diet supplemented with the distillate from rosemary (*Rosmarinus officinalis*) leaves. *Journal of Agricultural and Food Chemistry* 56, 3363–3367.
- Mourão JL, Pinheiro VM, Prates JAM, Bessa RJB, Ferreira LMA, Fontes CMGA and Ponte PIP 2008. Effect of dietary dehydrated pasture and citrus pulp on the performance and meat quality of broiler chickens. *Poultry Science* 87, 733–743.
- Niki E 2010. Assessment of antioxidant capacity *in vitro* and *in vivo*. *Free Radical Biology and Medicine* 49, 503–515.
- Nogata Y, Sakamoto K, Shiratsuchi H, Ishii T, Iano M and Ohto H 2006. Flavonoid composition of fruit tissues of citrus species. *Bioscience Biotechnology and Biochemistry* 70, 178–192.
- Pierre F, Taché S, Petit CR, Van der Meer R and Corpet DE 2003. Meat and cancer: haemoglobin and haemin in a low-calcium diet promote colorectal carcinogenesis at the aberrant crypt stage in rats. *Carcinogenesis* 24, 1683–1690.
- Sales J and Koukolová V 2011. Dietary vitamin E and lipid and colour stability of beef and pork: modeling of relationships. *Journal of Animal Science* 89, 2836–2848.
- Schüep W and Rettenmaier R 1994. Analysis of vitamin E homologous in plasma and tissue: high-performance liquid chromatography. *Methods in Enzymology* 234, 294–302.
- Siu GM and Draper HH 1978. A survey of the malonaldehyde content of retail meats and fish. *Journal of Food Science* 43, 1147–1149.
- Van Soest PJ, Robertson JB and Lewis BA 1991. Methods for dietary fibre, neutral detergent fibre, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* 74, 3583–3597.
- Zaspel JB and Csallany S 1983. Determination of alpha-tocopherol in tissue and plasma by high-performance liquid chromatography. *Analytical Biochemistry* 30, 146–150.