

Rapid decline of Philadelphia-positive metaphases after nilotinib treatment in a CML patient expressing a rare e14a3 *BCR-ABL1* fusion transcript: A case report

MICHELE MASSIMINO^{1,2*}, STEFANIA STELLA^{1,2*}, ELENA TIRRO^{1,2*}, MARIA LETIZIA CONSOLI³, MARIA STELLA PENNISI^{1,2}, ADRIANA PUMA^{1,2}, SILVIA RITA VITALE^{1,2}, CHIARA ROMANO^{1,2}, VALENTINA ZAMMIT³, FABIO STAGNO³, FRANCESCO DI RAIMONDO^{3,4} and LIVIA MANZELLA^{1,2}

¹Department of Clinical and Experimental Medicine, University of Catania;

²Center of Experimental Oncology and Hematology; ³Division of Hematology and Bone Marrow Transplant,

A.O.U. Policlinico-Vittorio Emanuele; ⁴Department of Surgery, Medical and Surgical Specialties, University of Catania, Catania I-95123, Italy

Received January 29, 2019; Accepted May 15, 2019

DOI: 10.3892/ol.2019.10558

Abstract. We report a case of chronic myeloid leukemia in a 52-year-old male expressing a rare e14a3 *BCR-ABL1* fusion transcript. Cytogenetic analysis showed the t(9;22) translocation and multiplex RT-PCR detected an atypical fragment of approximately 230 base pairs. Using two primers recognizing exon 10 of *BCR* and exon 4 of *ABL1*, a larger PCR product was identified, cloned, sequenced and defined as an e14a3 *BCR-ABL1* rearrangement. The patient was treated with nilotinib and monitored measuring cytogenetic and hematological parameters, while *BCR-ABL1* transcripts were surveyed by conventional and semi-nested PCR. The patient achieved a complete hematologic response after two months of treatment followed by a complete cytogenetic remission two months later. Furthermore, PCR and semi-nested PCR failed to detect the e14a3 *BCR-ABL1* mRNA after 15 and 21 months of nilotinib, respectively.

Introduction

The BCR-ABL oncoprotein is the culprit of chronic myeloid leukemia (CML) as it transforms the hematopoietic stem cell altering its survival, proliferation and interaction with both the cell cytoskeleton and the bone marrow microenvironment (1-6). In the last 15 years, multiple tyrosine kinase

inhibitors (TKIs) have been approved for the first line treatment of the disease including imatinib (IM), dasatinib (DAS), nilotinib (NIL) and bosutinib (BOS) (7-12). From a genetic standpoint, the *BCR-ABL1* chimeric oncogene derives from the t(9;22) reciprocal translocation that generates the Philadelphia (Ph) chromosome (13). Most patients express one of three fusion transcripts juxtaposing *BCR* exons 1, 13 or 14 with exon 2 of *ABL1* (14). However, several alternative breakpoints involving different *BCR* and/or *ABL1* exons have been previously described. Specifically, *BCR* exons 1, 6, 8, 13, 14 and 19 can rearrange with *ABL1* exons 2 or 3, generating e6a2, e8a2, e1a3, e13a3, e14a3, e19a3 fusions (15).

BCR-ABL1 fusions involving *ABL1* exon 3 are extremely rare (0.3%) and are usually associated with contrasting clinical outcome (16,17). In the present study we report a CML patient expressing an atypical e14a3 *BCR-ABL1* transcript that was successfully treated with NIL.

Case Report

In October 2016 a 52 year-old male (Table I) with a history of neutrophilia in the absence of thrombocytosis was admitted to the Hematology ward and subjected to a bone marrow aspirate in order to perform a karyotype analysis by G-banding. His cytogenetic profile showed the t(9;22) translocation in 100% of the analyzed metaphases (20/20) (Fig. 1A). At this time, his white blood cells (WBCs) were lysed and used to extract total messenger RNA (mRNA) that was reverse transcribed in cDNA and employed to perform a multiplex RT-PCR in order to determine his *BCR-ABL1* fusion transcript variant (18). An atypical band of approximately 230 base pairs was detected (Fig. 1B) with no amplification of the common *BCR-ABL1* variants in real-time PCR, performed as previously described (19,20). However, the detection of the Ph chromosome by G-banding led to exclude that this 230 base pair band was a non-specific RT-PCR product. Hence, we employed the cDNA for a new PCR reaction using forward (BCR-10 5'-TATGAC

Correspondence to: Dr Michele Massimino, Department of Clinical and Experimental Medicine, University of Catania, 78 Via Santa Sofia, Catania I-95123, Italy
E-mail: michedot@yahoo.it

*Contributed equally

Key words: BCR-ABL1, chronic myeloid leukemia, nilotinib, e14a3, case report

TGCAAATGGTACATTCC-3') and reverse (ABL-4 5'-TCG TAGTTGGGGGACACACC-3') primers recognizing exons 10 and 4 of *BCR* and *ABL1*, respectively. The 977 bp PCR product (Fig. 1C) generated by the Pfx platinum DNA polymerase enzyme was then cloned in the pcr4-TOPO-TA vector according to the manufacture's protocol (all from Thermofisher Scientific). Plasmid DNA obtained from ten individual bacterial colonies was then subjected to Sanger sequencing and each colony showed the e14a3 fusion transcript. One representative pherogram displaying the *BCRe14* and *ABL1a3* gene exons rearrangement is depicted in Fig. 1D. Finally, Sokal (0.79, low), Hasford (949, intermediate), EUTOS (11, low) and ELTS (1.39, low) scores were calculated as indicated in Table I. In conclusion, the patient was diagnosed with chronic phase CML expressing a rare e14a3 *BCR-ABL1* variant. Although IM represents an excellent first line treatment for CML, previous data show that not all patients expressing the e14a3 isoform benefit from this drug (16,21). Thus, the patient, with no additional medication, received NIL 300 mg *bis in die* (BID), as this compound is a more potent inhibitor of ABL1 catalytic activity.

Disease evolution was monitored measuring both hematological and cytogenetic parameters. After two months of NIL, the patient achieved a complete hematological response (CHR) and, in February 2017, a new cytogenetic analysis failed to detect Ph+ metaphases suggesting a complete cytogenetic response (CCyR). During the following months, the patient maintained both his CHR and CCRy (Fig. 2A and B).

To molecularly monitor the disease, we evaluated the presence of the e14a3 *BCR-ABL1* gene by PCR and the negative reaction was confirmed by semi nested-PCR (SN-PCR) at the time points indicated in Fig. 2A. A first round of PCR was performed using the primers specified above while for the SN-PCR we employed a Fw primer recognizing exon 12 of the *BCR* gene (BCR-12 5'-GTGCAGAGTGGAGGGAGA ACA-3') obtaining a PCR product of approximately 668 bp. In December 2016 the PCR detected the e14a3 fusion transcript. However, in July 2017, *BCR-ABL1* mRNA-reverse transcribed by Superscript III One-Step RT-PCR (Thermofisher Scientific)-was detected only after SN-PCR and by December 2017 and June 2018 neither reaction detected the e14a3 fusion transcript suggesting a strong reduction in the overall number of *BCR-ABL1*-positive cells (Fig. 2B).

In order to evaluate the depth of the response to NIL, in January 2019 we isolated CD34+ cells from peripheral blood performing a colony forming units (CFU) assay as previously described (22). Ten CD34+-derived colonies were collected and total RNA extracted using the Trizol reagent (Life Technologies). We subsequently employed the Superscript III One-Step RT-PCR (Life Technologies) to detect the e14a3 *BCR-ABL1* fusion. Negative colonies for the first *BCR-ABL1* amplification were subjected to a semi-nested PCR as described above. Interestingly, we failed to detect the fusion transcript in any of the analyzed colonies (data not show).

At the last control (January 2019), the patient is still receiving NIL with no clinical, hematological, cytogenetic or molecular signs of disease progression. Furthermore, he has not reported side effects to NIL treatment during the course of his regular outpatient visits.

Table I. Patient characteristics at the time of diagnosis.

Patient characteristic	Values
Complete blood count	
Platelets (μ l)	207.000
WBCs (μ l)	53910x10 ³
Neutrophils	64%
Eosinophils	3%
Basophils	1%
Lymphocytes	10%
Monocytes	1%
Metamyelocytes	5%
Myelocytes	10%
Promyelocytes	4%
Myeloblasts	2%
Haemoglobin (g/dl)	12.5
Cytogenetic analysis	
Karyotype	46, XY,100% (9;22)(q34;q11)
Fusion transcript	
<i>BCR-ABL1</i>	e14a3
Relative risk	
Sokal	0.79 (Low)
Hasford	949 (Intermediate)
EUTOS	11 (Low)
ELTS	1.39 (Low)

Discussion

BCR-ABL1 transcripts displaying breakpoints lacking *ABL1* exon 2 comprise e1a3, e13a3 and e14a3 rearrangements and represent infrequently occurring oncogenic isoforms (21). A methodological issue may have contributed to the uncommon detection of these transcripts as multiplex RT-PCR reactions can generate atypical PCR fragments often interpreted as nonspecific products. In our study, the detection of the t(9;22) translocation by G-banding indicated the presence of a Ph chromosome. We therefore decided to employ primers recognizing more distant exons from the common *BCR-ABL1* breakpoints and successfully identified the atypical *BCR-ABL1* e14a3 rearrangement. To date, different cases of CML patients expressing this isoform have been reported (16,21,23,24). Although IM represents an excellent first line therapy for most CML patients (7), extensive published data suggest that patients receiving this drug may more frequently develop both *BCR-ABL*-dependent and *BCR-ABL*-independent resistance to therapy (8,19,25) requiring alternative treatments (26). Furthermore, an inverse correlation has been reported between the size of the *BCR* portion retained in the oncogenic fusion protein and CML aggressiveness (27). Indeed, complex variant translocations, intron-derived insertions/truncations in the *BCR-ABL1* kinase domain or hyperdiploidy suggest that CML patients displaying these genetic alterations often present an unfavorable clinical outcome and inferior IM responses (13,16,21,28,29).




Figure 1. e14a3 *BCR-ABL1* fusion transcript identification. (A) The image shows the G-banded karyotype. Arrows indicate the t(9;22) translocation. (B) Multiplex RT-PCR of different *BCR-ABL1* fusion transcripts performed on total RNA extracted from immortalized cell lines SD1 (*BCR-ABL* p190-e1a2), K562 (*BCR-ABL* p210-e14a2) and BV-173 (p210-e13a2). Ctrl- indicates RNA derived from healthy donor and Pt-e14a3 indicates the atypical e14a3 fragment of 230 bps. The 808 bp BCR band represents a PCR internal reaction control amplified when the sample is negative for *BCR-ABL1* expression. Numbers indicate the size of the bands obtained by multiplex PCR (18). (C) The panel shows bands generated with a PCR reaction using BCR-10 and ABL-4 primers. Ctrl-(reaction mix missing cDNA) and K562 were used as negative and positive controls, respectively. (D) Schematic representation of the e14a3 *BCR-ABL1* fusion transcript and one representative pherogram obtained after Sanger sequencing of each bacterial colony showing the *BCRe14* and *ABLa3* exons junction.

The *ABL1* a2 exon encodes for a Src homology domain 3 (SH3) and, although its loss may lead to reduced leukemogenesis with a benign clinical course, its role as a negative regulator of the ABL kinase domain (SH1) may explain the reportedly more aggressive CML phenotype (30,31). Moreover, while the e14a3 *BCR-ABL1* breakpoint preserves the ATP binding

domain, the lack of the SH3 domain may modify the SH1 domain tertiary structure thus affecting drug response (32).

On the base of these findings and the controversial data on IM efficacy in subjects displaying the e14a3 transcript (16), we wanted to employ a second-generation TKI (2G-TKI) as first line treatment. Nilotinib was selected for his treatment




Figure 2. Hematologic, cytogenetic and molecular monitoring of the patient. (A) The graph depicts white blood cells (WBC) counts from the time of diagnosis (October 2016) to the last follow-up (January 2019) and indicates the time points of the cytogenetic and molecular analyses. (B) Follow-up of G-banded karyotype showing achievement and maintenance of CCRy at the indicated time points. Arrows indicate the normal chromosomes 9 and 22. (C) PCR and SN-PCR for e14a3 detection performed at the indicated time points. Ctrl- indicates the reaction mix missing cDNA. Nuclear receptor coactivator 4 (NCOA-4) was used as loading control and amplified by the Fw 5'-TGGAGAAGAGAGGCTGTATCT-3' and Rv 5'-ATTGAAGAAATGCAGGCTC-3' primers. WBC, White Blood Cells; NIL, nilotinib; Ph+, Philadelphia-positive metaphases; CHR, Complete Hematological Response; CCyR, Complete Cytogenetic Response; CFU, Colony Forming Units; SN, Semi Nested; BA, BCR-ABL.

because DAS was temporarily unavailable in our pharmacy at the time. Since the patient is an accountant who works from home, he has shown excellent compliance with the administration of the drug 1 h before meals.

Using NIL, a more potent ABL1 inhibitor (9), we observed a rapid decline in Ph metaphases that generated a CCyR after 4 months of NIL. Finally, employing a semi-nested RT-PCR we failed to detect *BCR-ABL1* transcripts both in peripheral WBCs and in CFUs grown in methylcellulose, indicating that the drug induced a rapid decline in the overall number of leukemic cells.

In summary, CML patients expressing the e14a3 *BCR-ABL1* fusion transcript may be undiagnosed because this rearrangement generates an atypical PCR product often misinterpreted as a nonspecific band that is then coupled to a negative real-time PCR result. Hence, performing a

cytogenetic analysis is critical to identify these CML patients. Moreover, the difficulty to perform a precise quantitative molecular monitoring and the rare incidence of atypical *BCR-ABL1* fusion transcripts does not allow the design of randomized clinical trials that may compare the efficacy of IM vs. 2G-TKIs.

We conclude that the *BCR-ABL1* oncoprotein, derived from *BCRe14* and *ABL1a3* exons rearrangement, is highly sensitive to NIL, suggesting that chronic phase CML patients exhibiting this rare rearrangement may quickly achieve CCyR followed by strong reductions in the size of the leukemic clone.

Acknowledgements

Not applicable.

Funding

Piano Sanitario Nazionale 2015 (grant no. Linea Progettuale 6-Azione 6.3).

Availability of data and materials

All data generated or analyzed during the present study are included in this published article.

Authors' contributions

MM drafted the manuscript; MM, ET and SS were responsible for study conception, and designed and performed the experiments; MM, ET, SS, MSP, AP, SRV and CR analyzed and interpreted the data; MLC performed cytogenetic analysis; FS and VZ made a critical revision of the paper and managed the patient; FDR supervised the project and contributed to study design; LM conceived the original idea and supervised the project. All authors approved the final version of the manuscript to be published.

Ethics approval and consent to participate

The patient provided written informed consent to participate. The study adheres to the declaration of Helsinki and the biological samples were collected following an institutionally approved protocol at the Azienda Ospedaliero-Universitaria 'Policlinico-Vittorio Emanuele' (Catania, Italy).

Patient consent for publication

Written informed consent obtained from patient for publication of this report.

Competing interest

The authors declare that they have no competing interests.

References

- Ren R: Mechanisms of BCR-ABL in the pathogenesis of chronic myelogenous leukaemia. *Nat Rev Cancer* 5: 172-183, 2005.
- Stella S, TIRRÒ E, Conte E, Stagno F, Di Raimondo F, Manzella L and Vigneri P: Suppression of survivin induced by a BCR-ABL/JAK2/STAT3 pathway sensitizes imatinib-resistant CML cells to different cytotoxic drugs. *Mol Cancer Ther* 12: 1085-1098, 2013.
- Ishii Y, Nhaiy MK, Tse E, Cheng J, Massimino M, Durden DL, Vigneri P and Wang JY: Knockout serum replacement promotes cell survival by preventing BIM from inducing mitochondrial cytochrome C release. *PLoS One* 10: e0140585, 2015.
- Manzella L, TIRRÒ E, Pennisi MS, Massimino M, Stella S, Romano C, Vitale SR and Vigneri P: Roles of interferon regulatory factors in chronic myeloid leukemia. *Curr Cancer Drug Targets* 16: 594-605, 2016.
- Preyer M, Vigneri P and Wang JY: Interplay between kinase domain autophosphorylation and F-actin binding domain in regulating imatinib sensitivity and nuclear import of BCR-ABL. *PLoS One* 6: e17020, 2011.
- Giallongo C, Tibullo D, La Cava P, Branca A, Parrinello N, Spina P, Stagno F, Conticello C, Chiarenza A, Vigneri P, *et al*: BRIT1/MCPH1 expression in chronic myeloid leukemia and its regulation of the G2/M checkpoint. *Acta Haematol* 126: 205-210, 2011.
- Stagno F, Stella S, Spitaleri A, Pennisi MS, Di Raimondo F and Vigneri P: Imatinib mesylate in chronic myeloid leukemia: Frontline treatment and long-term outcomes. *Expert Rev Anticancer Ther* 16: 273-278, 2016.
- Hochhaus A, Larson RA, Guilhot F, Radich JP, Branford S, Hughes TP, Baccarani M, Deininger MW, Cervantes F, Fujihara S, *et al*: Long-term outcomes of imatinib treatment for chronic myeloid leukemia. *N Engl J Med* 376: 917-927, 2017.
- Hochhaus A, Rosti G, Cross NC, Steegmann JL, le Coutre P, Ossenkoppele G, Petrov L, Masszi T, Hellmann A, Griskevicius L, *et al*: Frontline nilotinib in patients with chronic myeloid leukemia in chronic phase: Results from the European ENEST1st study. *Leukemia* 30: 57-64, 2016.
- Cortes JE, Saglio G, Kantarjian HM, Baccarani M, Mayer J, Boqué C, Shah NP, Chuah C, Casanova L, Bradley-Garelik B, *et al*: Final 5-year study results of DASISION: The dasatinib versus imatinib study in treatment-naïve chronic myeloid leukemia patients trial. *J Clin Oncol* 34: 2333-2340, 2016.
- Stagno F, Vigneri P, Cupri A, Stella S and Di Raimondo F: Personalized strategies for CML patients considering discontinuation of tyrosine kinase inhibitors treatment. *Leuk Res* 36: 1208-1209, 2012.
- Cortes JE, Gambacorti-Passerini C, Deininger MW, Mauro MJ, Chuah C, Kim DW, Dyagil I, Glushko N, Milojkovic D, le Coutre P, *et al*: Bosutinib versus imatinib for newly diagnosed chronic myeloid leukemia: Results from the randomized BFORE trial. *J Clin Oncol* 36: 231-237, 2018.
- Stagno F, Vigneri P, Del Fabro V, Stella S, Cupri A, Massimino M, Consoli C, Tambè L, Consoli ML, Antolino A, *et al*: Influence of complex variant chromosomal translocations in chronic myeloid leukemia patients treated with tyrosine kinase inhibitors. *Acta Oncol* 49: 506-508, 2010.
- Laurent E, Talpaz M, Kantarjian H and Kurzrock R: The BCR gene and philadelphia chromosome-positive leukemogenesis. *Cancer Res* 61: 2343-2355, 2001.
- Burmeister T and Reinhardt R: A multiplex PCR for improved detection of typical and atypical BCR-ABL fusion transcripts. *Leuk Res* 32: 579-585, 2008.
- Gui X, Zhang Y, Pan J, Qiu H, Cen J, Xue Y, Chen S, Shen H, Yao L, Zhang J, *et al*: Chronic myeloid leukemia with e14a3 BCR-ABL transcript: Analysis of characteristics and prognostic significance. *Leuk Lymphoma* 56: 3343-3347, 2015.
- Chisti MM and Sanders DS: Chronic myeloid leukemia with b3a3 (e14a3) fusion: A rare BCR/ABL rearrangement presenting with thrombocytosis-does MTHFR polymorphism matter. *Case Rep Oncol* 11: 485-492, 2018.
- Cross NCP: Detection of BCR-ABL in hematological malignancies by RT-PCR. *Methods Mol Med* 6: 25-36, 1996.
- Vigneri P, Stagno F, Stella S, Cupri A, Forte S, Massimino M, Antolino A, Siragusa S, Mannina D, Impera SS, *et al*: High BCR-ABL/GUS^{IS} levels at diagnosis of chronic phase CML are associated with unfavorable responses to standard-dose imatinib. *Clin Cancer Res* 23: 7189-7198, 2017.
- Vella V, Puppini C, Damante G, Vigneri R, Sanfilippo M, Vigneri P, Tell G and Frasca F: DeltaNp73alpha inhibits PTEN expression in thyroid cancer cells. *Int J Cancer* 124: 2539-2548, 2009.
- Jinawath N, Norris-Kirby A, Smith BD, Gocke CD, Batista DA, Griffin CA and Murphy KM: A rare e14a3 (b3a3) BCR-ABL fusion transcript in chronic myeloid leukemia: Diagnostic challenges in clinical laboratory practice. *J Mol Diagn* 11: 359-363, 2009.
- Massimino M, Consoli ML, Mesuraca M, Stagno F, TIRRÒ E, Stella S, Pennisi MS, Romano C, Buffa P, Bond HM, *et al*: IRF5 is a target of BCR-ABL kinase activity and reduces CML cell proliferation. *Carcinogenesis* 35: 1132-1143, 2014.
- Hu LH, Pu LF, Yang DD, Zhang C, Wang HP, Ding YY, Li MM, Zhai ZM and Xiong S: How to detect the rare BCR-ABL (e14a3) transcript: A case report and literature review. *Oncol Lett* 14: 5619-5623, 2017.
- Cai H, Yang L, Shen K, Zhang W, Xiong J, Zhang M, Mao X, Wang Y and Xiao M: A rare e14a3 BCR/ABL fusion transcript in acute lymphoblastic leukemia patient treated with CAR-modified T-cell therapy. *Oncol Lett* 15: 2491-2494, 2018.
- Buffa P, Romano C, Pandini A, Massimino M, TIRRÒ E, Di Raimondo F, Manzella L, Fraternali F and Vigneri PG: BCR-ABL residues interacting with ponatinib are critical to preserve the tumorigenic potential of the oncoprotein. *FASEB J* 28: 1221-1236, 2014.

26. Massimino M, Stella S, Tirrò E, Romano C, Pennisi MS, Puma A, Manzella L, Zanghì A, Stagno F, Di Raimondo F and Vigneri P: Non ABL-directed inhibitors as alternative treatment strategies for chronic myeloid leukemia. *Mol Cancer* 17: 56, 2018.
27. Yao J, Douer D, Wang L, Arcila ME, Nafa K and Chiu A: A case of acute myeloid leukemia with e6a2 BCR-ABL fusion transcript acquired after progressing from chronic myelomonocytic leukemia. *Leuk Res Rep* 7: 17-19, 2017.
28. Cayuela JM, Rousselot P, Nicolini F, Espinouse D, Ollagnier C, Bui-ThiMH, ChabaneK, RaffouxE, Callet-BauchuE, TigaudI, *etal*: Identification of a rare e8a2 BCR-ABL fusion gene in three novel chronic myeloid leukemia patients treated with imatinib. *Leukemia* 19: 2334-2336, 2005.
29. Stagno F, Vigneri P, Consoli ML, Cupri A, Stella S, Tambè L, Massimino M, Manzella L and Di Raimondo F: Hyperdiploidy associated with a high BCR-ABL transcript level may identify patients at risk of progression in chronic myeloid leukemia. *Acta Haematol* 127: 7-9, 2012.
30. Fujisawa S, Nakamura S, Naito K, Kobayashi M and Ohnishi K: A variant transcript, e1a3, of the minor BCR-ABL fusion gene in acute lymphoblastic leukemia: Case report and review of the literature. *Int J Hematol* 87: 184-188, 2008.
31. Snyder DS, McMahon R, Cohen SR and Slovak ML: Chronic myeloid leukemia with an e13a3 BCR-ABL fusion: Benign course responsive to imatinib with an RT-PCR advisory. *Am J Hematol* 75: 92-95, 2004.
32. Smith KM, Yacobi R and Van Etten RA: Autoinhibition of Bcr-Abl through its SH3 domain. *Mol Cell* 12: 27-37, 2003.