

The use of stoned olive cake and rolled linseed in the diet of intensively reared lambs: effect on the intramuscular fatty-acid composition

M. Mele^{1†}, A. Serra¹, M. Pauselli², G. Luciano³, M. Lanza³, P. Pennisi³, G. Conte¹, A. Taticchi⁴, S. Esposito⁴ and L. Morbidini²

¹Dipartimento di Scienze Agrarie, Alimentari e Agro-ambientali, University of Pisa, Via del Borghetto, 80, 56124 Pisa, Italy; ²Dipartimento di Biologia Applicata, University of Perugia, Borgo XX Giugno, 74, 06123 Perugia, Italy; ³Dipartimento di Scienze delle Produzioni Agrarie e Alimentari, University of Catania, Via Valdisavoia 5, 95123 Catania, Italy; ⁴Dipartimento di Scienze Economico-Estimative e degli Alimenti, University of Perugia, Via San Costanzo snc, 06126 Perugia, Italy

(Received 11 December 2012; Accepted 24 September 2013; First published online 30 October 2013)

The aim of the present study was to evaluate the effect of the inclusion of stoned olive cake and rolled linseed in a concentrate-based diet for lambs on the fatty-acid composition of polar and non-polar intramuscular lipids of the longissimus dorsi muscle. To achieve this objective, 32 Appenninica lambs were randomly distributed into four groups of eight lambs each and were fed conventional cereal-based concentrates (diet C); concentrates containing 20% on a dry matter (DM) basis of rolled linseed (diet L); concentrates containing 35% DM of stoned olive cake (diet OC); and concentrates containing both rolled linseed (10% DM) and stoned olive cake (17% DM; diet OCL). The concentrates were administered together with grass hay at a 20:80 forage:concentrate ratio. Growing performances and carcass traits were evaluated. The fatty-acid composition was analysed in the total intramuscular lipids, as well as in the polar and neutral lipids. The average feed intake and the growth performance of lambs were not affected by the dietary treatments, as a consequence of similar nutritional characteristics of the diets. The inclusion of rolled linseed in the L and OCL diets increased the content of C18:3 n-3 in intramuscular total lipids, which was threefold higher in meat from the L lambs and more than twofold higher in meat from the OCL lambs compared with the C and OC treatments. The n-6:n-3 ratio significantly decreased in the meat from lambs in the L and OCL groups, reaching values below 3. The L treatment resulted in the highest level of trans-18:1 fatty acids in the muscle. Regardless of the dietary treatment, the τ 10-18:1 was the major isomer, representing 55%, 45%, 49% and 45% of total trans-18:1 for C, L, OC and OCL treatments, respectively. Neutral lipids from the OC-fed lambs contained the highest amount of c9-18:1 (more than 36% of total fatty acids); however, the content of c9-18:1 did not differ between the OC and C lambs, suggesting an intensive biohydrogenation of dietary c9-18:1 in the case of OC treatment. The highest content of c9, τ 11-18:2 was detected in the intramuscular fat from the L-fed lambs, followed by the OCL treatment. A similar trend was observed in the neutral lipid fraction and, to a lower extent, in the polar lipids.

Keywords: lamb, fatty acids, olive cake, linseed, intramuscular fat

Implications

Rolling linseed improved the fatty-acid composition of intramuscular lipids towards a healthier profile, by promoting the deposition of polyunsaturated fatty acids and by improving the n-6:n-3 ratio, either alone or together stoned olive cake. In the last case, a reduction in the feeding cost may also be achieved. Finally, compared with a conventional cereal-based concentrate diet, the inclusion of up to 35% of stoned olive cake in the diet did not have detrimental effects on the

growing and slaughtering performances and had minor effects on the intramuscular fatty-acid composition.

Introduction

The fatty-acid composition of intramuscular lipids from intensively reared lambs is characterized by high levels of saturated fatty acids (SFA) and low levels of unsaturated fatty acids (especially polyunsaturated fatty acids (PUFA)) (Berthelot *et al.*, 2010). These characteristics are because of the biohydrogenation activity of rumen bacteria, which

[†] E-mail: marcello.mele@unipi.it

saturate most of the dietary PUFAs. Moreover, as the diet of fattening lambs is constituted mainly of cereals and other grains rich in n-6 fatty acids, the ratio n-6 : n-3 in fat depots of lambs is often unbalanced, reaching values higher than the threshold of 4 : 1 suggested by the World Health Organization in relation to the fat intake in the human diet (WHO, 2003). According to the WHO dietary guidelines, the total intake of SFA should be lowered, whereas the intake of n-3 fatty acids should be enhanced to reduce the risk for cardiovascular diseases. On the basis of the above-mentioned characteristics, the intake of meat from ruminants is often reduced by consumers without considering its overall nutritional value as a source of high biological value proteins, vitamins, minerals and for its content of other potentially bioactive substances such as conjugated linoleic acid (CLA) and branched-chain fatty acids (McAfee *et al.*, 2010). On the basis of the increasing request by consumers for food that is safe and healthy, in the last few years, several feeding strategies, mainly based on the dietary supplementation of unsaturated vegetable oils such as linseed, sunflower and soya bean, have been proposed to improve the fatty-acid composition of lamb meat (Shingfield *et al.*, 2013). In particular, several papers demonstrated that the inclusion of linseed at a percentage higher than 4% (as oil or seed) in the diet of lambs allowed to obtain an increase in the content of n-3 fatty acids (especially 18:3 n-3 and 20:5 n-3), CLA, vaccenic acid and other intermediates of the fatty-acid rumen biohydrogenation process and a decrease of the n-6 : n-3 ratio below the 4 : 1 value (Sinclair, 2007; Moloney *et al.*, 2012). Previous studies reported that dietary PUFA and their biohydrogenation products were selectively incorporated into membrane lipids or into neutral lipid fraction of the intramuscular fat (Jerónimo *et al.*, 2011). As some effect on human health may depend on the fatty-acid composition of membrane lipids assumed with the diets, an increasing interest has been devoted to the characterization of polar and neutral lipids in animal food, especially with regard to the *trans* fatty-acid content (Smith *et al.*, 2009).

The use of agro-industrial by-products as alternative feed resources may be a feasible strategy to reduce feeding cost in ruminant nutrition and, at the same time, to positively influence the quality of the milk and meat, especially in the case of by-products that contain appreciable amounts of vegetable oils and plant secondary compounds, which may contribute to improve the fatty-acid composition and the antioxidant power of meat from ruminants (Vasta and Luciano, 2011). In the Mediterranean area, olive cake is the main by-product of the olive oil industry. Until now, the use of olive cake in the ruminant nutrition has been limited by the high content of lignin, which worsens the digestibility of organic matter (Molina-Alcaide and Yáñez-Ruiz, 2008). Modern extraction technologies diffused in Italy are based on a three-phase extraction system at a low level of water addition that may include the preventive stoning of the olives to improve the quality of the oil. This technology allows to better control the activity of the peroxidase enzyme, which is mainly concentrated in the seed, otherwise virgin olive cake

can be stoned after oil mechanical extraction. Finally, coupling the three-phase extraction with a total or partial removal of seeds (stoning) allows to obtain an olive cake containing low levels of lignin and remarkable levels of residual oil and antioxidants, such as tocopherols and retinol, and bioactive phenols, such as secoiridoids and lignans (Servili *et al.*, 2011). Therefore, the use of stoned olive cake in substitution to a quote of the cereal in the diet of lamb could be a strategy to reduce the feeding cost and, at the same time, to introduce a source of oleic acid and of substances with antioxidant properties, which could contribute to improve the quality of meat (Molina-Alcaide and Yáñez-Ruiz, 2008).

In the present research, the effect of stoned olive cake supplementation, alone or in combination with rolled linseed, on the fatty-acid composition of polar and non-polar intramuscular lipids from the *longissimus dorsi* muscle of intensively reared lambs was evaluated.

Material and methods

Stoned olive cake production and polyphenols determination

Fresh olive cake was obtained after oil extraction was performed using an RCM Rapanelli three-phase decanter (Rapanelli Inc., Perugia, Italy). After oil extraction, virgin olive cakes were stoned and dried using a fluid-bed dryer. The initial temperature of the flow drying air was 120°C and did not exceed 45°C to prevent from oxidative deterioration (Servili *et al.*, 2011). The extraction of phenolic compounds from olive cake was performed according to Dal Bosco *et al.* (2012). Briefly, 10 g of the sample were homogenized with 100 ml of a solution containing 20 mg/l of sodium diethyl-dithiocarbamate in methanol; the extraction was performed in triplicate. After methanol removal, the aqueous extract was used for phenol separation by solid phase extraction (SPE) technique. The SPE procedure was applied by loading 1 ml of the aqueous extract into an Extractclean high-load C18 cartridge (Alltech Italia S.r.l., Milano, Italy) and eluting with methanol. After solvent removal under vacuum at 30°C, the phenolic extract was recovered and then dissolved in methanol. The HPLC analysis of the phenolic extracts was performed according to Dal Bosco *et al.* (2012).

Animals and diets

The study was carried out at the Experimental Station of the Department of Applied Biology at the University of Perugia (Italy) on 32 Appenninica male lambs that immediately after weaning (40 ± 5 days old, weighing 17.8 ± 1.6 kg) were randomly distributed into four groups of eight lambs each. After 20 days of adaptation, each group was kept in multiple pens (eight lambs for each pen) and was allowed one of the following dietary treatments for 40 days: a concentrate with a low level of lipids and a high content of non-structural carbohydrates (diet control, C); a concentrate containing 20% on a dry matter (DM) basis of rolled linseed (diet L); a concentrate containing 35% on DM of stoned olive cake (diet OC); and a concentrate containing a mixture of rolled linseed

(10% on DM) and stoned olive cake (17% on DM; diet OCL). All the concentrates were pelleted by a pelleting machine (CMS IEM; Colognola ai Colli, Verona, Italy), the pellet diameter was 3 mm and the pelleting temperature ranged from 35°C to 40°C. Diets were balanced according to energy and protein content to obtain isoenergetic and isonitrogenous treatments. The concentrates were administered together with grass hay according to the ratio 20:80 forage:concentrate, on the basis of the expected DM intake. The composition and the chemical characteristics of the diet are reported in Table 1. Feeds offered and refused were daily recorded per group andorts collected and mixed. Each week feed samples and orts were collected and stored at -30°C before the analysis.

CP, ether extract and ash were determined in feeds according to AOAC (1990) methods. Fibre fractions were analysed according to the method described by Van Soest *et al.* (1991). The metabolizable energy content of the experimental concentrates was estimated according to Cornell Net Carbohydrates and Protein System for sheep (Cannas *et al.*, 2004).

Lambs were weighed at the beginning of the experiment and every 10 days until the day before slaughter. At the end of the experiment, lambs were slaughtered and carcasses were immediately weighed to obtain hot carcass weight and kept at 4°C for 24 h. Samples from the *longissimus dorsi* muscle were removed between the 2nd and the 13th rib, vacuum-packed and stored at -80°C until further analysis.

Fatty-acid analysis

Total intramuscular lipids (TL) were extracted by means of a chloroform/methanol solution (2:1, v/v), according to Serra *et al.* (2009a). Polar (PL) and neutral (NL) lipids were separated according to Juaneda and Rocquelin (1985) using Sep-Pak® silica cartridges (Waters, Milford, MA, USA).

For each lipid fraction (i.e. TL, PL and NL), fatty-acid methyl esters (FAME) were prepared by the base-catalysed *trans*-methylation (Christie, 1982). Before methylation, 9:0 and 23:0 FAME were added together as internal standards. The FAMEs were determined by a GC 8000 Top ThermoQuest (Milan, Italy) gas-chromatograph apparatus equipped with a flame ionization detector (FID) and a high-polar fused silica capillary column (WCOT-fused silica CP-Select CB for FAME Varian, Middelburg, the Netherlands; 100 m × 0.25 mm i.d.; film thickness 0.25 µm). Helium was used as the carrier gas at a flow of 1 ml/min. The split ratio was 1:80. The oven temperature was programmed at 150°C and held for 1 min, then increased up to 175°C at a rate of 0.8°C/min, held for 14 min, increased up to 188°C at 2°C/min, held for 18 min and then increased up to 230°C at a rate of 2°C/min, held for 13 min. The injector and detector temperatures were set at 270°C and 300°C, respectively. The identification of individual FAME was based on a standard mixture of 52 Component FAME Mix (Nu-Chek Prep Inc., Elysian, MN, USA) and 77 individual FAME standards (Larodan Fine Chemicals, Malmo, Sweden). The identification of 18:1 and 18:2 isomers was based on commercial standard mixtures

(Larodan Fine Chemicals) and on chromatograms published by Kramer *et al.* (2008) and Alves and Bessa (2007). For each FA, response factors to FID and inter- and intra-assay coefficients of variation were calculated by using a reference standard butter (CRM 164, Community Bureau of Reference, Brussels, Belgium). Fatty acids were expressed as g/100 g of fatty acids.

With regard to feed samples, fat was extracted according to the method described by Folch *et al.* (1957). Fatty acids were esterified according to Christie (1982) with 19:0 as the internal standard, and were identified using the same procedure described above for fatty acids of meat samples. The fatty-acid composition of the four experimental concentrates is presented in Table 1.

Statistical analysis

Data were analysed by means of one-way analysis of variance (SAS, 1999), including diet as a fixed factor. Individual animals were considered experimental units for all the variables included in the statistical analysis and reported in the tables. Multiple comparisons of means were made using the Tukey's adjustment. Main effects and differences were considered significant when $P \leq 0.05$.

Results and discussion

Chemical composition of experimental diets

The chemical composition of olive cake is highly variable according to cultivar, oil extraction and stoning system applied (Molina-Alcaide and Yáñez-Ruiz, 2008). In the present trial, the olive cake was obtained by a three-phase extraction system coupled with a stoning system after the oil extraction. The values of NDF, ADF, ADL and CP content were higher than those found by Vera *et al.* (2009) in stoned olive cake obtained from two-phase system, whereas ether extract was lower (Supplementary Table S1). As expected, the fatty-acid composition of olive cake was high in c9-18:1 (more than 75% of total fatty acids). The low temperature used in the drying system determined that total polyphenol content was higher than that reported by Molina-Alcaide and Yáñez-Ruiz (2008) in their review, concerning the potential use of olive oil extraction by-products in animal nutrition. The most abundant polyphenols were 3,4-dihydroxyphenylethanol-elenolic acid di-aldehyde (3,4-DHPEA-EDA, also known as oleuropein-aglycone di-aldehyde) and verbascoside (Supplementary Table S1). Similar results were reported by Dal Bosco *et al.* (2012). The effect of olive cake polyphenols on the oxidative stability of lamb meat has been reported in a previous paper (Luciano *et al.*, 2013).

The inclusion of both rolled linseed and olive cake in the L, OC and OCL concentrates resulted in a higher content of fat, compared with C concentrates (Table 1). Furthermore, the inclusion of olive cake in the concentrates (OC and OCL treatments) increased the NDF, ADF and ADL content, compared with the C and L diets. The inclusion of olive cake in the OC and OCL diets resulted in a high content of c9-18:1,

Table 1 *Ingredients, chemical composition and fatty acids of the experimental concentrates and of the hay administered to the lambs*

	Grass hay	Experimental concentrates			
		C	L	OC	OCL
Ingredients (g/100 g of DM)					
Oat		35.0	18.0	18.0	18.0
Horse bean		10.0	6.0	17.0	6.0
Wheat bran		5.0	38.0	4.0	25.0
Corn gluten meal		11.0	6.0	13.0	11.0
Barley		37.0	10.0	11.0	11.0
Rolled linseed		–	20.0	–	10.0
Stoned olive cake		–	–	35.0	17.0
Mineral premix		2.0	2.0	2.0	2.0
Chemical composition					
DM ¹	93.34	87.45	87.74	88.70	87.85
CP ²	7.25	20.15	20.30	20.90	21.50
Ether extract ²	2.32	2.58	10.85	9.96	10.55
NDF ²	61.50	21.82	25.89	33.54	29.51
ADF ²	36.31	9.65	10.51	18.63	14.95
ADL ²	4.28	1.86	2.76	8.19	7.77
Ash ²	4.25	8.25	8.33	9.20	8.74
ME ³	7.99	12.02	12.81	12.06	12.52
Fatty acids (g/100 g of total fatty acids)					
16:0	20.15	21.08	9.49	14.68	12.42
18:0	6.15	1.62	1.80	1.95	2.70
18:1 c9	17.99	25.20	13.95	60.44	39.88
18:2 n-6	37.37	45.15	24.95	19.20	21.22
18:3 n-3	15.81	2.85	45.73	1.77	21.45

C = control concentrate containing neither linseed nor olive cake; L = concentrate containing 20% of rolled linseed; OC = concentrate containing 35% of stoned olive cake; OCL = concentrate containing 17% olive cake and 10% rolled linseed; DM = dry matter; ME = metabolizable energy.

¹Expressed as g/100 g of fresh weight.

²Expressed as g/100 g of DM.

³Expressed as MJ/kg of DM.

Table 2 *Effect of stoned olive cake and rolled linseed in the diet of intensively reared lambs on growth performance and carcass composition*

Items	Dietary treatments ¹				s.e.m.	P-values
	C	L	OC	OCL		
Number of lambs	8	8	8	8		
Initial live weight (kg)	17.50	17.88	17.93	17.72	0.61	0.900
Live weight at slaughter (kg)	29.52	27.67	28.52	29.23	1.00	0.750
Average daily gain (g/day)	204	165	180	196	13	0.560
Cold carcass weight (kg)	12.78	12.53	12.61	12.95	0.50	0.960

C = containing neither linseed nor olive cake; L = containing 20% of rolled linseed; OC = containing 35% of stoned olive cake; OCL = containing 10% of rolled linseed and 17% of stoned olive cake.

¹The dietary treatments consisted of grass hay and the experimental concentrates at a 20 : 80 forage : concentrate ratio.

whereas the supplementation with rolled linseed (L and OCL diets) increased the levels of 18:3 n-3.

Animal performance and carcass characteristics

The growth performance of lambs was not affected by the dietary treatments (Table 2), as a consequence of the similar nutritional characteristics of the four diets that allowed to obtain very similar values of DM intake (944.2, 914.4, 968.3

and 939.1 g/day of DM, for group C, L, OC and OCL, respectively).

In a previous study, Mioc *et al.* (2007) reported no adverse effects of the inclusion at 15% on DM intake of whole dried and whole olive cake in the diet of finishing lambs, respectively; although when olive cake was added at 30% on DM intake, an adverse effect on the average daily gain was found. In the present study, the OC diet contained nearly

25% of olive cake. This amount of olive cake had been proposed as the upper limit in the diet of sheep (Molina-Alcaide and Yáñez-Ruiz, 2008). However, the lack of negative effects on the growing performance suggested that the stoning process could have improved the digestibility of the diet as already observed by Sadeghi *et al.* (2009), using experimental diets containing 20% of stoned olive cake. The dietary treatment did not affect slaughtering performance according to previous studies that included comparable amounts of stoned olive cake or vegetable fat in the diet of finishing lambs (Mioc *et al.*, 2007; Berthelot *et al.*, 2010; Noci *et al.*, 2011).

Fatty-acid composition of intramuscular fat

As a consequence of the similar growth rate of lambs, the content of intramuscular fat in the *Longissimus dorsi* muscle was not affected by the dietary treatment (3.82, 3.80, 3.93 and 3.61 for C, L, OC and OCL diet, respectively, $P=0.96$; data not shown). Therefore, the content of total fatty acids and the proportion between polar and neutral fatty acids did not differ between treatments (Tables 3 to 5). This result allowed to avoid confounding effects of total fatness on meat fatty-acid composition (Noci *et al.*, 2011). Therefore, differences in fatty-acid composition were mainly because of the diet and the selectivity of individual fatty acids for polar and neutral lipids.

As expected, the inclusion of rolled linseed in the L and OCL diets resulted in an increase of 18:3 n-3 content in intramuscular total lipids, which was threefold higher in meat from the L lambs and more than twofold higher in meat from the OCL lambs, compared with that from lambs fed C and OC diets ($P<0.01$ in all cases). The content of long-chain PUFA n-3 was also enhanced, with the exception of 22:5 n-3 and 22:6 n-6. In particular, the 20:3 n-3 was detected only in total lipids of meat samples from the L and OCL lambs and, as the other long-chain PUFAs, was preferentially incorporated into the polar lipid fraction (Table 4). The presence of this fatty acid only in the intramuscular fat from lamb fed with linseed suggested a possible role of 20:3 n-3 as a marker of linseed supplementation. Moreover, dietary linseed enhanced the content of 20:5 n-3 in both intramuscular fat and polar PL from the L and OCL lambs, but not the content of 22:6 n-3, confirming what was reported in previous studies, about the long time needed to complete the elongation process from 18:3 n-3 to 22:6 n-3 (Scollan *et al.*, 2006; Sinclair, 2007). The overall effect of the L diet and, to a lower extent, of the OCL diet, was to increase the degree of unsaturation of the polar lipids of muscle cell membranes (Table 4). Although the content of 18:3 n-3 in the L diet was double than that in the OCL diet, the transfer of this fatty acid from diet to intramuscular fat was not proportional to the dietary intake, suggesting a lower apparent transfer of C18:3 n-3 from the L diet, when compared with the OCL diet (Table 3). This phenomenon was also confirmed by the polar lipid composition (Table 4). This result did not agree with Bas *et al.* (2007), who observed a linear increase in 18:3 n-3 content in intramuscular fat with the increase of extruded linseed in the concentrate from 0% to 9% on DM basis.

As the results of the present study did not allow to assess whether this trend was because of an interactive effect between stoned olive cake and rolled linseed, further studies are needed to examine the effect of olive polyphenols on the rumen biohydrogenation of 18:3 n-3. In fact, previous studies reported that some kinds of plant polyphenols might influence rumen microbe metabolism, affecting the biohydrogenation process of the dietary unsaturated fatty acids (Vasta *et al.*, 2009; Buccioni *et al.*, 2011; Jerónimo *et al.*, 2012).

Despite the high amount of rolled linseed included in our study on the L and OCL concentrates (20% and 10% of the concentrate, respectively), the enrichment of intramuscular fat with 18:3 n-3 was not as high as expected. Bas *et al.* (2007) obtained similar levels of enrichment in the intramuscular fat of lambs fed a concentrate with 9% of extruded linseed (1.52/100 g of fatty acids), whereas Berthelot *et al.* (2010) reported higher levels of 18:3 n-3 (more than 2.2/100 g of fatty acids), by feeding lambs with a concentrate containing 10% of extruded linseed. However, it is not easy to compare different feeding studies because the composition of the basal diet or the nature of the dietary fat source may significantly affect the biohydrogenation process of PUFAs in the rumen (Shingfield *et al.*, 2013). A higher duodenal flow of 18:3 n-3 and its biohydrogenation intermediates have been observed when extruded linseed or pure linseed oil have been compared with rolled linseed in the cow diet (Doreau *et al.*, 2009). Moreover, when lamb carcasses were very lean, as in the case of the study by Berthelot *et al.* (2010), the contribution of those fatty acids, which are selectively incorporated in the polar lipids (such as 18:3 n-3 and 18:2 n-6, Jerónimo *et al.*, 2011), increased.

The n-6:n-3 ratio significantly decreased in the meat samples from the L and OCL lambs reaching values below 3, which is considered an optimal value for human nutrition, although recent insights suggest that absolute amounts of n-3 and n-6 PUFA are more useful than the n-6:n-3 ratio as indices for cardiovascular diseases (McAfee *et al.*, 2010).

The content of $\text{c9}, \text{t11}, \text{c15-18:3}$ and $\text{t11}, \text{c15-18:2}$, which represent the first two intermediates of 18:3 n-3 rumen biohydrogenation, proportionally increased with increasing content of linseed in the concentrate. The content of $\text{t11}, \text{c15-18:2}$ was higher in neutral lipid than in polar lipids of L and OCL samples (Tables 4 and 5), confirming a higher affinity of this fatty acid for neutral lipids, as previously reported by Jerónimo *et al.* (2011).

The content of *trans*-18:1 fatty acids in the intramuscular fat decreased with a reduction in the degree of unsaturation of the dietary fat. In fact, the muscles from the L-fed lambs showed the highest level of *trans*-18:1 ($P<0.001$), followed by the muscle from lambs in the OCL, OC and C groups (Table 3). In all cases, t10-18:1 was the major isomer, representing 55%, 45%, 49% and 45% of total *trans*-18:1 fatty acids for C, L, OC and OCL treatments, respectively. High levels of t10-18:1 in the intramuscular fat have been previously reported in intensively reared lambs fed with a forage:concentrate ratio similar to that applied in the

Table 3 Effect of stoned olive cake and rolled linseed in the diet of intensively reared lambs on the concentration of total fatty acids (mg/g fresh muscle) and on the fatty-acid composition (g/100 g of fatty acids) of total lipids in the *Longissimus dorsi* muscle

	Dietary treatments ¹				s.e.m.	P-values
	C	L	OC	OCL		
Total fatty acids	28.95	28.54	29.89	27.14	3.69	0.852
Fatty-acid composition						
14:0	4.49	4.53	3.96	4.64	0.33	0.665
14:1 t9	0.07 ^a	0.05 ^a	0.03 ^b	0.03 ^b	0.01	0.002
14:1 c9	0.15	0.14	0.11	0.10	0.01	0.062
14-iso	0.06	0.05	0.04	0.05	0.01	0.278
15:0	0.54	0.62	0.42	0.51	0.01	0.143
15-anteiso	0.18	0.19	0.14	0.16	0.02	0.338
15-iso	0.14	0.10	0.08	0.09	0.01	0.095
16:0	22.37 ^a	19.51 ^b	19.99 ^b	20.42 ^b	0.61	0.045
16-iso	0.22 ^a	0.17 ^{ab}	0.12 ^b	0.14 ^b	0.01	0.007
16:1 c9	1.85 ^a	1.39 ^b	1.37 ^b	1.16 ^b	0.06	0.001
16:1 t9	0.09 ^b	0.13 ^a	0.06 ^b	0.08 ^b	0.01	0.001
16:1 c7	0.46	0.48	0.49	0.45	0.02	0.829
Other 16:1	0.32 ^a	0.35 ^a	0.23 ^b	0.24 ^b	0.01	0.001
17:0	1.31 ^{ab}	1.40 ^a	0.97 ^b	1.19 ^b	0.07	0.015
17:1 c9	0.60 ^a	0.52 ^a	0.38 ^b	0.37 ^b	0.02	0.001
17-anteiso	0.60 ^a	0.49 ^{ab}	0.37 ^b	0.44 ^{ab}	0.03	0.007
17-iso	0.47 ^a	0.42 ^{ab}	0.31 ^b	0.38 ^{ab}	0.03	0.027
18:0	15.19 ^b	15.13 ^b	17.68 ^a	19.26 ^a	0.74	0.017
18-iso	0.17 ^a	0.11 ^b	0.08 ^b	0.10 ^b	0.01	0.001
18:1 c9	34.23 ^a	27.36 ^b	34.17 ^a	28.01 ^b	0.73	0.001
18:1 c11	1.03 ^b	1.00 ^b	1.36 ^a	1.01 ^b	0.07	0.037
18:1 c12	0.38 ^{bc}	0.75 ^a	0.23 ^c	0.52 ^b	0.03	0.001
18:1 c13	0.10 ^b	0.23 ^a	0.08 ^c	0.14 ^b	0.01	0.001
18:1 c14	0.06 ^b	0.14 ^a	0.05 ^b	0.06 ^b	0.01	0.001
18:1 c15	0.07 ^{bc}	0.13 ^a	0.06 ^c	0.10 ^b	0.01	0.001
Total 18:1 cis	35.90 ^a	29.60 ^b	35.95 ^a	29.80 ^b	0.77	0.001
18:1 t4	0.01 ^c	0.02 ^b	0.04 ^a	0.04 ^a	0.01	0.001
18:1 t5	0.03 ^c	0.04 ^{bc}	0.08 ^a	0.06 ^{ab}	0.01	0.001
18:1 t6-8	0.32 ^b	0.54 ^a	0.62 ^a	0.54 ^a	0.03	0.001
18:1 t9	0.29 ^b	0.43 ^a	0.38 ^{ab}	0.36 ^{ab}	0.02	0.001
18:1 t10	2.74 ^b	5.00 ^a	3.54 ^{ab}	3.73 ^{ab}	0.15	0.001
18:1 t11	0.98 ^b	2.34 ^a	1.10 ^b	1.48 ^{ab}	0.11	0.001
18:1 t12	0.39 ^b	1.02 ^a	1.01 ^a	0.95 ^a	0.05	0.001
18:1 t15	0.14 ^c	0.70 ^a	0.17 ^c	0.48 ^b	0.02	0.001
18:1 t16	0.24 ^c	0.81 ^a	0.22 ^c	0.54 ^b	0.03	0.001
Total 18:1 trans	5.15 ^c	10.90 ^a	7.15 ^{bc}	8.20 ^b	0.54	0.001
18:2 c9, t11	0.54 ^b	0.87 ^a	0.50 ^b	0.58 ^{ab}	0.03	0.001
Other 18:2 conjugated	0.05 ^b	0.13 ^a	0.05 ^b	0.09 ^{ab}	0.01	0.001
18:2 t11, c15	0.16 ^c	1.93 ^a	0.15 ^c	0.97 ^b	0.06	0.001
18:2 n-6	4.93	4.82	5.11	5.04	0.43	0.765
Other 18:2 non-conjugated	0.38 ^c	1.28 ^a	0.34 ^c	0.79 ^{ab}	0.01	0.001
18:3 n-3	0.52 ^c	1.65 ^a	0.63 ^c	1.33 ^b	0.07	0.001
18:3 c9, t11, c15	0.00 ^b	0.05 ^a	0.00 ^b	0.03 ^a	0.01	0.001
20:0	0.16	0.17	0.20	0.19	0.01	0.161
20:1 c9	0.15	0.12	0.18	0.15	0.01	0.330
20:2 n-6	0.05	0.03	0.04	0.05	0.01	0.222
20:3 n-3	0.00 ^b	0.01 ^a	0.00 ^b	0.01 ^a	0.01	0.001
20:3 n-6	0.10	0.09	0.13	0.10	0.01	0.397
20:4 n-6	1.06	0.89	1.25	1.09	0.14	0.738
20:5 n-3	0.16 ^b	0.30 ^a	0.18 ^b	0.27 ^a	0.01	0.025
21:0	0.02	0.02	0.01	0.02	0.01	0.467
22:0	0.02	0.02	0.03	0.03	0.01	0.687
22:4 n-6	0.10 ^a	0.05 ^b	0.09 ^a	0.06 ^b	0.01	0.030

Table 3: (Continued)

	Dietary treatments ¹				s.e.m.	P-values
	C	L	OC	OCL		
22:5 n-6	0.03 ^a	0.01 ^b	0.04 ^a	0.02 ^b	0.01	0.024
22:5 n-3	0.32	0.36	0.31	0.37	0.01	0.248
22:6 n-3	0.11	0.12	0.14	0.15	0.01	0.684
Other fatty acids	0.87	0.76	0.70	0.83	0.02	0.564
SFAs	44.10	41.40	43.30	46.25	1.39	0.251
BCFAs	1.80 ^a	1.55 ^b	1.15 ^c	1.35 ^{bc}	0.11	0.008
MUFAs	44.60 ^a	43.65 ^{ab}	45.90 ^a	40.60 ^b	1.05	0.020
PUFAs	7.91 ^b	11.61 ^a	8.40 ^b	10.29 ^{ab}	0.76	0.007
n-6 : n-3 ratio	5.70 ^a	2.40 ^b	5.25 ^a	2.95 ^b	0.12	0.001

C = containing neither linseed nor olive cake; L = containing 20% of rolled linseed; OC = containing 35% of stoned olive cake; OCL = containing 10% of rolled linseed and 17% of stoned olive cake; SFA = saturated fatty acid; BCFA = branched-chain fatty acid; MUFA = monounsaturated fatty acid; PUFA = polyunsaturated fatty acid. ^{a-c}Within a row, values with different letters are significantly different ($P \leq 0.05$).

¹The dietary treatments consisted of grass hay and the experimental concentrates at a 20 : 80 forage : concentrate ratio.

present study, especially when unsaturated fat has been added to the diet (Daniel *et al.*, 2004; Bas *et al.*, 2007). All *trans*-18:1 isomers were preferentially incorporated in neutral lipids, where they represented 5.06%, 10.89%, 7.49% and 8.57% of the total fatty acids for C, L, OC and OCL treatments, respectively (Table 5). On the contrary, *trans*-18:1 fatty acids in polar lipids ranged from a minimum of 2.15% (C treatment) to a maximum of 4.92% (L treatment) (Table 4). Two isomers (*t*4- and *t*5-18:1) were detected only in neutral lipids. The higher affinity of *trans* fatty acids for neutral lipids was previously reported, especially when the lambs were fed a diet enriched with 18:3 n-3 (Jerónimo *et al.*, 2011; Noci *et al.*, 2011). According to the pattern of *trans*-18:1, the biohydrogenation of dietary *c*9-18:1 contained in the O and OL diets resulted in an accumulation of *t*5-18:1 *t*6-8-18:1, *t*9-18:1, *t*10-18:1 and *t*12-18:1, whereas the content of *t*11-18:1 did not significantly vary when compared with C treatment. Previous *in vitro* studies demonstrated that the biohydrogenation of *c*9-18:1 may produce a large spectrum of *trans* fatty acids from 6 to 16 double-bond position (Mosley *et al.*, 2002; Buccioni *et al.*, 2006), but with a lower efficiency in the case of *t*11-18:1 (Mosley *et al.*, 2002). In total lipids, the content of *c*9-18:1 did not differ between the OC- and C-fed lambs, suggesting an intensive biohydrogenation of dietary *c*9-18:1 in the case of the OC treatment. As a consequence, the level of 18:0 (the end product of biohydrogenation) in the intramuscular fat of the OC lambs was significantly higher than that of the C lambs. The lack of a significant effect of dietary olive cake on the *c*9-18:1 content in the intramuscular total lipids did not agree with previous studies on dairy sheep. Often, in fact, the inclusion of crude olive cake in the diet of dairy sheep resulted in an increase of *c*9-18:1 in milk fat (Molina-Alcaide and Yáñez-Ruiz, 2008). This result may be partly explained by the different composition between intramuscular and milk fat and by the different affinity of *c*9-18:1 to polar and neutral lipids. Milk fat is mainly composed of neutral lipids, whereas the intramuscular fat contains appreciable amount

of polar lipids, which can significantly contribute to the fatty-acid composition of total intramuscular lipids.

Although *t*10-18:1 was the predominant isomer among *trans* fatty acids, the content of *t*10,*c*12-CLA, which is the precursor of *t*10-18:1 in the biohydrogenation process, was negligible (data not shown). In the present study, the highest levels of *t*10-18:1 was found in the intramuscular fat from the L lambs, as a consequence of the biohydrogenation of 18:3 n-3. High levels of *t*10-18:1 were previously reported as effect of the inclusion of unsaturated vegetable oils in concentrate-rich diets (Shingfield *et al.*, 2013). However, as stated above, *t*10-18:1 may also originate from the biohydrogenation of *c*9-18:1 contained in the OC, as confirmed by the high levels of *t*10-18:1 also in intramuscular fat from the OC and OCL lambs (Table 3). The *c*9,*t*11-CLA was the predominant CLA isomer, representing from 86% to 90% of total CLA (Table 3). The highest content of *c*9,*t*11-CLA was detected in the intramuscular fat from the L lambs, followed by the OCL lambs, which showed an intermediate concentration between the L and C and OC lambs (Table 3). A similar trend was observed in the neutral lipid fraction and, to a lower extent, in the polar lipids (Tables 4 and 5). The trend observed for the levels of *c*9,*t*11-CLA can be associated with the content of *t*11-18:1, because the Stearoyl Co-A desaturase enzyme converts *t*11-18:1 to *c*9,*t*11-CLA in lamb tissues (Palmquist *et al.*, 2004). According to this pattern, the inclusion of linseed in the diet resulted in an increase in *t*11-18:1, which, in turn, led to an increase in CLA content in the intramuscular fat, especially in the neutral lipids. Similar results have been previously obtained by feeding lambs diets supplemented with linseed oil (Noci *et al.*, 2011). The CLA and *t*11-18:1 are considered potential bioactive compounds for human health (Mele and Banni, 2010) and the use of linseed in the diet of lambs may be a feasible strategy to enrich meat with these fatty acids. However, till now the optimal level of enrichment has not been well defined, although some recent data on dairy products seem to indicate a beneficial effect of CLA on human lipid profile

Table 4 Effect of stoned olive cake and rolled linseed in the diet of intensively reared lambs on the concentration of total fatty acids (mg/g fresh muscle) and on the fatty-acid composition (g/100 g of fatty acids) of polar lipids in the *Longissimus dorsi* muscle

	Dietary treatments ¹				s.e.m.	P-values
	C	L	OC	OCL		
Total fatty acids	6.60	6.84	6.52	6.13	0.67	0.563
Fatty-acid composition						
14:0	0.32	0.46	0.30	0.52	0.08	0.174
15:0	0.23	0.22	0.16	0.20	0.02	0.055
15-anteiso	0.04	0.04	0.03	0.05	0.01	0.131
15-iso	0.03	0.02	0.01	0.02	0.01	0.245
16:0	12.77	13.95	15.11	14.47	1.24	0.556
16-iso	0.12	0.19	0.05	0.06	0.06	0.339
16:1 c9	0.81 ^a	0.47 ^b	0.43 ^b	0.41 ^b	0.05	0.001
16:1 t9	0.12 ^b	0.33 ^a	0.09 ^b	0.14 ^b	0.02	0.001
16:1 c7	0.23 ^a	0.17 ^b	0.16 ^b	0.15 ^b	0.01	0.002
Other 16:1	0.24 ^b	0.41 ^a	0.20 ^b	0.28 ^b	0.03	0.001
17:0	0.81 ^a	0.66 ^{ab}	0.50 ^b	0.52 ^b	0.05	0.001
17:1 c9	0.47 ^a	0.29 ^b	0.23 ^b	0.20 ^b	0.04	0.001
17-anteiso	0.23 ^a	0.16 ^{ab}	0.10 ^b	0.14 ^b	0.02	0.002
17-iso	0.44	0.43	0.31	0.33	0.06	0.347
18:0	15.73	15.13	16.32	16.80	0.56	0.363
18-iso	0.08 ^a	0.05 ^{ab}	0.03 ^b	0.03 ^b	0.01	0.001
18:1 c9	24.58 ^a	16.48 ^b	23.68 ^a	19.19 ^b	1.13	0.001
18:1 c11	2.69 ^{ab}	2.47 ^b	3.15 ^a	2.91 ^{ab}	0.16	0.031
18:1 c12	0.65 ^c	1.91 ^a	0.48 ^c	1.10 ^b	0.10	0.001
18:1 c13	0.10 ^b	0.16 ^a	0.06 ^c	0.10 ^b	0.01	0.001
18:1 c15	0.22 ^b	0.38 ^a	0.10 ^c	0.20 ^b	0.01	0.001
18:1 t6-8	0.16	0.20	0.24	0.19	0.02	0.172
18:1 t9	0.14	0.21	0.16	0.21	0.03	0.186
18:1 t10	1.01 ^b	1.99 ^a	1.16 ^{ab}	1.27 ^{ab}	0.26	0.032
18:1 t11	0.40 ^b	1.13 ^a	0.59 ^b	0.62 ^b	0.08	0.001
18:1 t12	0.22 ^b	0.59 ^a	0.57 ^a	0.55 ^a	0.04	0.001
18:1 t15	0.11 ^c	0.34 ^a	0.12 ^c	0.25 ^b	0.01	0.001
18:1 t16	0.12 ^c	0.45 ^a	0.11 ^c	0.30 ^b	0.01	0.001
18:2 c9, t11	0.17 ^b	0.25 ^a	0.12 ^b	0.17 ^b	0.01	0.001
Other 18:2 conjugated	0.24	0.19	0.25	0.22	0.03	0.409
18:2 t11, c15	0.32 ^b	0.56 ^a	0.31 ^b	0.40 ^{ab}	0.01	0.001
18:2 n-6	22.38	22.32	21.41	22.85	1.52	0.386
Other 18:2 non-conjugated	0.24 ^b	0.68 ^a	0.17 ^b	0.37 ^{ab}	0.04	0.001
18:3 n-3	1.02 ^c	4.07 ^a	1.04 ^c	3.00 ^b	0.11	0.001
18:3 c9, t11, c15	0.00 ^b	0.06 ^a	0.00 ^b	0.03 ^{ab}	0.01	0.001
20:0	0.19	0.17	0.18	0.16	0.01	0.098
20:1 c9	0.06	0.04	0.05	0.04	0.01	0.704
20:2 n-6	0.16	0.14	0.14	0.12	0.01	0.339
20:3 n-3	0.01 ^b	0.08 ^a	0.01 ^b	0.09 ^a	0.01	0.001
20:3 n-6	0.76	0.64	0.74	0.65	0.02	0.148
20:4 n-6	7.22	6.42	7.28	6.20	0.22	0.131
20:5 n-3	0.98 ^b	1.97 ^a	0.98 ^b	1.47 ^{ab}	0.13	0.001
22:0	0.04	0.04	0.05	0.05	0.01	0.779
22:4 n-6	0.63 ^a	0.32 ^b	0.47 ^{ab}	0.32 ^b	0.05	0.001
22:5 n-6	0.16 ^a	0.05 ^b	0.14 ^a	0.06 ^b	0.01	0.001
22:5 n-3	1.74 ^b	2.34 ^a	1.62 ^b	1.91 ^{ab}	0.13	0.001
22:6 n-3	0.41 ^b	0.58 ^a	0.47 ^b	0.60 ^a	0.01	0.045
Other fatty acids	0.11	0.15	0.14	0.16	0.01	0.895

C = containing neither linseed nor olive cake; L = containing 20% of rolled linseed; OC = containing 35% of stoned olive cake; OCL = containing 10% of rolled linseed and 17% of stoned olive cake.

^{a-c}Within a row, values with different letters are significantly different ($P \leq 0.05$).

¹The dietary treatments consisted of grass hay and the experimental concentrates at a 20 : 80 forage : concentrate ratio.

Table 5 Effect of stoned olive cake and rolled linseed in the diet of intensively reared lambs on the concentration of total fatty acids (mg/g fresh muscle) and on the fatty-acid composition (g/100 g of fatty acids) of neutral lipids in the Longissimus dorsi muscle

	Dietary treatments ¹				s.e.m.	P-values
	C	L	OC	OCL		
Total fatty acids	22.36	21.69	23.37	21.02	3.22	0.568
Fatty-acid composition						
14:0	4.59	4.47	3.88	4.15	0.33	0.411
14:1 n9	0.08 ^a	0.06 ^a	0.04 ^b	0.04 ^b	0.01	0.003
14:1 n9	0.20	0.18	0.14	0.15	0.02	0.102
14-iso	0.07	0.06	0.05	0.04	0.01	0.478
15:0	0.58	0.67	0.48	0.53	0.05	0.07
15-anteiso	0.19	0.22	0.16	0.19	0.02	0.132
15-iso	0.15	0.12	0.09	0.11	0.02	0.096
16:0	23.59 ^a	19.91 ^b	21.08 ^b	20.16 ^b	0.55	0.001
16-iso	0.20 ^a	0.17 ^{ab}	0.13 ^b	0.15 ^b	0.02	0.005
16:1 n9	1.83 ^a	1.52 ^b	1.37 ^b	1.33 ^b	0.09	0.002
16:1 n9	0.10 ^b	0.14 ^a	0.08 ^b	0.08 ^b	0.01	0.030
16:1 n7	0.35	0.41	0.43	0.39	0.03	0.271
Other 16:1	0.23 ^b	0.37 ^a	0.22 ^b	0.34 ^a	0.03	0.004
17:0	1.16 ^{ab}	1.39 ^a	0.99 ^b	1.11 ^{ab}	0.11	0.041
17:1 n9	0.62 ^a	0.59 ^a	0.40 ^b	0.45 ^b	0.04	0.001
17-anteiso	0.60 ^a	0.53 ^{ab}	0.40 ^b	0.47 ^{ab}	0.04	0.007
17-iso	0.47 ^a	0.38 ^{ab}	0.29 ^b	0.34 ^{ab}	0.01	0.045
18:0	17.17 ^{ab}	15.22 ^b	19.05 ^a	18.66 ^a	0.83	0.012
18-iso	0.17 ^a	0.10 ^b	0.08 ^b	0.10 ^b	0.01	0.001
18:1 n9	34.69 ^{ab}	30.16 ^c	36.16 ^a	33.20 ^{bc}	0.75	0.001
18:1 n11	1.00 ^c	1.45 ^a	1.24 ^b	1.28 ^b	0.52	0.001
18:1 n12	0.32 ^b	0.64 ^a	0.25 ^b	0.41 ^b	0.05	0.001
18:1 n13	0.11 ^b	0.24 ^a	0.09 ^b	0.16 ^{ab}	0.03	0.011
18:1 n14	0.07 ^b	0.15 ^a	0.05 ^b	0.09 ^b	0.01	0.001
18:1 n15	0.26 ^c	0.68 ^a	0.23 ^c	0.48 ^b	0.05	0.001
18:1 n4	0.01 ^b	0.02 ^{ab}	0.04 ^a	0.03 ^a	0.01	0.001
18:1 n5	0.02 ^b	0.04 ^{ab}	0.07 ^a	0.06 ^a	0.01	0.001
18:1 n6-8	0.40 ^b	0.67 ^a	0.66 ^a	0.62 ^a	0.06	0.013
18:1 n9	0.29 ^b	0.44 ^a	0.37 ^a	0.38 ^a	0.03	0.004
18:1 n10	2.58 ^b	4.81 ^a	3.44 ^{ab}	3.75 ^{ab}	0.13	0.001
18:1 n11	1.02 ^b	2.39 ^a	1.52 ^{ab}	1.73 ^{ab}	0.15	0.001
18:1 n12	0.40 ^b	1.07 ^a	1.03 ^a	0.96 ^a	0.06	0.001
18:1 n15	0.13 ^c	0.71 ^a	0.18 ^c	0.49 ^b	0.02	0.001
18:1 n16	0.22 ^c	0.77 ^a	0.22 ^c	0.57 ^b	0.04	0.001
18:2 n9, n11	0.52 ^b	0.89 ^a	0.51 ^b	0.64 ^{ab}	0.05	0.001
Other 18:2 conjugated	0.06 ^b	0.14 ^a	0.08 ^b	0.11 ^a	0.01	0.004
18:2 n11, n15	0.17 ^c	1.92 ^a	0.12 ^c	0.93 ^b	0.06	0.001
18:2 n-6	3.25	3.32	3.38	3.41	0.20	0.565
Other 18:2 non-conjugated	0.34 ^b	0.72 ^a	0.27 ^b	0.49 ^{ab}	0.04	0.001
18:3 n-3	0.43 ^b	1.34 ^a	0.46 ^b	1.18 ^a	0.05	0.001
18:3 n9, n11, n15	0.01 ^c	0.06 ^a	0.01 ^c	0.03 ^b	0.01	0.001
20:0	0.16	0.16	0.17	0.17	0.01	0.879
20:1 n9	0.01	0.01	0.01	0.02	0.01	0.379
20:2 n-6	0.02	0.01	0.02	0.01	0.01	0.146
20:3 n-3	0.00 ^b	0.02 ^a	0.00 ^b	0.01 ^a	0.01	0.001
20:3 n-6	0.02	0.01	0.01	0.03	0.01	0.114
20:4 n-6	0.10	0.07	0.07	0.06	0.01	0.678
20:5 n-3	0.01	0.02	0.02	0.03	0.01	0.534
21:0	0.01	0.02	0.01	0.01	0.01	0.857
22:0	0.01	0.01	0.01	0.02	0.01	0.600
22:4 n-6	0.02 ^a	0.00 ^b	0.02 ^a	0.00 ^b	0.01	0.024
22:5 n-3	0.09	0.08	0.08	0.08	0.01	0.213
22:6 n-3	0.02	0.02	0.02	0.02	0.01	0.854
Other fatty acids	0.86	0.74	0.67	0.81	0.02	0.564

C = containing neither linseed nor olive cake; L = containing 20% of rolled linseed; OC = containing 35% of stoned olive cake; OCL = containing 10% of rolled linseed and 17% of stoned olive cake.

^{a-c} Within a row, values with different letters are significantly different ($P \leq 0.05$).

¹ The dietary treatments consisted of grass hay and the experimental concentrates at a 20 : 80 forage : concentrate ratio.

when the daily intake exceeds 500 mg (Pintus *et al.*, 2013). As the CLA selectivity is higher for neutral lipids than for polar lipids (Jerónimo *et al.*, 2011), the overall content of intramuscular fat may significantly affect the total content of CLA. For instance, Berthelot *et al.* (2010) reported very low levels (<0.2/100 g of fatty acids) of CLA in the muscle fat in response to a concentrate-based diet supplemented with 10% of extruded linseed, as a consequence of the very lean muscles obtained in that trial (<2/100 g of muscle). Nevertheless, previous studies on the fatty-acid composition of suckling lambs reported very high levels of CLA in the intramuscular fat from very lean carcasses (Serra *et al.*, 2009a, 2009b), which highlights that the incorporation of CLA in the lipid fraction may also be affected by individual, breed and physiological factors.

The level of SFA was poorly affected by the dietary treatment. In particular, the C diet resulted in a higher level of 16:0, whereas the OC and OCL diets, which included olive oil, resulted in a higher content of 18:0. Meat from the lambs in the L group tended to have the lower content of SFA, in particular 16:0. This pattern was particularly appreciable in the neutral lipid fraction (Table 5), where the SFAs are selectively esterified (Jerónimo *et al.*, 2011).

The branched-chain fatty acids (BCFA) originate from the metabolism of branched-chain amino acids in the rumen and their content in milk fat has been proposed as a marker of rumen metabolism (especially for volatile fatty acid and methane production) and as predictor of rumen acidosis (Fievez *et al.*, 2012). Some rumen microorganisms (especially fibrolytic bacteria) are sensitive to the presence of unsaturated fatty acids and their growth may be significantly inhibited by diets enriched with vegetable or fish oil (Shingfield *et al.*, 2013). In the present study, meat from the L or OC lambs contained significantly lower amounts of C16 iso, C17 iso, C17 anteiso and C18 iso than meat from the C lambs (Table 3), probably as an effect of the higher level of unsaturated fatty acids in the L and OC diet. This effect is more evident in neutral lipids, where BCFA were preferentially incorporated, than in polar lipids (Tables 4 and 5). However, the content of BCFA in meat from the OC lambs was lower than that in meat from the L lambs ($P < 0.01$), whereas the level of BCFA in meat from the OCL lambs was intermediate (Tables 4 and 5). This pattern could be because of an additive effect of polyphenols and olive oil contained in the cake. Previous *in vitro* and *in vivo* studies, in fact, demonstrated a significant effect of dietary polyphenols on the rumen bacterial community, which resulted in a decrease of BCFA content in both rumen liquor and intramuscular fat (Vasta *et al.*, 2009; Buccioni *et al.*, 2011).

Conclusions

The use of rolled linseed at 20% concentrate DM allowed to modify the fatty-acid composition of intramuscular lipids towards a healthier profile, by increasing the unsaturation degree of both membrane lipids and NL. In particular, the

ratio n-6:n-3 reached values below 3, which is considered an optimal value for human nutrition. This positive effect was also observed, to a lower extent, when rolled linseed was included at 10% of concentrate DM in combination with stoned olive cake. However, the level enrichment of 18:3 n-3 obtained in the intramuscular fat from the OCL lambs was higher than expected, considering the level of rolled linseed in the diet (when compared with the OC and L treatments), suggesting a putative synergic effect of stoned olive cake with linseed. Further studies are needed to assess the role of olive polyphenols in the C18:3 n-3 biohydrogenation. The use of high amount of stoned olive cake in the concentrate (35%) did not affect the growing performances of lambs and only marginal effects were observed on the fatty-acid composition of intramuscular fat, as a consequence of the intense biohydrogenation of oleic acid contained in the olive cake. In conclusion, the use of rolled linseed to improve the fatty-acid profile of intramuscular fat of lambs may be combined with stoned olive to decrease the feeding costs, without any detrimental effect on growing performances.

Acknowledgements

Research was funded by the University of Pisa under the research project: Progetto di Ricerca di Ateneo 'Improvement of nutritional and dietetic properties of beef and lamb meat: an interdisciplinary approach'.

Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S1751731113001924>.

References

- Alves SP and Bessa RJB 2007. Identification of *cis*-12, *cis*-15 octadecadienoic acid and other minor polyenoic fatty acids in ruminant fat. *European Journal of Lipid Science and Technology* 109, 879–883.
- Association of Official Analytical Chemists 1990. Official method of analysis, vol. 2, 15th edition. AOAC, Arlington, VA, USA.
- Bas P, Berthelot V, Pottier E and Normand J 2007. Effect of level of linseed on fatty acid composition of muscles and adipose tissues of lambs with emphasis on *trans* fatty acids. *Meat Science* 77, 678–688.
- Berthelot V, Bas P and Schmidely P 2010. Utilization of extruded linseed to modify fatty composition of intensively-reared lamb meat: effect of associated cereals (wheat vs. corn) and linoleic acid content of the diet. *Meat Science* 84, 114–124.
- Buccioni A, Minieri S, Rapaccini S, Antongiovanni M and Mele M 2011. Effect of chestnut and quebracho tannins on fatty acid profile in rumen liquid- and solid-associated bacteria: an *in vitro* study. *Animal* 5, 1521–1530.
- Buccioni A, Antongiovanni M, Petacchi F, Mele M, Serra A, Secchiari P and Benvenuti D 2006. Effect of dietary fat quality on C18:1 fatty acids and conjugated linoleic acid production: an *in vitro* rumen fermentation study. *Animal Feed Science and Technology* 127, 268–282.
- Cannas A, Tedeschi LO, Fox DG, Pell AN and Van Soest PJ 2004. A mechanistic model for predicting the nutrients requirements and feed biological values for sheep. *Journal of Animal Science* 82, 149–169.
- Christie WW 1982. A simple procedure for rapid transmethylation of glycerolipids and cholesteryl esters. *Journal of Lipid Research* 23, 1072–1075.
- Dal Bosco A, Mourvaki E, Cardinali R, Servili M, Sebastiani B, Ruggeri S, Mattioli S, Taticchi A, Esposto S and Castellini C 2012. Effect of dietary

- supplementation with olive pomaces on the performance and meat quality of growing rabbits. *Meat Science* 92, 783–788.
- Daniel ZCTR, Wynn RJ, Salter AM and Buttery PJ 2004. Differing effects of forage and concentrate diets on the oleic acid and conjugated linoleic acid content of sheep tissues: the role of stearoyl-CoA desaturase. *Journal of Animal Science* 82, 747–758.
- Doreau M, Laverroux S, Normand J, Chesneau G and Glasser F 2009. Effect of linseed fed as rolled seeds, extruded seeds or oil on fatty acid rumen metabolism and intestinal digestibility in cows. *Lipids* 44, 53–62.
- Fievez V, Colman E, Castro-Montoya JM, Stefanov I and Vlaeminck B 2012. Milk odd- and branched-chain fatty acids as biomarkers of rumen function – an update. *Animal Feed Science and Technology* 172, 51–65.
- Folch J, Lees M and Stanley GHS 1957. A simple method for the isolation and purification of total lipids from animal tissue. *Journal of Biological Chemistry* 226, 497–509.
- Jerónimo E, Alves SP, Alfaia CM, Prates JAM, Santos-Silva J and Bessa RJB 2011. Biohydrogenation intermediates are differentially deposited between polar and neutral intramuscular lipids of lambs. *European Journal of Lipid Science and Technology* 113, 924–934.
- Jerónimo E, Alfaia CMM, Alves SP, Dentinho MTP, Prates JAM, Vasta V, Santos-Silva J and Bessa RJB 2012. Effect of dietary grape seed extract and *Cistus ladanifer* L. in combination with vegetable oil supplementation on lamb meat quality. *Meat Science* 92, 841–847.
- Juaneda P and Rocquelin G 1985. Rapid and convenient separation of phospholipids and non-phosphorus lipids from rat heart using silica cartridges. *Lipids* 20, 40–41.
- Kramer JK, Hernandez M, Hernandez CC, Kraft J and Dugan ME 2008. Combining results of two GC separations partly achieves determination of all *cis* and *trans* 16:1, 18:1, 18:2 and 18:3 except CLA isomers of milk fat as demonstrated using Ag-ion SPE fractionation. *Lipids* 43, 259–273.
- Luciano G, Pauselli M, Servili M, Mourvaki E, Serra A, Monahan FJ, Lanza M, Priolo A, Zinnai A and Mele M 2013. Dietary olive cake reduces the oxidation of lipids, including cholesterol, in lamb meat enriched in polyunsaturated fatty acids. *Meat Science* 93, 703–714.
- McAfee AJ, McSorley EM, Cuskelly GJ, Moss BW, Wallace JMW, Bonham MP and Fearon AM 2010. Red meat consumption: an overview of the risks and benefits. *Meat Science* 84, 1–13.
- Mele M and Banni S 2010. Lipid supplementation in small ruminant nutrition and dairy products quality: implications for human nutrition. In *Energy and protein metabolism and nutrition*. EAAP Publication 127 (ed. GM, Crovetto), pp. 653–663. Wageningen Academic Publishers, Wageningen, The Netherlands.
- Mioc B, Pavic V, Vnucac I, Prpic Z, Kostelic A and Susic V 2007. Effect of olive cake on daily gain, carcass characteristics and chemical composition of lamb meat. *Czech Journal of Animal Science* 52, 31–36.
- Molina-Alcaide E and Yáñez-Ruiz DR 2008. Potential use of olive by-products in ruminant feeding: a review. *Animal Feed Science and Technology* 147, 247–264.
- Moloney AP, Kennedy C, Noci F, Monahan FJ and Kerry JP 2012. Lipid and colour stability of *M. longissimus* muscle from lambs fed camelina or linseed as oil or seeds. *Meat Science* 92, 1–7.
- Mosley E, Powell GL, Riley MB and Jenkins TC 2002. Microbial biohydrogenation of oleic acid to *trans* isomers in vitro. *Journal of Lipid Research* 43, 290–296.
- Noci F, Monahan FJ and Moloney AP 2011. The fatty acid profile of muscle and adipose tissue of lambs fed camelina or linseed as oil or seeds. *Animal* 5, 134–147.
- Palmquist DL, St-Pierre N and McClure KE 2004. Tissue fatty acid profiles can be used to quantify endogenous rumenic acid synthesis in lambs. *Journal of Nutrition* 134, 2407–2414.
- Pintus S, Murru E, Carta G, Cordeddu L, Batetta B, Accossu S, Pistis D, Uda S, Elena Ghiani M, Mele M, Secchiari P, Almerighi G, Pintus P and Banni S 2013. Sheep cheese naturally enriched in α -linolenic, conjugated linoleic and vaccenic acids improves the lipid profile and reduces anandamide in the plasma of hypercholesterolaemic subjects. *British Journal of Nutrition* 109, 1453–1462.
- Sadeghi H, Yansari AT and Ansari-Pirsarai Z 2009. Effects of different olive cake by products on dry matter intake, nutrient digestibility and performance of Zel sheep. *International Journal of Agriculture and Biology* 11, 39–43.
- SAS 1999. User's guide: statistics, version 8.0 edition. SAS Inst. Inc., Cary, NC, USA.
- Scollan N, Hocquette JF, Nuernberg K, Dannenberger D, Richardson I and Moloney A 2006. Innovations in beef production systems that enhance the nutritional and health value of beef lipids and their relationship with meat quality. *Meat Science* 74, 17–33.
- Serra A, Mele M, La Comba F, Conte G, Buccioni A and Secchiari P 2009a. Conjugated linoleic acid (CLA) content of meat from three muscles of Massese suckling lambs slaughtered at different weights. *Meat Science* 81, 396–404.
- Serra A, Macciotta NPP, Mele M, Nudda A, Conte G and Secchiari P 2009b. Effect of weight of slaughter and feeding regimen on conjugated linoleic acid and *trans* fatty acid content in lamb meat: a meta-analysis approach. *Italian Journal of Animal Science* 8 (suppl. 2), 540–542.
- Servili M, Esposito S, Taticchi A, Urbani S, Selvaggini R, Di Maio I and Veneziani G 2011. Innovation in extraction technology for improved virgin olive oil quality and by-product valorisation. *Acta Horticulturae* 888, 303–316.
- Shingfield KJ, Bonnet M and Scollan ND 2013. Recent developments in altering the fatty acid composition of ruminant-derived foods. *Animal* 7 (suppl. 1), 132–162.
- Sinclair LA 2007. Nutritional manipulation of the fatty acid composition of sheep meat: a review. *The Journal of Agricultural Science* 145, 419–434.
- Smith BK, Robinson LE, Nam R and Ma DWL 2009. *Trans*-fatty acids and cancer: a mini-review. *British Journal of Nutrition* 102, 1254–1266.
- Van Soest PJ, Robertson JB and Lewis BA 1991. Methods for dietary fiber, neutral detergent fiber, and non starch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* 74, 3583–3597.
- Vasta V and Luciano G 2011. The effect of dietary consumption of plants secondary compounds on small ruminants' products quality. *Small Ruminant Research* 101, 150–159.
- Vasta V, Mele M, Serra A, Scerra M, Luciano G, Lanza M and Priolo A 2009. Metabolic fate of fatty acids involved in ruminal biohydrogenation in sheep fed concentrate or herbage with or without tannins. *Journal of Animal Science* 87, 2674–2684.
- Vera R, Aguilar C, Lira R, Toro P, Barrales L, Pena I, Squella F, Pérez P, Quenaya J, Yutronic H and Briones I 2009. Feeding dry olive cake modifies subcutaneous fat composition in lambs, noting cake resistance to degradation and peroxidation. *Chilean Journal of Agricultural Research* 69, 548–559.
- World Health Organization 2003. Diet, nutrition and the prevention of chronic diseases. Report of a joint WHO/FAO expert consultation. WHO technical report series 916. WHO, Geneva, Suisse.