Endocrine

The 2039 A/G FSH receptor gene polymorphism influences glucose metabolism in healthy men --Manuscript Draft--

Manuscript Number:	ENDO-D-20-00430R2
Full Title:	The 2039 A/G FSH receptor gene polymorphism influences glucose metabolism in healthy men
Article Type:	Original Article
Corresponding Author:	Rossella Cannarella, M.D. University of Catania Catania, ITALY
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	University of Catania
Corresponding Author's Secondary Institution:	
First Author:	Rossella Cannarella, M.D.
First Author Secondary Information:	
Order of Authors:	Rossella Cannarella, M.D.
	Nicolò Musso, Ph.D.
	Rosita A. Condorelli, M.D., Ph.D.
	Marco Musmeci, M.D.
	Stefania Stefani
	Antonio Aversa
	Sandro La Vignera, M.D., Ph.D.
	Aldo E. Calogero, M.D.
Order of Authors Secondary Information:	
Funding Information:	
Abstract:	OBJECTIVE. To assess the role of c. 2039 A/G (p. Asp680Ser) (rs 6166) and c29 G/A (rs 1394205) follicle-stimulating hormone receptor (FSHR) gene single nucleotide polymorphisms (SNPs) in a cohort of healthy men. METHODS. One-hundred twenty-seven healthy men underwent evaluation of the anthropometric parameters, assessment of metabolic and lipid profile, measurement FSH serum levels and genotyping of both the aforementioned FSHR SNPs. Data grouped according to the FSHR rs6166 or rs1394205 genotypes underwent to statistical analysis. MAIN RESULTS. The three groups of men for each FSHR SNP did not differ statistically significantly for BMI and serum FSH levels. As for FSHR rs6166 SNP, glucose levels were significantly lower in men with the GG genotype compared to those with the AA genotype. Men with AG had lower insulin levels and HOMA index values compared with those carrying the genotype AA (p<0.05). The GG group showed a negative correlation between serum FSH levels and HOMA index (p<0.05). In contrast, men grouped according to the FSHR rs1394205 genotype showed no significant difference in blood glucose, serum insulin levels and HOMA index. The AG group showed a negative correlation between serum FSH levels and HOMA index (p<0.05). CONCLUSION. Men with the genotype GG of the FSHR rs6166 SNP have lower blood glucose levels than those with the AA genotype. Their FSH levels inversely correlated with insulin and HOMA index. In contrast, the genotype FSHR rs6166 A/G did not reveal any role of FSH on glucose metabolism in healthy men. The inverse

	relationship between FSH and insulin or HOMA index in the group with the genotype GG of the FSHR rs6166 SNP suggests a possible cross-talk between FSH and insulin.
Response to Reviewers:	Manuscript ID ENDO-D-20-00430 Comment 1: Reviewer #2: The responses to comments made by the reviewer have been done for the main part. Anyway, I still suggest that Figure 1 is unnecessary and the data should be included in Table 1. Answer to comment 1: Done. Thank you. Comment 2: Response to question 11 (see above) is not contested in the text: The authors say that thy added the following comment in the conclusion (please see lines 212-213). "Molecular mechanisms underlying the interaction between FSH and insulin are still matter of debate. Unpublished data which we are working on, support that incubation with insulin down-regulates FSH receptor in an in-vitro culture of porcine Sertoli cells. The in-vivo findings of a recently published study (ref. 29) may somehow reflect this molecular mechanism. With this in mind, the inverse relationship between FSH and insulin and between FSH and HOMA-index in the GG group (encoding for a more resistant FSHR to signal transduction) could confirm the occurrence of a molecular interaction between insulin and the FSHR. Clearly, this topic represents a black hole and deserves further investigation.
	But this paragraph is not found in the revised version. Answer to comment 2: Added. Thank you.

Click here to view linked References

1	The 2039 A/G FSH receptor gene polymorphism influences glucose metabolism in healthy men
₁ 2	
2 3 3	
4 5 4	Short title: FSHR polymorphisms and glycaemia
б	Short title. I Srik polymorphisms and grycaenna
⁷ 5 ⁹ 6	
9 10 6	
11 12 7	Rossella Cannarella ¹ , Nicolò Musso ² , Rosita A. Condorelli ¹ , Marco Musmeci ¹ , Stefania Stefani ² , Antonio
13 14 8	Aversa ³ , Sandro La Vignera ¹ , Aldo E. Calogero ¹
15	Aversa, Sandro La Vignera, Aldo E. Calogero
16 17 9	
18 19 10	
20 21 11	
22 23 12	¹ Department of Clinical and Experimental Medicine, University of Catania, Catania, Italy
24	
²⁵ 13 26	² Bio-nanotech Research and Innovation Tower (BRIT), University of Catania, Catania, Italy
²⁷ 28 ¹ 4	³ Department of Experimental and Clinical Medicine, "Magna Graecia" University, Catanzaro, Italy
29 30 15	
31 3 216	
33	
³⁴ 17 35	
³⁶ 18	
38 39 19	For submission to: Endocrine
40 41 20	
42 4 321	
44	
45 22 46	Summary characters count: 287
47 48 23	Number of figures: 2
49 50 24	Number of tables: 2
51 52 25	
53 54 1 6	
54 26 55 56	
56 57 27 58 28	Corresponding author: Prof. Aldo E. Calogero, Department of Clinical and Experimental Medicine, University of Catania, via S. Sofia 78, 95123 Catania, Italy. FAX ++39-95-3781180; Email: <u>acaloger@unict.it</u>
59	
60 61	
62 63	1
64 65	

37 17

²⁵41 26

28**42**

³⁴45 35

³⁶46 37

39**47**

46

48<mark>51</mark>

₁30 **OBJECTIVE.** To assess the role of c. 2039 A/G (p. Asp680Ser) (rs 6166) and c. -29 G/A (rs 1394205) follicle-stimulating hormone receptor (FSHR) gene single nucleotide polymorphisms (SNPs) in a cohort of healthy ⁵32 6 ⁷33 ⁹34 men.

METHODS. One-hundred twenty-seven healthy men underwent evaluation of the anthropometric parameters, assessment of metabolic and lipid profile, measurement FSH serum levels and genotyping of both the aforementioned FSHR SNPs. Data grouped according to the FSHR rs6166 or rs1394205 genotypes underwent to statistical analysis.

MAIN RESULTS. The three groups of men for each FSHR SNP did not differ statistically significantly for body mass index and serum FSH levels. As for FSHR rs6166 SNP, glucose levels were significantly lower in men with the GG genotype compared to those with the AA genotype. Men with AG had lower insulin levels and HOMA index values compared with those carrying the genotype AA (p<0.05). The GG group showed a negative correlation between serum FSH levels and insulin and between serum FSH levels and HOMA index (p<0.05). In contrast, men grouped according to the FSHR rs1394205 genotype showed no significant difference in blood glucose, serum insulin levels and HOMA index. The AG group showed a negative correlation between FSH insulin and between serum FSH levels and HOMA index (p<0.05).

CONCLUSION. Men with the genotype GG of the FSHR rs6166 SNP have lower blood glucose levels than those with the AA genotype. Their FSH levels inversely correlated with insulin and HOMA index. In contrast, the genotype FSHR rs6166 A/G did not reveal any role of FSH on glucose metabolism in healthy men. The inverse relationship between FSH and insulin or HOMA index in the group with the genotype GG of the FSHR rs6166 SNP suggests a possible cross-talk between FSH and insulin.

Keywords: FSH; 2039 A/G FSHR; FSHR polymorphism; glucose; metabolism; insulin; insulin resistance

54 Introduction

The follicle-stimulating hormone (FSH) is a glycoprotein made of an α and a β subunit. It exerts its peripheral effects by binding to its receptor (FSHR), a 678 amino acid protein belonging to the G-coupled receptor family, expressed in granulosa and Sertoli cells, where FSH stimulates follicle genesis and spermatogenesis, respectively [1].

The *FSHR*, mapping in the 2p16.3 chromosome, contains 10 exons and 9 introns. The first nine exons encode for the majority of the extracellular domain. The exon 10 encodes for the C-terminal region of the extracellular domain, the trans-membrane and the intracellular domains [1]. The HapMap database (http://hapmap.ncbi.nlm.nih.gov) lists more than 900 *FSHR* single nucleotide polymorphisms (SNPs). A large body of evidence has been released on the c. 2039 A/G (p. Asp680Ser) (rs 6166) and the c. -29 G/A (rs 1394205) *FSHR* SNPs so far. The first maps in the exon 10, which encodes for the transmembrane domain of the FSHR, and it is known to influence the efficiency of signal transduction. In detail, the c. 2039 A/G genotype impacts on the expression of the amino acid 680 and the FSHR Ser680Ser (GG) results to be more resistant to the FSH signal compared with the Asp680Asp (AA) [2]. The c. -29 G/A SNP maps in the *FSHR* gene promoter and influences *FSHR* expression. Particularly, the A allele is associated with a lower (by 56%) transcription activity of the promoter [3].

Classically, the effects of FSH have been restricted to gonads and a large body of evidence of *FSHR* SNPs has been published on fertility in both genders [4-6]. However, recent data point to the existence of extra-gonadal effects of FSH. Accordingly, the *FSHR* is expressed in human bone cells and in human bone marrow-adipocytes [7,8]. Furthermore, FSH has been already addressed as possibly implicated in the post-menopausal changes of bone mineral density (BMD) [7,9] and in lipid distribution, adiposity and metabolism in male patients with hypergonadotropic hypogonadism [8].

To further explore the role that FSH may play in glucose and lipid metabolism in the male gender, we investigated the possible influence of the c. 2039 A/G (p. Asp680Ser) (rs 6166) and the c. -29 G/A (rs 1394205) *FSHR* SNPs on blood glucose, insulin, total, HDL and LDL cholesterol, triglycerides, on the homeostasis model assessment (HOMA) index and the body mass index (BMI) in a cohort of healthy men.

80 Subjects and methods

Ethical approval

This study was approved by the Teaching Hospital "G. Rodolico" Ethics Committee (Protocol n. 178/2017/PO) and informed written consent was obtained from each participant after full explanation of the purpose and nature of all procedures used. The study has been carried out in accordance with the principles expressed in the Declaration of Helsinki.

Patient selection

The study has been carried out in male patients referring to the Division of Andrology and Endocrinology, University of Catania, for an endocrinologic or andrologic counselling. A detailed medical history was collected and weight and height were assessed for each man. The BMI was calculated using the formula: weight (Kg)/[height (m) x height (m)]. A total of 146 men were evaluated for inclusion in this study. The following exclusion criteria were used and this resulted in the exclusion of a number of men that is indicated in parentheses: azoospermia (n=3), tumor (n=2), assumption of metformin, insulin, hypoglycemic or hypocholesterolemic drugs (n=5), diabetes (n=2), use of gonadotoxic drugs (n=2), hypogonadism (n=3), testosterone replacement therapy (n=1), FSH therapy (n=1), major adverse cardiovascular events (n=0), major comorbilities (n=0). Taking all this into account, 127 Caucasian men from Eastern Sicily were included in the study.

Hormone evaluation was performed by electro chemiluminescence (Hitachi-Roche equipment, Cobas 6000, Roche Diagnostics, Indianapolis, IN, USA). Reference values were as follows: FSH 0.95-11.95 IU/l, glycemia 74-100 mg/dl, insulin 1.9-23 μ IU/ml, total cholesterol 0-200 mg/dl, HDL cholesterol 35-55 mg/dl, LDL cholesterol <160 mg/dl, triglycerides 35-200 mg/dl. The homeostasis model assessment (HOMA) index was evaluated using the formula: [glycemia (mg/dl) x insulin (μ IU/ml)]/405. The upper normal value of HOMA index was 2.5.

FSHR analysis

Genomic DNA was extracted from blood cells using the PureLink® Genomic DNA Kits (invitrogen Catalog Numbers K1821-04) for purification of genomic DNA according to the manufacturer's instructions. The concentration and the quality of the DNA was determined using a ND-1000 spectrophotometer (NanoDrop, Thermo Scientific, USA). Allelic Discrimination was performed with TaqMan assay in order to show a different genotyping distribution of *FSHR* polymorphism. Probes and primers for rs 6166 *FSHR* and

rs 1394205 *FSHR* polymorphisms were chosen on https://www.thermofisher.com/it/en/home/life science/pcr/real-time-pcr/real-time-pcr-assays/snp-genotyping-taqman-assays.html?SID=fr-taqman-2. The
 reaction was carried out according to manufacturer's instructions (cod 4371355, Applied Biosystems, CA,
 USA). Each DNA sample was analyzed in triplicate [10]. Allelic Discrimination real-time PCR analysis was
 performed using LightCycler® 480 System (Roche Molecular Systems, Inc).

Statistical analysis

The normality of the variables was evaluated with the Shapiro-Wilk test. Descriptive statistical results have been reported as mean±standard deviation (SD) for not-skewed variables. Statistical analysis was performed by one-way analysis of variance (ANOVA), followed by the Duncan's Multiple Range Test, using SPSS 22.0 for Windows (SPSS Inc., Chicago, IL, USA). For correlation analysis, the Pearson or the Kendall tests were used, according to the data distribution. A p value less than 0.05 was accepted as statistically significant.

Results

The number and the prevalence (in parentheses) of *FSHR* rs6166 genotypes were: 33 (26.0%), 63 (49.6%) and 31 (24.4%) for AA, AG and GG, respectively. For the genotype *FSHR* rs1394205, they were: 14 (11.0%), 38 (29.9%) and 75 (59.1%) for AA, AG and GG genotypes, respectively. The *FSHR* allelic frequencies and the estimated *FSHR* haplotype frequencies found in this cohort were consistent with previously published data from unrelated Caucasian populations [9, 11-13].

Data were normally distributed. Anthropometric data and biochemical parameters of men grouped according to the *FSHR* rs6166 or rs1394205 genotype are shown in **Table 1**. No significant difference in all parameters was observed between men classified according to the different genotype (p>0.05 in all cases), with the exception of total cholesterol, which was higher in men with *FSHR* rs1394205 AA genotype compared to those with the AG genotype (p <0.05). According to their BMI, 42.2% of patients were overweight, 20% obese; 8.3% had hyperglicemia, 20.5% insulin-resistance, and 26.3% hypercholesterolemia. The prevalence of these features did not differ after classification by the *FSHR* genotype.

Serum glucose levels were lower in men with *FSHR* rs6166 GG vs. men with the AA genotype. Men with AG SNP had lower insulin levels and HOMA index values (**Table 1**) compared to AA (p < 0.05). Men

137 grouped for the FSHR rs1394205 genotype did not show any difference in glycaemia, insulin and HOMA

index (Figure 1).

No significant correlation was found between FSH and glycaemia (r= 0.01), insulin (r= -0.26) or HOMA index (r= -0.19) in the entire cohort. Correlation analysis in men grouped by *FSHR* rs6166 or rs1394205 genotypes are shown in **Table 2**. A significant negative correlation between FSH and insulin and FSH and HOMA index was observed in men with the genotype *FSHR* rs6166 GG and for the genotype *FSHR* rs1394205 AG. All the other correlations were not significant.

Discussion

The results of the present study showed that the polymorphism rs6166 in exon 10 of the *FSHR* gene influences significantly blood glucose levels in healthy men. Particularly, the GG genotype, encoding for a less efficiently transducing FSHR protein, is associated with lower glucose levels compared to the AA one, a SNP that more efficiently transduce the FSH signal. Since the groups resulted to have a similar age, BMI and FSH levels, these findings suggest that the *FSHR* gene plays a role on glucose metabolism in men. The significant lower insulin levels and HOMA index in men with the AG genotype compared those with AA suggest a sort of compensatory mechanism to avoid hyperglycemia, which is not observed in men with GG due to the normal glucose levels. Interestingly, an inverse relationship between FSH and insulin levels or HOMA index was found in men with the genotype GG. Furthermore, no effect on such parameters was observed when men were classified according to the rs1394205 genotype, aside a significant indirect correlation between FSH and insulin and a negative trend between FSH and HOMA index in the AG group.

Scanty evidence has been published on the impact of FSH on glucose metabolism so far. A preclinical study reported that female mice FSHR gene deficient (*Fshr+/–* mice) do not develop obesity as wild type ovariectomized mice do. These effects were also reported in male *Fshr+/–* mice. Furthermore, the administration of a blocking antibody against FSH was found to efficiently induce thermogenic adipose tissue and to reduce body fat in both sexes [14]. This indicates that FSH may be involved in the development of obesity. In addition, the FSHR has been found expressed in pancreatic β cells in animals but its function in this district is still unclear [15].

A recent clinical study investigated whether FSH may impact on glucose and lipid metabolism in a large number of patients (more then 300) with hypergonadotropic hypogonadism, including patients with

Klinefelter syndrome. Remarkably, the authors reported that FSH could enhance soluble-RANKL (sRANKL) secretion from human adipocytes. sRANKL levels, in turn, are negatively associated with fat percentage, fasting insulin and glucose [8]. sRANKL has been shown to decrease pancreatic β cell proliferation and to decrease glucose tolerance [16]. Accordingly, the administration of the sRANKL inhibitor denosumb improves glucose tolerance, but not HbA1c levels in non-diabetic women [17,18]. Thus, the association between FSH and glucose metabolism may be indirect being mediated by the FSH-induced sRANKL production.

The inverse relationship between FSH and insulin or HOMA index observed in men with the GG genotype of the FSHR rs6166 SNP suggests the hypothesis that FSH and insulin may share a common molecular signaling. Generally, tyrosine kinase receptors (e.g. insulin receptor or insulin-like growth factor receptor) are able to influence the FSHR signaling [19,20,21]. In this regard, the insulin receptor substrate 1 (IRS1), which represents a co-factors linked to tyrosine kinase receptors, seems to play a role in the pathogenesis of insulin-resistance [22,23]. Interestingly, it has been described as the hub that links FSH signaling pathway and the tyrosine kinase receptor-dependent cascade [24,25]; thus relating FSHR to the insulin receptor. The possible cross-talk between FSH and insulin has been further investigated in an *in-vitro* study on porcine pre-pubertal Sertoli cells aimed to evaluate the secretion of anti-Müllerian hormone (AMH) and inhibin in response to FSH and insulin. The results showed that insulin significantly decreased FSHstimulated AMH and inhibin secretion, suggesting that insulin may likely interfere with the FSH signaling [26]. In line with this evidence, a recent clinical study reported that insulin is able to decrease the response of granulosa cells to FSH stimulation in women with obesity-related infertility during *in-vitro* fertilization [27]. Unfortunately, clinical evidence is very limited in men, due to the small duration of FSH stimulation of the poorly standardized protocols used for the treatment of infertile male patients [28]. FSH administration to patients with oligozoospermia, normal serum FSH levels and insulin-resistance showed to increase its efficacy on sperm parameters and fertility outcome when it was co-administered with metformin [29]. These findings add further evidence on the possible interference of insulin in FSH signaling also in men. Conversely, FSH may somehow influence insulin pathway and glucose metabolism, which may be dimly supported by the significantly increase of FSH serum levels in diabetic patients, as results of meta-analysis have shown [30]. In addition, an inverse correlation between FSH and insulin, and FSH and HOMA index were found in the rs1394205 AG group. The reason of this result is hardly to be interpreted. This strengthen the importance of fully addressing the molecular relationship between FSHR and insulin.

7

Finally, the results of the present study expand the view of the possible clinical applications of *FSHR* SNPs assessment. A possible *FSHR* SNPs usefulness has been classically searched on fertility so far. However, the research promoted after the identification of FSH effects on bone tissue has led to studies aimed at exploring its protective role against the development of osteoporosis in women with the *FSHR* rs6166 GG genotype; thus supporting extra-gonadal effects that FSH may likely have [9]. Similarly, the importance of *FSHR* SNPs in the metabolic field needs to be further addressed.

Our results need to be taken with care due to the relatively small sample size examined. Hence, further studies on greater cohorts are warranted.

In conclusion, we found significantly lower blood glucose levels in men with the GG genotype of the FSHR rs6166 SNP, and significantly lower insulin levels and HOMA index in AG compared with AA men. Furthermore, FSH serum levels correlated inversely with insulin levels and the HOMA index in men with GG, No effects were observed on these parameters when men were classified according to the rs1394205 genotype, although a significant indirect correlation between FSH and insulin and a negative trend between FSH and HOMA index was found in the AG group. FSHR rs6166 A/G genotypization could reveal the impact of FSH on glucose metabolism in healthy men. The inverse relationship of FSH with insulin and the HOMA index in the FSHR rs6166 GG group may support the existence of a cross-talk between FSH and insulin. Accordingly, the higher is insulin, the lower is FSH, thus reflecting the importance of treating insulin-resistance prior to FSH administration in oligozoospermic patients [29]. Molecular mechanisms underlying the interaction between FSH and insulin are still matter of debate. Unpublished data support that incubation with insulin down-regulates FSH receptor in an in-vitro culture of porcine Sertoli cells. The in-vivo findings of a recently published study [29] may somehow reflect this molecular mechanism. With this in mind, the inverse relationship between FSH and insulin and between FSH and HOMA-index in the GG group (encoding for a more resistant FSHR to signal transduction) could confirm the occurrence of a molecular interaction between insulin and the FSHR. Clearly, this topic represents a black hole and deserves further investigation.

Compliance with Ethical Standards

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of interest: Rossella Cannarella declares no conflict of interest. Nicolò Musso declares no conflict of interest. Rosita A. Condorelli declares no conflict of interest. Marco Musmeci declares no conflict of interest. Stefania Stefani declares declares no conflict of interest. Antonio Aversa declares no conflict of interest. La Vignera declares no conflict of interest. Aldo E. Calogero declares no conflict of interest.

Ethical approval: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent: Informed consent was obtained from all individual participants included in the study.

Acknowledgements: None.

References

- Gromoll, J., Pekel, E., Nieschlag, E.: The structure and organization of the human follicle-stimulating hormone receptor (FSHR) gene. Genomics. 35, 308-311 (1996)
- Casarini, L., Moriondo, V., Marino, M., Adversi, F., Capodanno, F., Grisolia, C., La Marca, A., La Sala, G.B., Simoni, M.: FSHR polymorphism p.N680S mediates different responses to FSH in vitro. Mol. Cell. Endocrinol. 393, 83-91 (2014)
- Nakayama, T., Kuroi, N., Sano, M., Tabara, Y., Katsuya, T., Ogihara, T., Makita, Y., Hata, A., Yamada, M., Takahashi, N., Hirawa, N., Umemura, S., Miki, T., Soma, M.: Mutation of the folliclestimulating hormone receptor gene 5'-untranslated region associated with female hypertension. Hypertension. 48, 512-518 (2006)
- 4. Ferlin, A., Vinanzi, C., Selice, R., Garolla, A., Frigo, A.C., Foresta, C.: Toward a pharmacogenetic approach to male infertility: polymorphism of follicle-stimulating hormone b-subunit promoter. Fert.
 & Steril. 96, 1344-1349 (2011)
- Tüttelmann, F., Laan, M., Grigorova, M., Punab, M., Sõber, S., Gromoll, J.: Combined effects of the variants FSHB -211G>T and FSHR 2039A>G on male reproductive parameters. J. Clin. Endocrinol. Metab. 97(10), 3639-3647 (2012)
 - Simoni, M., Casarini, L.: Mechanisms in endocrinology: Genetics of FSH action: a 2014-and-beyond view. Eur. J. Endocrinol. 170(3), 91-107 (2014)

252	7.	Sun, L., Peng, Y., Sharrow, A.C., Iqbal, J., Zhang, Z., Papachristou, D.J., Zaidi, S., Zhu, L.L.,
253		Yaroslavskiy, B.B., Zhou, H., Zallone, A., Sairam, M.R., Kumar, T.R., Bo, W., Braun, J., Cardoso-
2 254 4		Landa, L., Schaffler, M.B., Moonga, B.S., Blair, H.C., Zaidi, M.: FSH directly regulates bone mass.
2555 6		Cell 125, 247–260 (2006)
2/56	8.	Juel Mortensen, L., Lorenzen, M., Jørgensen, N., Andersson, A.M., Nielsen, J.E., Petersen, L.I.,
9 1 2 57		Lanske, B., Juul, A., Hansen, J.B., Blomberg Jensen, M.: Possible link between FSH and RANKL
11 1 2258 13		release from adipocytes in men with impaired gonadal function including Klinefelter syndrome. Bone
13 1 2159 15		123, 103-114 (2019)
1260 17	9.	Rendina, D., Gianfrancesco, F., De Filippo, G., Merlotti, D., Esposito, T., Mingione, A., Nuti, R.,
18 1 261		Strazzullo, P., Mossetti, G., Gennari, L.: FSHR gene polymorphisms influence bone mineral density
20 2 2<u>1</u>62		and bone turnover in postmenopausal women. Eur. J. Endocrinol. 163, 165-172 (2010)
22 2 263 24	10.	Barresi, V., Signorelli, S.S., Musso, N., Anzaldi, M., Fiore, V., Alberghina, M., Condorelli, D.F.
² 264		ICAM-1 and SRD5A1 gene polymorphisms in symptomatic peripheral artery disease. Vascular
27 28 28		Medicine.: 19(3), 175–181 (2014)
29 3 266	11.	Sudo, S., Kudo, M., Wada, S., Sato, O., Hsueh, A.J., Fujimoto, S. Genetic and functional analyses of
31 3 2267 33		polymorphisms in the human FSH receptor gene. Mol. Hum. Reprod. 8(10), 893-899 (2002)
3 268 35	12.	Gromoll J, Simoni M.: Genetic complexity of FSH receptor function. Trends Endocrinol Metab
36 3769		2005;16(8):368-73.
38 3 2 570	13.	Tamburino, L., La Vignera, S., Tomaselli, V., Condorelli, R.A., Cannarella, R., Mongioì, L.M.,
40 4 2171 42		Calogero, A.E.: The $-29G/A$ FSH receptor gene polymorphism is associated with higher FSH and LH
4 2372 44		levels in normozoospermic men. J. Assist. Reprod. Genet. 34(10), 1289-1294 (2017)
45 273 46	14.	Liu, P., Ji, Y., Yuen, T., Rendina-Ruedy, E., De Mambro, V.E., Dhawan, S., Abu-Amer, W., Izadmehr,
47 428 74		S., Zhou, B., Shin, A.C., Latif, R., Thangeswaran, P., Gupta, A., Li, J., Shnayder, V., Robinson, S.T.,
49 5 275 51		Yu, Y.E., Zhang, X., Yang, F., Lu, P., Zhou, Y., Zhu, L.L., Oberlin, D.J., Davies, T.F., Reagan, M.R.,
52276 53		Brown, A., Kumar, T.R., Epstein, S., Iqbal, J., Avadhani, N.G., New, M.I., Molina, H., van Klinken,
5 277 55		J.B., Guo, E.X., Buettner, C., Haider, S., Bian, Z., Sun, L., Rosen, C.J., Zaidi, M.: Blocking FSH
56 5 278		induces thermogenic adipose tissue and reduces body fat. Nature. 546(7656), 107-112. (2017)
58 59		
60 61 62		
63 64		10

279	15.	Zhang, R., Zhang, S., Zhu, X., Zhou, Y., Wu, X.: Follicle-stimulating hormone receptor (FSHR) in
2 <u>4</u> 80		Chinese alligator, Alligator sinensis: molecular characterization, tissue distribution and mRNA
2 2381		expression changes during the female reproductive cycle. Anim. Reprod. Sci. 156, 40-50 (2015)
4 2 582 6	16.	Kondegowda, N.G., Fenutria, R., Pollack, I.R., Orthofer, M., Garcia-Ocaña, A., Penninger, J.M.,
283		Vasavada, R.C.: Osteoprotegerin and Denosumab Stimulate Human Beta Cell Proliferation through
9 1 2 84		Inhibition of the Receptor Activator of NF-κB Ligand Pathway. Cell. Metab. 22(1), 77-85 (2015)
11 1 285	17.	Passeri, E., Benedini, S., Costa, E., Corbetta, S.: A Single 60 mg Dose of Denosumab Might Improve
13 1 2486 15		Hepatic Insulin Sensitivity in Postmenopausal Nondiabetic Severe Osteoporotic Women. Int. J.
13 1287 17		Endocrinol. 2015, 352858 (2015)
18 1 288	18.	Lasco, A., Morabito, N., Basile, G., Atteritano, M., Gaudio, A., Giorgianni, G.M., Morini, E., Faraci,
20 2 2<u>1</u>89		B., Bellone, F., Catalano, A.: Denosumab Inhibition of RANKL and Insulin Resistance in
22 2 2390		Postmenopausal Women with Osteoporosis. Calcif. Tissue Int. 98(2), 123-128 (2016)
24 2 591 26	19.	Cannarella, R., Mattina, T., Condorelli, R.A., Mongioì, L.M., Pandini, G., La Vignera, S., Calogero,
27 2792 28		A.E.: Chromosome 15 structural abnormalities: effect on the IGF1R gene expression and function.
29 3 293		Endocr Connect. 6(7), 528-539 (2017)
31 3 2294	20.	Cannarella, R., Mancuso, F., Condorelli, R.A., Arato, I., Mongioì, L.M., Giacone, F., Lilli, C.,
33 3 295 35		Bellucci, C., La Vignera, S., Calafiore, R., Luca, G., Calogero, A.E. Effects of GH and IGF1 on Basal
3296 3796		and FSH-Modulated Porcine Sertoli Cells In-Vitro. J Clin Med 8(6) (2019). pii: E811. doi:
38 3 297		10.3390/jcm8060811.
40 4 2198	21.	Cannarella, R., Arato, I., Condorelli, R.A., Luca, G., Barbagallo, F., Alamo, A., Bellucci, C., Lilli, C.,
42 4 2999 44		La Vignera, S., Calafiore, R., Mancuso, F., Calogero, A.E. The IGF1 Receptor Is Involved in Follicle-
4500 46		Stimulating Hormone Signaling in Porcine Neonatal Sertoli Cells. J Clin Med 8(5) (2019). pii: E577.
$^{47}_{4301}$		doi: 10.3390/jcm8050577.
49 5 302	22.	Honma, M., Sawada, S., Ueno, Y., Murakami, K., Yamada, T., Gao, J., Kodama, S., Izumi, T.,
51 5 3403		Takahashi, K., Tsukita, S., Uno, K., Imai, J., Kakazu, E., Kondo, Y., Mizuno, K., Kawagishi, N.,
53 5 304 55		Shimosegawa, T., Katagiri, H.: Selective insulin resistance with differential expressions of IRS-1 and
56 5 305		IRS-2 in human NAFLD livers. Int. J. Obes. (Lond) 42(9), 1544-1555 (2018)
58 59		
60 61		
62 63		11

306	23.	Shakeri, H., Khoshi, A., Kaffash Bajestani, M., Farahi, A., Javadzadeh, M.S., Hosseini, Z.,
3 <u>1</u> 07		Mohammadi, R.: Association of IRS1 GLY971ARG gene polymorphism with insulin resistance in
2 308 4		Iranian newly diagnosed diabetic adults. Acta Endocrinol. (Buchar.) 15(3), 317-322 (2019)
3 09 6	24.	Law, N.C., Hunzicker-Dunn, M.E.: Insulin receptor substrate 1, the hub linking follicle-stimulating
3710		hormone to phosphatidylinositol 3-kinase activation. J. Biol. Chem. 291, 4547–4560 (2016)
9 1 311 11	25.	Cannarella, R., Condorelli, R.A., La Vignera, S., Calogero, A.E.: Effects of the insulin-like growth
1 3212 13		factor system on testicular differentiation and function: a review of the literature. Andrology. 6(1), 3-
1 3413 15		9 (2018)
¹ 314 17	26.	Cannarella, R., Arato, I., Condorelli, R.A., Mongioì, L.M., Lilli, C., Bellucci, C., La Vignera, S., Luca,
18 1 315		G., Mancuso, F., Calogero, A.E.: Effects of Insulin on Porcine Neonatal Sertoli Cell Responsiveness
20 2 316 22		to FSH In Vitro. J. Clin. Med. 8(6). (2016)
2 3317 24	27.	Xu, P., Huang, B.Y., Zhan, J.H., Liu, M.T., Fu, Y., Su, Y.Q., Sun, Q.Y., Wang, W.H., Chen, D.J., Liu,
2 518 26		J.Q.: Insulin Reduces Reaction of Follicular Granulosa Cells to FSH Stimulation in Women With
27 28 28		Obesity-Related Infertility During IVF. J. Clin. Endocrinol. Metab. 104(7), 2547-2560 (2019)
29 3 320 31	28.	Cannarella, R., La Vignera, S., Condorelli, R.A., Mongioì, L.M., Calogero, A.E.: FSH dosage effect
3 3221 333		on conventional sperm parameters: a meta-analysis of randomized controlled studies. Asian. J. Androl.
3 322 35		(2019). doi: 10.4103/aja.aja_42_19.
³ 3723	29.	La Vignera, S., Condorelli, R.A., Duca, Y., Cannarella, R., Giacone, F., Calogero, A.E.: FSH treatment
38 3 3 24		for normogonadotropic male infertility: a synergistic role for metformin? Eur. Rev. Med. Pharmacol.
40 4 3125 42		Sci. 23(13), 5994-5998 (2019)
4 326 44	30.	Pergialiotis, V., Prodromidou, A., Frountzas, M., Korou, L.M., Vlachos, G.D., Perrea, D.: Diabetes
4527 46 47 4328		mellitus and functional sperm characteristics: A meta-analysis of observational studies. J. Diab.
47 4 328		Compl. 30(6), 1167-1176 (2016)
49 50		
51 52		
53 54		
55		
55 56 57		
57 58		
50		

- 60 61 62 63 64 65

329	Legend to the Figures
3 <u>3</u> 30	Figure 1. Glycaemia, insulin and HOMA index classified according to the rs1394205 <i>FSHR</i> genotype.
2 3381 4	No significant difference was found in AA, AG and GG groups for glycaemia (Panel A), insulin (Panel B)
582 6 3333	and HOMA index (Panel C).
3333	
9 10	
11 12 13	
14 15	
16 17	
18 19	
20 21	
22 23	
24 25 26	
27 28	
29 30	
31 32	
33 34	
35 36 37	
38 39	
40 41	
42 43	
44 45	
46 47 48	
40 49 50	
51 52	
53 54	
55 56	
57 58	
59 60 61	
62 63	13
64 65	

Table 1. Antropometric data and biochemical parameters in 127 healthy men classified according to rs6166 or rs1394205
 polymorphisms of the FSH receptor gene.


Parameters		rs6166			rs1394205	
	AA	AG	GG	AA	AG	GG
Number	33	63	31	14	38	75
Age (years)	37.0±8.7	36.6±9.2	35.7±7.6	40.4±8.5	36.9±10.9	35.4±7.4
BMI (Kg/m ²)	27.0±4.9	27.3±5.4	26.5±3.8	27.0±3.4	26.6±5.5	27.1±4.9
Glycaemia (mg/dl)	<mark>90.5±7.4</mark>	<mark>86.2±8.6</mark>	82.2±5.5*	87.4±12.7	88.2±7.9	<mark>85.2±7.8</mark>
<mark>Insulin (µIU/ml)</mark>	11.0±4.8	7.3±3.0*	<mark>9.0±1.4</mark>	7.5±2.9	8.1±3.7	<mark>9.4±3.5</mark>
HOMA index	2.5±1.3	1.6±0.7*	1.8±1.0	1.6±0.9	1.8±0.9	2.0±1.0
Cholesterol (mg/dl)	184.7±46.6	179.4±36.6	178.6±30.1	208.5±45.5	174.3±28.6*	184.1±37.7
HDL cholesterol (mg/dl)	46.8±14.2	50.4±13.6	47.0±9.3	52.2±12.8	49.7±10.0	46.9±11.7
LDL cholesterol (mg/dl)	134.4±46.6	115.8±32.1	120.9±31.1	145.3±40.7	115.7±39.8	121.6±32.7
Triglycerides (mg/dl)	88.6±49.3	81.4±64.4	93.1±40.2	165.8±33.7	92.5±57.2	92.2±63.1
FSH (IU/l)	5.0±3.2	4.5±2.9	3.9±2.5	4.0±2.3	4.5±3.0	4.4±2.8

Data are expressed as mean \pm standard deviation for continual variables. FSH, follicle-stimulating hormone; BMI, body mass index; HOMA, homeostasis model assessment. The *p* values were calculated using ANOVA for continuous variables. *p<0.05 vs. AA (ANOVA followed by Duncan Multiple Range Test).

Table 2. Correlation analysis of glycaemia, insulin, and HOMA index with the rs6166 or rs1394205 polymorphisms of the FSH receptor gene, in 127 healthy men.

Parameters		rs6166		rs1394205				
	AA	AG	GG	AA	AG	GG		
FSH vs. Glycaemia	r= -0.29	r= 0.07	r= 0.12	r= 0.11	r= -0.20	r=0.08		
FSH vs. Insulin	r= -0.02	r= -0.16	r= -0.69*	r= -0.31	r= -0.67*	r= -0.19		
FSH vs. HOMA index	r= -0.11	r= -0.07	r= -0.71*	r= -0.17	r= -0.70*	r= -0.07		

HOMA, homeostasis model assessment; FSH, follicle-stimulating hormone. $*p \le 0.05$.



SPRINGER NATURE Springer Nature is through the combination of Nature Publishing Group, one of t), Palgra

Authorship change form

behind the name. 1 st author First 2 nd author NIC	Hirst name(s)
1 st author	ROSECUA
2 nd author	Nicolo
3 rd author	POSTAP A.
4 th author	MARCO
5 th author	SARDE
6 th author	87680
7 th author	AUD E.
Please use an additi	n additional sheet if there are mor

This form should be used by authors to request a


Please note: In author collaborations wher

provide each author's full name with the

Please read the important information

macmillan le jole o 3

Bohn Stafleu van Loghum



the world's leading we Macmillan, Maci	than 7 authors.								ip, in the order	JE POLYHOR	nuscript itle and/or DOI DOI if available.	e is formal agreem nge Form on beha	ny change in au me first followe
global research, educational and profession millan Education and Springer Science+Bu	Gete	SUCCESS	5	HUSHEC	GEOREL	HUNO	GNNRAUS	Family name	shown on the manuscript before	SAISH INFUENCES GLUCO	if available.)	reement for representing the collab behalf of all authors.	thorship including changes in correct by the family name.

Springer on page 4 before 3 97 you **M** O egin Q siness nal 2 0 esponding the ation, ADA publisher changes Media DOI ISBN Manuscript Ŧ (p S S Desi D authors F sufficient S created in (for W case er P 00 え ----Please May of oks for O 0 -0 T publish 2015 3 the 0 ce ... fully P . 2 3 rep U complete case lease resentative ed S manuscript: E indicate 0 unpublish all or sections. 0 C the

(For

books: Please provide the title, ISBN a

(For journals: Please

provide the

manuscri

Title:

コエー

2039

Þ

5

FI

RA

5

3

Section

2: Please provide


the

previous


au


Section 1: Please provide the current title

author) to complete and sign the Authorship Cha given na pt ID, ti e there thorsh Ind/or of mai £ D 0




Section 3: Please authorship? removed on the submitted manuscr Section adding 4th 55 2nd 5 th 3rd 1 7th author Please use JS 6 ス author author author author author author P is NGER VATURE Springer (CA) behind the name. 4: SWE D Proposed new authorship. 9 Please refer to an additional ワシ Y RO ろう provide First name(s) B D \$ 5 Z Bohn Stafleu van Loghum 3 7 PZ 6 F 2 sheet D 0 justification for change. the ma d -ti there are Bouillau うちろろ ipt.






(journal) policy pages for more information about authorship. Please explain why omitted Please use this section to explain your reasons for changing the authorship authors were not originally 01



If the corresponding author has changed, please indicate the reason under section Please provide your new authorship list



more than 7 authors.

Nature is one of the world's leading global research, educational and professional publishers, Group, Palgrave Macmillan, Macmillan Education and Springer Science+Business Media. created in May 2015

SP2


through the combination of Nature


Publishing

nge of authorship r n Q Cost

mily name (this name will appear in full on the final publication and will be

ω.





Springer

Bohn Stafleu van Loghum

90

-

macm

an

Springer

0

Rent

OC

D

D

dis

removed on authorship? Section 3 Please the Please refer to submitted provide 2 justification the manuscript (journal) OF policy change.

pages for

more

information about

Please use

this

section

to

exp

ain

N

our

Schauthor 6th 4th 7th Srd Nnd б 5 author author author author author author First name(s) 1 1 2 2 1

adding Section (CA) 4: Proposed behind the new name. authorship If the corresponding author has Please provide your new changed, please in authorship list in t he

one of the Palgrave Macmillan, world's leading global research, Macmillan Education educat and Springer ional and

SPRINGER NATURE

Springer Nature

S

through the

combination

of

Nature

Publishing

Group,

more than ノ authors.

Please

use

an


additional

sheet

Ŧ

there

are



S through State have New Your New Author Contributions Section 5: Author contribution, 三丁 New Disclosures (financial D 2 0 S manuscript - after approval of the change(s) - as our production department will not transfer the information in this acknowledgements: been removed as authors NGER NATURE Springer Nature is one of the world's leading global research, educational and professional publishers, 'Not the dis S combination of Nature 0 applicable' AND AND Sack Bohn Stafleu van Loghum 010 ÷ . there -20A and non-financial interests, funding): statemen g A STAN are no polorave macmillan 创 Springer and Acknowledgement ensure you state what contribution any t (if applicable per the journal policy):


Publishing Group, Palgrave Macmillan, Macmillan

new authors.

のうろうろ

R

and Disclosures. Please use this



SPRINCER NATURE Springer Natur through the combination of Nature Publishing Gr Please author ~ 6 H author 55 author 45 author authors β author th N author 1---nd st use an 4 Paro いなりとう 币 505 2 ROSE additional -6000 R 4 之う U 0° 5 D . sheet ÷ ゴ \$ SI th \bigcirc 0 77 5 ere Y 26 4 D いま 25 P 5 Z 2

Section the signature box are (NB: Please 6: Declaration of agreem print the form, unacceptabl sign and return e. ent. * All authors, a scanned copy. Please note that signatures that have been inserted unchanged, new and rem oved must sign this declaration.

First


name


רד

amily

Bohn Staffen Van Loghum mach

dis






-

ricase uciece da	s appropriate. Derete all of the boid if you v	vere on the original au	thorship list an
/ name		Signature	Affiliated inst
RES	I agree to the proposed new authorship shown in section 4 /and the addition/removal*of my name to the authorship list.	For the second	
	I agree to the proposed new authorship shown in section 4 /and the addition/removal*of my name to the authorship list.	Furs	E TERS
AL	I agree to the proposed new authorship shown in section 4 /and the addition/removal*of my name to the authorship list.	les es	SIGO
E.	I agree to the proposed new authorship shown in section 4 /and the addition/removal*of my name to the authorship list.	Kultur	CECO
2	I agree to the proposed new authorship shown in section 4 /and the addition/removal*of my name to the authorship list.		Even
5	I agree to the proposed new authorship shown in section 4 /and the addition/removal*of my name to the authorship list.		CHAR AND
GEB S	I agree to the proposed new authorship shown in section 4 /and the addition/removal*of my name to the authorship list.	4	GRANS
re more than 7	5		

Nature is one of the world's leading global research, educational and professional publishers, created in May 2015 ing Group, Palgrave Macmillan, Macmillan Education and Springer Science+Business Media.


of authorship reques



ptable as long as it is handwritten. Typed names nd are remaining as an author.

as an

ima




shown in section 4 /and the addition/removal*of my name to the authorship list.				
4 /and the 1*of my name to the	authorship list.			
4 /and the	addition/removal*	1		2
	shown in section 4	T	/	, nuthor
proposed new authorship	I agree to the prop		/	7 th
	authorship list.	/		
Il*of my name to the	addition/removal*	1		
shown in section 4 / and the	shown in section 4		/	author
posed new authorship	I agree to the prop			6 th
	authorship list.	/		
It of my name to the	addition/removal*	/		
4/and the	shown in section 4		/	author
posed new authorship	I agree to the prop			5#
	authorship list.	/		
ddition/removal*of my name to the	addition/removal*	1		
4 /and the	shown in section 4		/	authors
l agree to the proposed new authorship	l agree to the prop			4 th
	authorship list.	1		
Il*of my name to the	addition/removal*	7		
4 /and the	shown in section 4			author
lagree to the proposed new authorship	l agree to the prop			3 rd
	authorship list.	/		
I*of my name to the	addition/removal*	/		
4 /and the	shown in section 4			author
posed new authorship	I agree to the prop			2 nd
authorship list.	authorship list.			
addition/removal* of my name to the 1242 CARA	addition/removal*		760 5.	
A land the authorship 1 2 2 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	Shown in certion 4 /and	うろう	237	Author
Signature Annated Instit		Family name	First name	2

through the

--1

combination of Nature Publishing Group, Palgrave Macmillan, Macmillan Education a



SPR through ZGED the combination Z D TUR of Nature Publishing Springer Na ture Group, Palgrave S. one of the world's leading global research, Macmillan, Macmillan Education ed an 0 ucational and professional Springer Science+Business publishers, Media. created F May

COLUMN ST mportant information. Please read. dis

Bohn Stafleu van Loghum

macn

millan

Springer

06

Ne

Please We may return this choose to contact your institution form, fully completed, to

for

more

information

or unde

ertake

Springer

Nature.

We

will

con

isider

the

discipline, By signing and this declaration, al that only autho ors have authors

been added who made a meaningful

guarantee that the order of the a

uthors

are

0

must refer the matter to Please note, we cannot investigate your institution(s) for investigation. Please inform

or

mediate

any authorship disputes.

Th

You

are

Sn

.

Q.

cannot publish manuscripts v If you are not able to return a fully vhere completed form within 30 days authorship has not been agreed by a of th P 2 date authors that (including those who it was sent to the author have been reques ting 0 the ved

Incomplete forms will be rejected.

0

contribution to the if you need to a further information unable to ī accordance with investigation, obtained do this. you have work. their = agreement provided to appropriate scientific T befor ecide 0 ntribution, 0 whether makin change 90 -----= 2 5 applicabl fin appr din ma 2 decisio 0 P P as -P di -+ tions O P 3 apply auti emo orshi red) D σ 1 P Vou P