

The treatment of Kaposi's sarcoma: present and future options, a review of the literature

M. CECCARELLI^{1,2}, A. FACCIOLÀ¹, R. TAIBI³, G.F. PELLICANÒ⁴,
G. NUNNARI¹, E. VENANZI RULLO^{1,2}

¹Department of Clinical and Experimental Medicine, University of Messina, Messina, Italy

²Department of Pathology and Laboratory Medicine, School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

³Department of Medical Oncology, Centro di Riferimento Oncologico (CRO), Istituto Nazionale Tumori Aviano, Pordenone, Italy

⁴Department of Human Pathology of the Adult and the Developmental Age "G. Barresi", University of Messina, Messina, Italy

Abstract. – Kaposi's Sarcoma (KS) is an angiogenic tumor involving skin, mucosa and splanchnic organs. It is caused by Human Herpes Virus 8 (HHV8), when in the presence of other cofactors, such as an immune dysregulation. KS is particularly frequent in HIV-infected individuals. The major goals of treatment are to prevent disease progression, to reduce tumor and edema, to avoid organ compromise, and to relieve psychological stress. The importance and the high cancer risk offered by this co-infection, together with the spread of both these viruses, and the fact that angiogenesis is such an important characteristic of KS led to a lively interest in finding a definitive therapy. Most of the ongoing studies are focused on finding an application of old drugs in KS. Unfortunately, given the number of studies with different targets, it seems we are still far from completely understanding this disease and obtaining a "cure" which could be effective and safe for everyone. Further studies will hopefully offer new and definitive solutions.

Key Words:

Kaposi's Sarcoma, Human herpesvirus 8, HIV, PLWH, Angiogenesis, Treatment.

Introduction

The introduction of the combined antiretroviral therapy (cART) against the Human Immunodeficiency Virus (HIV) was followed by both positive and negative effects. A decrease of the comorbidities linked to the HIV-disease and an increase of the life expectancy for People Living With HIV (PLWH) was, as a matter of fact, accompanied by a relative increase in comorbidities

linked to ageing¹⁻²¹. Moreover, due to the increase of the life expectancy, cancer has become the first cause of death in PLWH⁷. Among the AIDS-related comorbidities, we assisted to a decrease of the so-called AIDS-defining cancers (ADCs). However, during the last few years, the increase in the numbers of late-presenters halted this positive trend, causing a recrudescence of pathologies such as the Kaposi's Sarcoma (KS)²². KS is an angiogenic tumor involving skin, mucosa and splanchnic organs. It is caused by Human Herpes Virus 8 (HHV8), when in presence of other cofactors, such as an immune deregulation. Its lesions are characterized by a chronic inflammatory infiltrate, and the infection with HHV8 leads to activation of cell pathways producing angiogenic and pro-inflammatory chemokines, and it also produces proteins having a cytokine-like action. As effect or a cause of this association with immune deregulation, KS is observed almost exclusively in immune depressed hosts, with the principal sub-population involved being represented by PLWH^{2,23-27}. On March 31st, 2019 we performed a systematic review of the literature regarding the link existing between angiogenesis and Kaposi's sarcoma in PLWH. We searched PubMed and Scopus applying "(angiogenesis) and (Kaposi's sarcoma)" as search terms. References of the articles retrieved were also reviewed. We limited the inclusion only to articles written in English. No age-related exclusion was performed. We identified 1314 records and, after duplicates were removed, we screened 983 of them. We excluded 689 articles by title and abstract, because they reported just one out of the two queries, and we assessed 294 full-texts for eligibility. After

assessment, we excluded 224 full-text articles because they reported about findings that were not confirmed in more recent research. At the end of the assessment, we included in our review 70 full-text articles reporting about the importance of angiogenesis in Kaposi's sarcoma in PLWH and the treatment available targeting this stage of the cancer. The aim of this paper is to review the available literature on KS and shed light on the role of antiangiogenic agents in PLWH.

Epidemiology

Moritz Kaposi first described the KS in 1872 as an angio-proliferative tumor of the elder, characterized by a survival not longer than 5 years²⁸⁻³⁰. Before the AIDS-era, it was particularly spread in Central Africa, Eastern Europe, Central Europe and Mediterranean regions²⁸⁻³¹.

Its distribution is similar to that of Human Herpes Virus 8 (HHV8), a gamma-herpesvirus discovered in 1994, also called Kaposi Sarcoma Herpes Virus (KSHV) to highlight its causal relationship with KS²⁸⁻³².

Since its discovery, it has become apparent that HHV8 infection prevalence is high in the general population, with child-to-child transmission being the most frequent transmission³³. However, in PLWH, and in particular among men who have sex with men (MSM), the transmission is mainly sexual, because of the use of saliva for lubrication during sexual intercourses^{28,29,31,32}. Another way of transmission, less frequent, is via blood transfusion and organ transplant. This iatrogenic transmission accounts for less than 3%, even where the seroprevalence is high^{28,29}. On the other hand, the transmission via organ transplant, though infrequent, is important because of the possible onset of iatrogenic KS^{28,29}.

KS is a burden in PLWH. In 1997 it was estimated that PLWH were 20,000 more likely to develop KS than the general population, and this estimate was confirmed to be true in 2010, highlighting HIV as the most important cofactor in developing KS^{29,32}. As a matter of fact, almost 50% of PLWH acquiring a HHV8 infection after a HIV infection develop KS.

Pathogenesis

KS, caused by HHV8, a B lymphotropic herpesvirus, with a latency/lytic reactivation lifecycle, is characterized by an intense angiogenesis^{34,35}.

Most of the proteins encoded by HHV8 genome are, as a matter of fact, involved in the promotion of the host cell survival and proliferation

and in the upregulation of angiogenesis. Viral FLICE inhibitory protein (v-FLIP), viral cyclin (v-cyclin), the latency-associated nuclear antigen (LANA) together with non-structural HHV8 membrane proteins, such as vGPCR, K1 and K15 are only a few of the viral products that variously interact with the host³⁴⁻³⁶. In particular, vFLIP increases uncontrolled angiogenesis by downregulating the Sin3A Associated Protein 18/Histone Deacetylase 1 (SAP18/HDAC1) complex. SAP18/HDAC1 action is mediated by the nuclear factor kappa-light-chain of activated B cells (NF-κB) and apoptosis activation³⁴⁻³⁶.

HHV8 genome also includes 12 MicroRNA precursor (pre-miRNA), which encode 25 microRNAs (miRNA)³⁵. It has been demonstrated that miRNA, and especially miRNA-K3 and miRNA-K6-5p, interact with the host upregulating the expression of different growth factors, cytokines and pathways. Among them: Platelet-Derived Growth Factor (PDGF), Epidermal Growth Factor-like domain 7 (EGFL7), Vascular Endothelial Growth Factor (VEGF), Angiopoietin-2 (Ang-2), rapamycin (mTOR) and c-kit signaling. All these elements are involved in angiogenesis upregulation^{37,38}. Upregulation of angiogenesis is important in the setting of KS and constitutes one of the fundamentals of this disease³⁹.

As a matter of fact, as previously hinted, KSHV interacts with the host in a number of ways, all converging in increasing the cell turnover and angiogenesis.

KS in PLWH

KS is particularly frequent in HIV-infected individuals. It is common knowledge that PLWH are affected by an increase in aberrant alternative splicing. This leads to an abnormal production of proteins and a dysregulation of the cell cycle⁴⁰.

Moreover, HIV and HHV8 variously interact, enhancing each other pathogenicity, and this is the main reason why KS incidence is so important in PLWH.

It has been demonstrated that HIV Viral Protein R (Vpr) inhibits HHV8 lytic cycle by upregulating a cellular miRNA which activates the NF-κB signaling⁴¹. Also HIV-Tat and HIV-Nef have been recognized to interact with HHV8 as promoters of angiogenesis and tumorigenesis⁴².

The importance and the high cancer risk offered by this co-infection, together with the spread of both these viruses, and the fact that angiogenesis is such an important characteristic of KS led to a lively interest in finding a definitive therapy.

Diagnosis

KS diagnosis is mainly clinical, especially in PLWH. The patient presents with various symptoms, but their principal complains are about red-violet cutaneous lesions in the lower extremities, face, and genitalia. These lesions are typically multifocal, with the appearance of papules, patches, plaques, or nodules³⁰.

However, KS can also have a visceral expression, for the diagnosis of which different techniques were studied. During the first decade of the 2000 it was demonstrated that a combination of thermography, laser Doppler imaging and near-infrared spectroscopy could be useful in diagnosis, staging and follow up of KS lesions⁴³.

Treatment

As for other herpes viruses, an eradicating treatment for HHV-8 does not exist. This fact makes scientists question whether it is possible to cure any form of KS. So, the major goals of treatment are to prevent disease progression, to reduce tumor and edema, to avoid organ impairment, and to relieve psychological stress.

Despite KS pathogenesis and treatment have been the object of many prospective randomized trials, they did not have any effect on the management of the disease over the last 20 years⁴⁴. cART is the pillar of KS treatment. Local therapy, radiotherapy and chemotherapy are applied on a case-severity basis. The choice is based on the rapidity of tumor growth, the extension and the number of the lesions, HIV viral load and CD4⁺ T-cell count, the general conditions of the patient and the existence of possible drug to drug interactions⁴⁵.

Since its introduction, cART dramatically changed the KS natural history^{25,46,47}. The observed decrease in incidence rates, progression of illness, visceral involvement and related mortality is likely due to an improved immune control in comparison with a pre-cART era.

cART seems to be so important that even when it is not completely successful in controlling HIV, its benefits among patients with KS are clear and independent from CD4⁺ T-cell count and HIV-RNA plasmatic viral load⁴⁷.

Several studies tried to demonstrate a direct effect of specific antiretroviral drugs in KS. A study conducted by Pati et al⁴⁸ showed the *in vitro* effects of ritonavir (RTV), a drug belonging to the Protease Inhibitors (PIs) class. RTV inhibits the production and secretion of several inflammatory cytokines and chemokines from the endo-

thelial cells, leading to a decreased proliferation and a stimulated apoptosis in KS cells. A similar study about PIs, performed *in vivo* on mice by Monini et al⁴⁹, showed that saquinavir and indinavir inhibits the secretion of basic fibroblast growth factor (bFGF) and VEGF. Despite these brilliant results *in vitro*, regimens containing PIs do not show superiority when compared to non-PI ones in observational studies in PLWH^{50,51}.

As a consequence, the onset of KS does not represent a reason for a cART regimen modification. Moreover, PI-containing regimens have the same effect on preventing KS compared to non-PI regimens⁴⁶.

Local Therapy

When a systemic treatment is not required for the severity of the disease, some options for a local treatment are available.

Vinblastine (VNB) is a natural Vinca alkaloid. It shows an antimetabolic cytotoxic effect and antiangiogenic activity even at low concentrations⁵². Typically, lesions treated with VNB regress but not resolve completely⁵³⁻⁵⁵. A different number of doses are required, according to the surface area of the lesions. For lesions larger than 0.5 cm², a second treatment cycle after a month is often required⁵³⁻⁵⁵.

When the area affected is wide, treatment with VNB is not possible. However, before resorting to systemic therapy, a cycle of radiation therapy can represent a valid option⁵⁶⁻⁵⁷.

Alitretinoin application is a second line option for local therapy. It showed a response *vs.* placebo of 35-37% *vs.* 7-8%^{58,59}.

Chemotherapy

Systemic chemotherapy should be added to cART whenever there is a progression of KS, presence of an extensive edema, massive skin involvement or symptomatic visceral involvement. Chemotherapy is also required in case of immune reconstitution inflammatory syndrome (IRIS).

Pegylated liposomal doxorubicin and liposomal daunorubicin represent the first line chemotherapeutic drugs in KS. Other agents are vinblastine, bleomycin, paclitaxel, vincristine and etoposide⁶⁰. Most of these drugs are currently under investigation in clinical trials, in various combinations between them (Table I).

First Line

Pegylated liposomal doxorubicin and liposomal daunorubicin represent the first choice in

Table I. Current Clinical Trials for Kaposi's Sarcoma.

Study title	Status	Study phase	NCT ID	Drugs
Pomalidomide in Combination With Liposomal Doxorubicin in People With Advanced or Refractory Kaposi Sarcoma	Recruiting	Phase 1	NCT02659930	Liposomal doxorubicin/ pomalidomide
sEphB4-HSA in Treating Patients With Kaposi Sarcoma	Recruiting	Phase 2	NCT02799485	Recombinant EphB4-HSA Fusion Protein
Nivolumab and Ipilimumab in Classical Kaposi Sarcoma (CKS)	Recruiting	Phase 2	NCT03219671	Nivolumab/Ipilimumab
An Evaluation of Tc 99m Tilmanocept by Intravenous (IV) and Subcutaneous (SC) Injection in Kaposi Sarcoma (KS)	Recruiting	Phase 1	NCT03157167	Tc99m-tilmanocept
Nelfinavir Mesylate in Treating Patients With Kaposi Sarcoma	Recruiting	Phase 2	NCT03077451	Nelfinavir Mesylate
Pomalidomide for Kaposi Sarcoma in People With or Without HIV	Active, not recruiting	Phase 1/ Phase 2	NCT01495598	Pomalidomide
Investigating Chemotherapy Treatments, Response and Subsets of HIV-associated Kaposi Sarcoma in Malawi	Recruiting		NCT03160183	ART (different regimens)
Evaluating Quality of Life in Patients With AIDS-Associated Kaposi Sarcoma Treated With Bleomycin and Vincristine	Recruiting	Phase 1	NCT03596918	Bleomycin Sulfate/Vincristine Sulfate
Compression Therapy for Leg Ulcers and Kaposi Sarcoma in Western Kenya	Recruiting	Not Applicable	NCT03404297	
Intra-lesional Nivolumab Therapy for Limited Cutaneous Kaposi Sarcoma	Recruiting	Early Phase 1	NCT03316274	Intra-lesional injection of nivolumab
Valganciclovir Four Weeks Prior to cART Initiation Compared to Standard Therapy for Disseminated Kaposi Sarcoma	Active, not recruiting	Phase 2	NCT03296553	Valganciclovir/ART
Phase II Multicentric Study of Pembrolizumab in Classic or Endemic Kaposi's Sarcoma	Recruiting	Phase 2	NCT03469804	Pembrolizumab
Antiretroviral Therapy (ART) Alone or With Delayed Chemo Versus ART With Immediate Chemo for Limited AIDS-related Kaposi's Sarcoma	Active, not recruiting	Phase 3	NCT01352117	Etoposide and efavirenz/emtricitabine/tenofovir disoproxil fumarate
Tocilizumab for KSHV-Associated Multicentric Castlemans Disease	Recruiting	Phase 2	NCT01441063	Zidovudine/Tocilizumab/Valganciclovir
History of the KSHV Inflammatory Cytokine Syndrome (KICS)	Recruiting	Phase 2	NCT01419561	Zidovudine/Liposomal Doxorubicin/Valganciclovir/Rituximab
Virotherapy and Natural History Study of KHSV-Associated Multicentric Castlemans Disease With Correlates of Disease Activity	Recruiting	Phase 2	NCT00092222	Etoposide/Interferon-alpha/Rituximab/Zidovudine/Liposomal Doxorubicin/Bortezomib/Valganciclovir/Vincristine/Cyclophosphamide/Filgrastim (G-CSF)/Prednisone/Sirolimus
Nivolumab and Ipilimumab in Treating Patients With HIV Associated Relapsed or Refractory Classical Hodgkin Lymphoma or Solid Tumors That Are Metastatic or Cannot Be Removed by Surgery	Recruiting	Phase 1	NCT02408861	Ipilimumab/Nivolumab
Pembrolizumab in Treating Patients With HIV and Relapsed, Refractory, or Disseminated Malignant Neoplasms	Recruiting	Phase 1	NCT02595866	Pembrolizumab
Nivolumab With or Without Varlilumab in Treating Patients With Relapsed or Refractory Aggressive B-cell Lymphomas	Recruiting	Phase 2	NCT03038672	Nivolumab/Varlilumab
Phase II Multicentric Study of Digoxin Per os in Classic or Endemic Kaposi' s Sarcoma	Recruiting	Phase 2	NCT02212639	Digoxin
Alpha Radiation Emitters Device for the Treatment of Cutaneous, Mucosal or Superficial Soft Tissue Neoplasia (DaRT)	Recruiting	Not Applicable	NCT03737734	Radiation: Diffusing Alpha Radiation Emitters Therapy (DaRT)
Study to Evaluate Safety and Tolerability of XmAb13676 in Patients With CD20-expressing Hematologic Malignancies	Recruiting	Phase 1	NCT02924402	XmAb13676

case of systemic chemotherapy. Both achieve the goal of decreasing edema and cancer volume. Moreover, they improve the typical KS hyperpigmentation with a response rate of 30-60%⁶¹⁻⁶⁴.

In a double-blind comparison between these two liposomal anthracyclines both tumor responses and clinical benefits were better in the doxorubicin arm⁶⁵. Side effects are usually mild in comparison with conventional combination chemotherapy regimens^{64,66}.

Taxanes are a family of chemotherapeutic agents, which act disrupting the microtubule function and cellular division. They showed promising results in KS.

In particular, paclitaxel demonstrated a great efficacy in KS. Compared to other drugs of the same family, paclitaxel is able to downregulate angiogenesis acting on different pathways. Its first effect on the microtubules leads to a decrease of cell proliferation, motility and migration. Paclitaxel also downregulates both VEGF and Angiopoietin-1 (Ang-1), and upregulates thrombospondin-1 (TSP-1)⁶⁷. Thus, paclitaxel is a valid second line alternative to liposomal anthracyclines or even the first choice in selected cases⁶⁸⁻⁷¹. However, paclitaxel high toxicity limits its use⁶⁸⁻⁷¹.

Docetaxel and cabazitaxel are other taxane drugs able to downregulate angiogenesis. However, their most common adverse effect is severe neutropenia and their use is not suggested in PLWH suffering from KS. Two important limitations in taxanes use are the necessity of premedication with glucocorticoids (not ideal for already immunosuppressed patients) and their metabolism through cytochrome P450.

Second Line

In case of failure of a first line treatment or when drug-drug interactions make their use difficult (i.e. paclitaxel and ritonavir) few second line options are available. Among them, the most used are etoposide^{72,73} (once daily and oral dose), vinorelbine^{74,75} (active even after first line treatment failure) and interferon-alfa⁷⁶ that has been tested in association with zidovudine⁷⁷.

Other Targets

Angiogenesis

Inhibitors of mTOR pathway, such as temsirolimus and rapamycin have been studied in the last years for the treatment of KS^{38,78-81}.

Following the observation that patients who experienced KS after transplant and have been treated successfully with rapamycin, this drug was applied in HIV-related KS obtaining only a partial success⁸²⁻⁸³.

VEGF could be a target for KS treatment^{84,85}. Bevacizumab, a monoclonal antibody against VEGF, showed a complete or partial response in 29% of the cases in a phase II study. Moreover, it results in stable disease in 64% of the cases⁸⁶. Anti-VEGF/VEGFR therapy is not currently used in KS.

Imatinib, an inhibitor of the PDGF and c-kit receptors, has been studied in a multicenter phase II trial³⁷⁻⁸⁷. It showed a partial response in 33 %, with an acceptable toxicity profile⁸⁷.

Fumagillin and thalidomide were studied in KS setting because of their known action on angiogenesis. However, their toxicities and lack of prolonged controls put them on the back burner⁸⁸⁻⁹².

Sorafenib, a VEGF, angiogenesis and tyrosine kinase inhibitor, has been studied in a phase Ib trial (NCT00287495). The importance of this study was in pointing out that most of the anti-angiogenic drugs were CYP3A4 inhibitors, with negative drug-drug interactions with cART⁹³.

PTC299 is an inhibitor of VEGFA mRNA translation showing promising results in a phase Ib trial (NCT00686842)⁹⁴. In three patients a partial response was observed and in eleven the disease did not progress. Notably, no differences by cART regimen were observed and the patients did not experience the typical VEGF-inhibitors adverse effects (bleeding, renal vascular injury, hypertension). Unfortunately, the redundancies in the VEGF feedback pathway implicates that PTC299 should be combined with other antiangiogenic agents to be effective.

HHV-8

Even if HHV-8 has been clearly identified as the etiologic agent of KS, there is no drug capable of its complete eradication.

Drugs used for the management of infections caused by other herpes viruses, like *ganciclovir* (and its prodrug *valganciclovir*), *foscarnet* and *acyclovir*, have been evaluated in HIV patients with KS⁹⁵⁻⁹⁷. All of them showed anti-HHV-8 activity, except for acyclovir. However, all these studies were carried out on patients treated with different cART regimens and under treatment for other HIV-related infections or complications.

The most robust data in our opinion are the ones on ganciclovir, which showed to reduce the risk of KS by 75-93% when compared with placebo⁹⁷.

Conclusions

Given the suboptimal results obtained so far with targeted therapies, a future combined approach should be considered. Table I resumes the currently active clinical trials. It is clear, from what we stated before and what is shown in the table, that targeting just one pathway of angiogenesis is not the right choice. As a matter of fact, the research is currently focused on finding the most effective combination of antiretrovirals, anti-HHV8 drugs and chemotherapeutic agents with the smallest burden in terms of toxicity for the patient.

Most of the ongoing studies are focused on finding an application of old drugs in KS. We can see how the efforts rotate around evaluating the efficacy of classic chemotherapeutic agents such as doxorubicin and bleomycin, two antimicrobials, and etoposide, a Topoisomerase II inhibitor, in new therapeutic combinations with antiretrovirals of every class (nelfinavir, efavirenz/emtricitabine/tenofovir disoproxil fumarate, zidovudine).

The interest in monoclonal antibodies is also definitively high. These drugs can effectively target specific pathways, leading to a combined block of an aberrant function. A lot of drugs are already available on the market and are currently being tested for their efficacy in the KS landscape.

A particular attention should be given to the studies regarding pomalidomide, a thalidomide derivative with an anti-angiogenic activity, bortezomib, a proteasome inhibitor, also with an anti-angiogenic activity, and sirolimus, or rapamycin, an antimicrobial with known effectiveness in Castleman's Disease, a hematologic disease also related to HHV8.

Unfortunately, given the number of studies with different targets, it seems we are still far from completely understanding this disease, obtaining a "cure" which could be effective and safe for everyone. Further studies will hopefully offer new and definitive solutions.

Conflict of Interest

The Authors declare that they have no conflict of interests.

Funding

The authors did not receive funds for this research.

Authors' Contribution

MC, AF and EVR searched the literature; MC, AF, MB, GFP, GN and EVR independently assessed the full-text articles; MC, AF and EVR wrote the article; GN and EVR reviewed the manuscript. All the authors read and accepted the final manuscript.

References

- 1) D'ALEO F, CECCARELLI M, VENANZI RULLO E, FACCIOLA A, DI ROSA M, PINZONE MR, CONDORELLI F, VISALLI G, PICERNO I, BERRETTA M, PELLICANÒ GF, NUNNARI G. Hepatitis C-related hepatocellular carcinoma: diagnostic and therapeutic management in HIV-patients. *Eur Rev Med Pharmacol Sci* 2017; 21: 5859-5867.
- 2) FACCIOLA A, VENANZI RULLO E, CECCARELLI M, D'ALEO F, DI ROSA M, PINZONE MR, CONDORELLI F, VISALLI G, PICERNO I, FISICHELLA R, NUNNARI G, PELLICANÒ GF. Kaposi's sarcoma in HIV-infected patients in the era of new antiretrovirals. *Eur Rev Med Pharmacol Sci* 2017; 21: 5868-5879.
- 3) D'ALEO F, VENANZI RULLO E, CECCARELLI M, FACCIOLA A, CONDORELLI F, PINZONE MR, CACOPARDO B, DI ROSA M, NUNNARI G, PELLICANÒ GF. HIV and colorectal cancer. New insights and review of the literature. *WCRJ* 2018; 5: e1122.
- 4) VISALLI G, FACCIOLA A, D'ALEO F, PINZONE MR, CONDORELLI F, PICERNO I, NUNNARI G, PELLICANÒ GF, CECCARELLI M, VENANZI RULLO E. HPV and urinary bladder carcinoma: a review of the literature. *WCRJ* 2018; 5: e1038.
- 5) CECCARELLI M, VENANZI RULLO E, FACCIOLA A, MADEDDU G, CACOPARDO B, TAIBI R, D'ALEO F, PINZONE MR, PICERNO I, DI ROSA M, VISALLI G, CONDORELLI F, NUNNARI G, PELLICANÒ GF. Head and neck squamous cell carcinoma and its correlation with human papillomavirus in people living with HIV: a systematic review. *Oncotarget* 2018; 9: 17171-17180.
- 6) FACCIOLA A, CECCARELLI M, VENANZI RULLO E, D'ALEO F, CONDORELLI F, VISALLI G, CACOPARDO B, PINZONE MR, DI ROSA M, NUNNARI G, PELLICANÒ GF. Prostate cancer in HIV-positive patients- a review of the literature. *WCRJ* 2018; 5: e1136.
- 7) D'ANDREA F, CECCARELLI M, VENANZI RULLO E, FACCIOLA A, D'ALEO F, CACOPARDO B, IACOBELLO C, COSTA A, ALTAVILLA G, PELLICANÒ GF, NUNNARI G. Cancer screening in HIV-infected patients: early diagnosis in a high-risk population. *WCRJ* 2018; 5: e1130.
- 8) D'ALEO F, CAMA BAV, PAOLUCCI IA, VENANZI RULLO E, CONDORELLI F, FACCIOLA A, DI FRANZIA R, SAVASTA A, PINZONE MR, PICERNO I, VISALLI G, NUNNARI G, PELLICANÒ GF, CECCARELLI M. New and old assumptions on lung cancer in people living with HIV. *WCRJ* 2018; 5: e1036.

- 9) D'ANDREA F, CECCARELLI M, FACCIOLA A, NUNNARI G, PELLICANÒ GF, VENANZI RULLO E. Breast cancer in women living with HIV. *Eur Rev Med Pharmacol Sci* 2019; 23: 1158-1164.
- 10) CECCARELLI M, VENANZI RULLO E, VACCARO M, FACCIOLA A, D'ALEO F, PAOLUCCI IA, CANNAVÒ SP, CACOPARDO B, PINZONE MR, PELLICANÒ GF, CONDORELLI F, NUNNARI G, GUARNERI C. HIV-associated psoriasis: epidemiology, pathogenesis, and management. *Dermatol Ther* 2019; 32: e12806
- 11) SPINA M, BERRETTA M, TIRELLI U. Hodgkin's disease in HIV. *Hematol Oncol Clin North Am* 2003; 17: 843-858.
- 12) HERNÁNDEZ-RAMÍREZ RU, QIN L, LIN H, LEYDEN W, NEUGEBAUER RS, ALTHOFF KN, HESSOL NA, ACHENBACH CJ, BROOKS JT, GILL MJ, GROVER S, HORBERG MA, LI J, MATHEWS WC, MAYOR AM, PATEL P, RABKIN CS, RACHLIS A, JUSTICE AC, MOORE RD, ENGELS EA, SILVERBERG MJ, DUBROW R; NORTH AMERICAN AIDS COHORT COLLABORATION ON RESEARCH AND DESIGN OF THE INTERNATIONAL EPIDEMIOLOGIC DATABASES TO EVALUATE AIDS. Association of immunosuppression and HIV viremia with anal cancer risk in persons living with HIV in the United States and Canada. *Clin Infect Dis* 2019; pii: ciz329. doi: 10.1093/cid/ciz329
- 13) COGHILL AE, HAN X, SUNEJA G, LIN CC, JEMAL A, SHIELS MS. Advanced stage at diagnosis and elevated mortality among US patients with cancer infected with HIV in the National Cancer Data Base. *Cancer* 2019; doi: 10.1002/cncr.32158.
- 14) BERRETTA M, MARTELOTTA F, DI FRANZIA R, SPINA M, VACCHER E, BALESTRERI L, BORSATTI E, BEARZ A, DE PAOLI P, TIRELLI U. Clinical presentation and outcome of non-AIDS defining cancers, in HIV-infected patients in the ART-era: the Italian Co-operative Group on AIDS and tumors activity. *Eur Rev Med Pharmacol Sci* 2015; 19: 3619-3634.
- 15) ATTERITANO M, MIRARCHI L, VENANZI RULLO E, SANTORO D, IARIA C, CATALANO A, LASCO A, ARCORACI V, GULLO LO A, BITTO A, SQUADRITO F, CASCIO A. Vitamin D status and the relationship with bone fragility fractures in HIV-infected patients: a case control study. *Int J Mol Sci* 2018; 19: 119.
- 16) VENANZI RULLO E, CECCARELLI M, CONDORELLI F, FACCIOLA A, VISALLI G, D'ALEO F, PAOLUCCI I, CACOPARDO B, PINZONE MR, DI ROSA M, NUNNARI G, PELLICANÒ GF. Investigational drugs in HIV: pros and cons of entry and fusion inhibitors (review). *Mol Med Rep* 2019; 19: 1987-1995.
- 17) ZANET E, BERRETTA M, MARTELOTTA F, CACOPARDO B, FISICHELLA R, TAVIO M, BERRETTA S, TIRELLI U. Anal cancer: focus on HIV-positive patients in the HAART-era. *Curr HIV Res* 2011; 9: 70-81.
- 18) PINZONE MR, CECCARELLI M, VENANZI RULLO E, MARESCA M, BRUNO R, CONDORELLI F, DI ROSA M, MADEDDU G, FOCÀ E, CALCAGNO A, CELESIA B. M, CACOPARDO B, NUNNARI G, PELLICANÒ GF. Circulating angiotensin-like protein 2 levels are associated with decreased renal function in HIV+ subjects on cART: a potential marker of kidney disease. *Biomed Rep* 2019; 10: 140-144.
- 19) FACCIOLA A, VENANZI RULLO E, CECCARELLI M, D'ALEO F, D'ANDREA F, VISALLI G, PINZONE MR, PICERNO I, CACOPARDO B, CONDORELLI F, MADEDDU G, NUNNARI G, PELLICANÒ GF. Hodgkin's lymphoma in people living with HIV: epidemiology and clinical management. *WCRJ* 2019; 6: e1295.
- 20) D'ANDREA F, PELLICANÒ GF, VENANZI RULLO E, D'ALEO F, FACCIOLA A, MICALI C, COCO M, VISALLI G, PICERNO I, CONDORELLI F, PINZONE MR, CACOPARDO B, NUNNARI G, CECCARELLI M. Cervical cancer in women living with HIV: a review of the literature. *WCRJ* 2019; 6: e1224.
- 21) CECCARELLI M, CONDORELLI F, VENANZI RULLO E, PELLICANÒ GF. Editorial - Improving access and adherence to screening tests for cancers: a new, though old, challenge in the HIV epidemics. *WCRJ* 2018; 5: e1030.
- 22) MARTÍNEZ SANZ J, PÉREZ ELÍAS MJ, MURIEL A, GÓMEZ AYERBE C, VIVANCOS GALLEGO MJ, SÁNCHEZ CONDE M, HERRERO DELGADO M, PÉREZ ELÍAS P, POLO BENITO L, DE LA FUENTE CORTÉS Y, BAREA R, SULLIVAN AK, FUSTER RUIZ DE APODACA MJ, GALINDO MJ, MORENO S; DRIVE 03, OPTTEST WP5 AND FOCO STUDY GROUPS. Outcome of an HIV education program for primary care providers: Screening and late diagnosis rates. *PLoS One* 2019; 14: e0218380.
- 23) LA FERLA L, PINZONE MR, NUNNARI G, MARTELOTTA F, LLESHI A, TIRELLI U, DE PAOLI P, BERRETTA M, CACOPARDO B. Kaposi's sarcoma in HIV-positive patients: the state of art in the HAART-era. *Eur Rev Med Pharmacol Sci* 2013; 17: 2354-2365.
- 24) PINZONE MR, BERRETTA M, CACOPARDO B, NUNNARI G. Epstein-barr virus- and Kaposi sarcoma-associated herpesvirus-related malignancies in the setting of human immunodeficiency virus infection. *Semin Oncol* 2015; 42: 258-271.
- 25) FRANCESCHI S, MASO L D, RICKENBACH M, POESEL J, HIRSCHEL B, CAVASSINI M, BORDONI A, ELZI L, ESS S, JUNDT G, MUELLER N, CLIFFORD GM. Kaposi sarcoma incidence in the Swiss HIV cohort study before and after highly active antiretroviral therapy. *Br J Cancer* 2008; 99: 800-804.
- 26) COLAFIGLI M, BONADIES A, FERRARESI V, TONACHELLA R, CRISTAUDDO A, LATINI A. Kaposi Sarcoma in HIV-infected patients: an infectious-dermatological outpatient service experience. *Infect Dis Trop Med* 2017; 3: e410.
- 27) LA FERLA L, LO PRESTI COSTANTINO MR, MONDELLO P. Kaposi's sarcoma in HIV-infected patients: a review of the literature. *Infect Dis Trop Med* 2016; 2: e239.
- 28) GONCALVES PH, ULDRICK TS, YARCHOAN R. HIV-associated Kaposi sarcoma and related diseases. *AIDS* 2017; 31: 1903-1916.
- 29) SUNG JCY, LOUIE SG, PARK SY. Kaposi's sarcoma: Advances in tumor biology and pharmacotherapy. *Pharmacotherapy* 1997; 17: 670-683.
- 30) CAI Q, YUAN Z, LAN K. Infectious Agents Associated Cancers: Epidemiology and Molecular Biology. Springer Nature, 2017.
- 31) SPANO JP, ATLAN D, BREAU JL, FARGE D. AIDS and non-AIDS-related malignancies: a new vex-

- ing challenge in HIV-positive patients. Part II. Cervical and anal squamous epithelial lesions, lung cancer, testicular germ cell cancers, and skin cancers. *Eur J Intern Med* 2002; 13: 170-179.
- 32) MESRI EA, CESARMAN E, BOSHOFF C. Kaposi's sarcoma and its associated herpesvirus. *Nature reviews*. *Cancer* 2010; 10: 707-719.
 - 33) HENGGE UR, RUZICKA T, TYRING SK, STUSCHKE M, ROGGENDORF M, SCHWARTZ RA, SEEBER S. Update on Kaposi's sarcoma and other HHV8 associated diseases. Part 1: epidemiology, environmental predispositions, clinical manifestations, and therapy. *Lancet Infect Dis* 2002; 2: 281-292.
 - 34) LI X, BHADURI-MCINTOSH S. A central role for STAT3 in gammaherpesvirus-life cycle and -diseases. *Front Microbiol* 2016; 7: 1052.
 - 35) LI W, JIA X, SHEN C, ZHANG M, XU J, SHANG Y, ZHU K, HU M, YAN Q, QIN D, LEE MS, ZHU J, LU H, KRUEGER BJ, RENNE R, GAO SJ, LU C. A KSHV microRNA enhances viral latency and induces angiogenesis by targeting GRK2 to activate the CXCR2/AKT pathway. *Oncotarget* 2016; 7: 32286-32305.
 - 36) ABERE B, SCHULZ TF. KSHV non-structural membrane proteins involved in the activation of intracellular signaling pathways and the pathogenesis of Kaposi's sarcoma. *Curr Opin Virol* 2016; 20: 11-19.
 - 37) CAVALLIN LE, MA Q, NAIPAUE J, GUPTA S, KURIAN M, LOCATELLI P, ROMANELLI P, NADJI M, GOLDSCHMIDT-CLERMONT PJ, MESRI EA. KSHV-induced ligand mediated activation of PDGF receptor-alpha drives Kaposi's sarcomagenesis. *PLoS Pathog* 2018; 14: e1007175-28.
 - 38) KERR DA, BUSARLA SVP, GIMBEL DC, SOHANI AR, NAZARIAN RM. mTOR, VEGF, PDGFR, and c-kit signaling pathway activation in Kaposi sarcoma. *Hum Pathol* 2017; 65: 157-165.
 - 39) MARIGGIÒ G, KOCH S, SCHULZ TF. Kaposi sarcoma herpesvirus pathogenesis. *Phil Trans R Soc B* 2017; 372: 20160275-20.
 - 40) MTHEMBU NN, MBITA Z, HULL R, DLAMINI Z. Abnormalities in alternative splicing of angiogenesis-related genes and their role in HIV-related cancers. *HIV AIDS (Auckl)* 2017; 9: 77-93.
 - 41) YAN Q, SHEN C, QIN J, LI W, HU M, LU H, QIN D, ZHU J, GAO SJ, LU C. HIV-1 Vpr inhibits Kaposi's sarcoma-associated herpesvirus lytic replication by inducing microRNA miR-942-5p and activating NF- κ B signaling. *J Virol* 2016; 90: 8739-8753.
 - 42) NUNNARI G, SMITH JA, DANIEL R. HIV-1 Tat and AIDS-associated cancer: targeting the cellular anti-cancer barrier? *J Exp Clin Cancer Res* 2008; 27: 3.
 - 43) VOGEL A, DASGEB B, HASSAN M, AMYOT F, CHERNOMORDIK V, TAO Y, DEMOS SG, WYVILL K, ALEMAN K, LITTLE R, YARCHOAN R, GANDJBAKHCHE AH. Using quantitative imaging techniques to assess vascularity in AIDS-related Kaposi's sarcoma. *IEEE* 2006; 1: 232-235.
 - 44) DEZUBE BJ. Management of AIDS-related Kaposi's sarcoma: advances in target discovery and treatment. *Expert Rev Anticancer Ther* 2002; 2: 193-200.
 - 45) STEBBING J, SANITT A, NELSON M, POWLES T, GAZZARD B, BOWER M. A prognostic index for AIDS-associated Kaposi's sarcoma in the era of highly active antiretroviral therapy. *Lancet* 2006; 367: 1495-1502.
 - 46) GRABAR S, ABRAHAM B, MAHAMAT A, DEL GIUDICE P, ROSENTHAL E, COSTAGLIOLA D. Differential impact of combination antiretroviral therapy in preventing Kaposi's sarcoma with and without visceral involvement. *J Clin Oncol* 2006; 24: 3408-3414.
 - 47) GALLAFENT JH, BUSKIN SE, DE TURK PB, ABOULAFIA DM. Profile of patients with Kaposi's sarcoma in the era of highly active antiretroviral therapy. *J Clin Oncol* 2005; 23: 1253-1260.
 - 48) PATI S, PELSER CB, DUFRANE J, BRYANT JL, REITZ MS, WEICHHOLD, FF. Antitumorigenic effects of HIV protease inhibitor ritonavir: inhibition of Kaposi sarcoma. *Blood* 2002; 99: 3771-3779.
 - 49) MONINI P, SGADARI C, BARILLARI G, ENSOLI B. HIV protease inhibitors: antiretroviral agents with anti-inflammatory, anti-angiogenic and anti-tumour activity. *J Antimicrob Chemother* 2003; 51: 207-211.
 - 50) SGADARI C, BARILLARI G, TOSCHI E, CARLEI D, BACIGALUPO I, BACCARINI S, PALLADINO C, LEONE P, BUGARINI R, MALAVASI L, CAFARO A, FALCHI M, VALDEMBRI D, REZZA G, BUSSOLINO F, MONINI P, ENSOLI B. HIV protease inhibitors are potent anti-angiogenic molecules and promote regression of Kaposi sarcoma. *Nat Med* 2002; 8: 225-232.
 - 51) GILL J, BOURBOULIA D, WILKINSON J, HAYES P, COPE A, MARCELIN A. G, CALVEZ, V, GOTCH F, BOSHOFF C, GAZZARD B. Prospective study of the effects of antiretroviral therapy on Kaposi sarcoma - Associated herpesvirus infection in patients with and without Kaposi sarcoma. *J Acquir Immune Defic Syndr* 2002; 31: 384-390.
 - 52) CHAPLIN DJ. The effect of therapy on tumour vascular function. *Int J Radiat Biol* 1991; 60: 311-325.
 - 53) RAMÍREZ-AMADOR V, ESQUIVEL-PEDRAZA L, LOZADA-NUR F, DE LA ROSA-GARCÍA E, VOLKOW-FERNÁNDEZ P, SÚCHIL-BERNAL L, MOHAR A. Intralesional vinblastine vs. 3% sodium tetradecyl sulfate for the treatment of oral Kaposi's sarcoma. A double blind, randomized clinical trial. *Oral Oncol* 2002; 38: 460-467.
 - 54) MCCORMICK SU. Intralesional vinblastine injections for the treatment of oral Kaposi's sarcoma: Report of 10 patients with 2-year follow-up. *J Oral Maxillofac Surg* 1996; 54: 583-587.
 - 55) EPSTEIN JB. Treatment of oral Kaposi sarcoma with intralesional vinblastine. *Cancer* 1993; 71: 1722-1725.
 - 56) DONATO V, GUARNACCIA R, DOGNINI J, DE PASCALIS G, CARUSO C, BELLAGAMBA R, MORRONE A. Radiation therapy in the treatment of HIV-related Kaposi's sarcoma. *Anticancer Res* 2013; 33: 2153-2157.
 - 57) OLWENY CLM, BOROK M, GUDZA I, CLINCH J, CHEANG M, KIIRE CF, LEVY L, OTIM-OYET D, NYAMASVE J, SCHIPPER H. Treatment of AIDS-associated Kaposi's

- sarcoma in Zimbabwe: results of a randomized quality of life focused clinical trial. *Int J Cancer* 2004; 113: 632-639.
- 58) WALMSLEY S, NORTHFELT DW, MELOSKY B, CONANT MA, FRIEDMAN-KIEN AE, WAGNER B. Treatment of AIDS-related cutaneous Kaposi's sarcoma with topical alitretinoin (9-cis-retinoic acid) gel. Panretin Gel North American Study Group. *J Acquir Immune Defic Syndr* 1999; 22: 235-246.
 - 59) BODSWORTH N. J, BLOCH M, BOWER M, DONNELL D, YOCUM R. International Panretin Gel KS Study Group Phase III vehicle-controlled, multi-centered study of topical alitretinoin gel 0.1% in cutaneous AIDS-related Kaposi's sarcoma. *Am J Clin Dermatol* 2001; 2: 77-87.
 - 60) LEE FC, MITSUYASU RT. Chemotherapy of AIDS-related Kaposi's sarcoma. *Hematol Oncol Clin North Am* 1996; 10: 1051-1068.
 - 61) HAART is the centre of graduated systemic treatment of AIDS-associated Kaposi's sarcoma. *Drugs Ther Perspect* 2005; 21: 14-17.
 - 62) STEWART S, JABLONOWSKI H, GOEBEL FD, ARASTEH K, SPITILE M, RIOS A, ABOULAFIA DM, GALLESCHAW J, DEZUBE BJ. Randomized comparative trial of pegylated liposomal doxorubicin versus bleomycin and vincristine in the treatment of AIDS-related Kaposi's sarcoma. International Pegylated Liposomal Doxorubicin Study Group. *J Clin Oncol* 1998; 16: 683-691.
 - 63) NORTHFELT DW, DEZUBE BJ, THOMMES JA, MILLER BJ, FISCHL MA, FRIEDMAN-KIEN A, KAPLAN LD, MOND DU C, MAMELOK RD, HENRY DH. Pegylated-liposomal doxorubicin versus doxorubicin, bleomycin, and vincristine in the treatment of AIDS-related Kaposi's sarcoma: results of a randomized phase III clinical trial. *J Clin Oncol* 1998; 16: 2445-2451.
 - 64) GILL PS, WERNZ J, SCADDEN DT, COHEN P, MUKWAYA GM, VON ROENN JH, JACOBS M, KEMPIN S, SILVERBERG I, GONZALES G, RARICK MU, MYERS AM, SHEPHERD F, SAWKA C, PIKE MC, ROSS ME. Randomized phase III trial of liposomal daunorubicin versus doxorubicin, bleomycin, and vincristine in AIDS-related Kaposi's sarcoma. *J Clin Oncol* 1996; 14: 2353-2364.
 - 65) COOLEY TP, HENRY D, TONDA M, SUN S, O'CONNELL M, RACKOFF W. A randomized, double-blind study of pegylated liposomal doxorubicin for the treatment of AIDS-related Kaposi's sarcoma. *Oncologist* 2007; 12: 114-123.
 - 66) YOUNG AM, DHILLON T, BOWER M. Cardiotoxicity after liposomal anthracyclines. *Lancet Oncol* 2004; 5: 654.
 - 67) BOCCI G, DI PAOLO A, DANESI R. The pharmacological bases of the antiangiogenic activity of paclitaxel. *Angiogenesis* 2013; 16: 481-492.
 - 68) WELLES L, SAVILLE MW, LIETZAU J, PLUDA JM, WYVILL KM, FEUERSTEIN I, FIGG WD, LUSH R, ODOM J, WILSON WH, FAJARDO MT, HUMPHREY RW, FEIGAL E, TUCK D, STEINBERG SM, BRODER S, YARCHOAN R. Phase II trial with dose titration of paclitaxel for the therapy of human immunodeficiency virus-associated Kaposi's sarcoma. *J Clin Oncol* 1998; 16: 1112-1121.
 - 69) GILL PS, TULPUL A, ESPINA BM, CABRIALES S, BRESNAHAN J, ILAW M, LOUIE S, GUSTAFSON NF, BROWN MA, ORCUTT C, WINOGRAD B, SCADDEN DT. Paclitaxel is safe and effective in the treatment of advanced AIDS-related Kaposi's sarcoma. *J Clin Oncol* 1999; 17: 1876-1883.
 - 70) TULPUL A, GROOPMAN J, SAVILLE MW, HARRINGTON W, FRIEDMAN-KIEN A, ESPINA BM, GARCES C, MANTELLE L, METTINGER K, SCADDEN DT, GILL PS. Multicenter trial of low-dose paclitaxel in patients with advanced AIDS-related Kaposi sarcoma. *Cancer* 2002; 95: 147-154.
 - 71) STEBBING J, WILDFIRE A, PORTSMOUTH S, POWLES T, THIRLWELL C, HEWITT P, NELSON M, PATTERSON S, MANDALIA S, GOTCH F, GAZZARD BG, BOWER M. Paclitaxel for anthracycline-resistant AIDS-related Kaposi's sarcoma: clinical and angiogenic correlations. *Ann Oncol* 2003; 14: 1660-1666.
 - 72) SPRINZ E, CALDAS AP, MANS DR, CANCELA A, DI LEONE L, DALLA COSTA T, SCHWARTSMANN G. Fractionated doses of oral etoposide in the treatment of patients with aids-related Kaposi sarcoma: a clinical and pharmacologic study to improve therapeutic index. *Am J Clin Oncol* 2001; 24: 177-184.
 - 73) EVANS SR, KROWN SE, TESTA MA, COOLEY TP, VON ROENN JH. Phase II evaluation of low-dose oral etoposide for the treatment of relapsed or progressive AIDS-related Kaposi's sarcoma: an AIDS clinical trials group clinical study. *J Clin Oncol* 2002; 20: 3236-3241.
 - 74) BRIASOULIS E, PAPPAS P, PUOZZO C, TOLIS C, FOUNTZILAS G, DAFNI U, MARSELOS M, PAVLIDIS N. Dose-ranging study of metronomic oral vinorelbine in patients with advanced refractory cancer. *Clin Cancer Res* 2009; 15: 6454-6461.
 - 75) NASTI G, MARTELLOTTA F, BERRETTA M, MENA M, FASAN M, DI PERRI G, TALAMINI R, PAGANO G, MONTRONI M, CINELLI R, VACCHER E, D'ARMINIO MONFORTE A, TIRELLI U; GICAT; ICONA. Impact of highly active antiretroviral therapy on the presenting features and outcome of patients with acquired immunodeficiency syndrome-related Kaposi sarcoma. *Cancer* 2003; 98: 2440-2446.
 - 76) KROWN SE, LI P, VON ROENN JH, PAREDES J, HUANG J, TESTA MA. Efficacy of low-dose interferon with antiretroviral therapy in Kaposi's sarcoma: a randomized phase II AIDS Clinical Trials Group Study. *J Interferon Cytokine Res* 2002; 22: 295-303.
 - 77) SHEPHERD FA, BEAULIEU R, GELMON K, THUOT CA, SAWKA C, READ S, SINGER J. Prospective randomized trial of two dose levels of interferon alfa with zidovudine for the treatment of Kaposi's sarcoma associated with human immunodeficiency virus infection: a Canadian HIV Clinical Trials Network study. *J Clin Oncol* 1998; 16: 1736-1742.
 - 78) DITTMER DP, KROWN SE. Targeted therapy for Kaposi's sarcoma and Kaposi's sarcoma-associated herpesvirus. *Curr Opin Oncol* 2007; 19: 452-457.
 - 79) WANG L, DITTMER DP, TOMLINSON CC, FAKHARI FD, DAMANIA B. Immortalization of primary endothelial cells by the K1 protein of Kaposi's sarcoma-associated herpesvirus. *Cancer Res* 2006; 66: 3658-3666.

- 80) STALLONE G, INFANTE B, PONTRELLI P, RANIERI E, LOVERRE A, SCHEA A, CORMIO L, CARRIERI G, SCHEA FP, GRANDALIANO G, GESUALDO L. ID2-VEGF-related pathways in the pathogenesis of Kaposi's sarcoma: a link disrupted by rapamycin. *Am J Transplant* 2009; 9: 558-566.
- 81) CHAISUPARAT R, HU J, JHAM BC, KNIGHT ZA, SHOKAT KM, MONTANER S. Dual inhibition of PI3K and mTOR as an alternative treatment for Kaposi's sarcoma. *Cancer Res* 2008; 68: 8361-8368.
- 82) STALLONE G, SCHEA A, INFANTE B, DI PAOLO S, LOVERRE A, MAGGIO G, RANIERI E, GESUALDO L, SCHEA FP, GRANDALIANO G. Sirolimus for Kaposi's sarcoma in renal-transplant recipients. *New Engl J Med* 2005; 352: 1317-1323.
- 83) KROWN SE, ROY D, LEE JY, DEZUBE BJ, REID EG, VENKATARAMANAN R, HAN K, CESARMAN E, DITTMER DP. Rapamycin with antiretroviral therapy in AIDS-associated Kaposi sarcoma. *J Acquir Immune Defic Syndr* 2012; 59: 447-454.
- 84) HADDAD L, EL HAJJ, H, ABOU-MERHI R, KFOURY Y, MAHIEUX R, EL-SABBAN M, BAZARBACHI A. KSHV-transformed primary effusion lymphoma cells induce a VEGF-dependent angiogenesis and establish functional gap junctions with endothelial cells. *Leukemia* 2007; 22: 826-834.
- 85) ABLANEDO-TERRAZAS Y, ALVARADO-DE LA BARRERA C, ORMSBY CE, RUIZ-CRUZ M, REYES-TERÁN G. Intralesional bevacizumab in patients with human immunodeficiency virus-associated Kaposi's sarcoma in the upper airway. *Laryngoscope* 2014; 125: E132-E137.
- 86) ULDRICK TS, WYVILL KM, KUMAR P, O'MAHONY D, BERNSTEIN W, ALEMAN K, POLIZZOTTO MN, STEINBERG SM, PITLUGA S, MARSHALL V, WHITBY D, LITTLE RF, YARCHOAN R. Phase II study of bevacizumab in patients with HIV-associated Kaposi's sarcoma receiving antiretroviral therapy. *J Clin Oncol* 2012; 30: 1476-1483.
- 87) KOON HB, KROWN SE, LEE JY, HONDA K, RAPISUWON S, WANG Z, ABOULAFIA DM, REID EG, RUDEK MA, DEZUBE BJ, NOY A. Phase II trial of imatinib in AIDS-associated Kaposi's sarcoma: AIDS malignancy consortium protocol 042. *J Clin Oncol* 2014; 32: 402-408.
- 88) DEZUBE BJ, VON ROENN JH, HOLDEN-WILTSE J, CHEUNG TW, REMICK SC, COOLEY TP, MOORE J, SOMMADOSSI JP, SHRIVER SL, SUCKOW CW, GILL PS. Fumagillin analog in the treatment of Kaposi's sarcoma: a phase I AIDS Clinical Trial Group study. AIDS Clinical Trial Group No. 215 Team. *J Clin Oncol* 1998; 16: 1444-1449.
- 89) MOORE JD, DEZUBE BJ, GILL P, ZHOU XJ, ACOSTA EP, SOMMADOSSI JP. Phase I dose escalation pharmacokinetics of O-(chloroacetylcarbonyl) fumagillin (TNP-470) and its metabolites in AIDS patients with Kaposi's sarcoma. *Cancer Chemother Pharmacol* 2000; 46: 173-179.
- 90) STEFF M, JOLY V, DI LUCCA J, FELDMAN J, BURG S, SARDA-MANTEL L, PEYTAVIN G, MARINHO E, CRICKX B, RAYMOND E, LARIVEN S, MAUBEC E. Clinical activity of lenalidomide in visceral human immunodeficiency virus-related Kaposi sarcoma. *JAMA Dermatol* 2013; 149: 1319-1322.
- 91) BUNN MR. Use of thalidomide in the management of three HIV seroreactive children with Kaposi's sarcoma referred for palliative care. *Malawi Med J* 2008; 20: 143-146.
- 92) LITTLE RF, WYVILL KM, PLUDA JM, WELLES L, MARSHALL V, FIGG WD, NEWCOMB FM, TOSATO G, FEIGAL E, STEINBERG SM, WHITBY D, GOEDERT JJ, YARCHOAN R. Activity of thalidomide in AIDS-related Kaposi's sarcoma. *J Clin Oncol* 2000; 18: 2593-2602.
- 93) ULDRICK TS, GONCALVES PH, WYVILL KM, PEER CJ, BERNSTEIN W, ALEMAN K, POLIZZOTTO MN, VENZON D, STEINBERG SM, MARSHALL V, WHITBY D, LITTLE RF, WRIGHT JJ, RUDEK MA, FIGG WD, YARCHOAN R. A phase Ib study of sorafenib (BAY 43-9006) in patients with Kaposi sarcoma. *Oncologist* 2017; 22: e505-e549.
- 94) BENDER IGNACIO RA, LEE JY, RUDEK MA, DITTMER DP, AMBINDER RF, KROWN SE, ABOULAFIA DM. AIDS Malignancy Consortium (AMC)-059 Study Team. A phase 1b/pharmacokinetic trial of PTC299, a novel posttranscriptional VEGF inhibitor, for AIDS-related Kaposi's sarcoma: AIDS malignancy consortium trial 059. *J Acquir Immune Defic Syndr* 2016; 72: 52-57.
- 95) MOCROFT A, YOULE M, GAZZARD B, MORCINEK J, HALAI R, PHILLIPS AN. Anti-herpesvirus treatment and risk of Kaposi's sarcoma in HIV infection. Royal Free/Chelsea and Westminster Hospitals Collaborative Group. *AIDS* 1996; 10: 1101-1105.
- 96) GLESBY MJ, HOOVER DR, WENG S, GRAHAM NMH, GRAHAM NM, PHAIR JP, DETELS R, HO M, SAAH AJ. Use of antiherpes drugs and the risk of Kaposi's sarcoma: data from the Multicenter AIDS Cohort Study. *J Infect Dis* 1996; 173: 1477-1480.
- 97) MARTIN DF, KUPPERMANN BD, WOLITZ RA, PALESTINE AG, LI H, ROBINSON CA. Oral ganciclovir for patients with cytomegalovirus retinitis treated with a ganciclovir implant. Roche Ganciclovir Study Group. *N Engl J Med* 1999; 340: 1063-1070.