


REVIEW

Open Access



β -Secretase1 biological markers for Alzheimer's disease: state-of-art of validation and qualification

Harald Hampel¹, Simone Lista^{1,2,3}, Eugeen Vanmechelen⁴, Henrik Zetterberg^{5,6,7,8}, Filippo Sean Giorgi⁹, Alessandro Galgani¹⁰, Kaj Blennow^{6,7}, Filippo Caraci^{11,12}, Brati Das¹³, Riqiang Yan¹³, Andrea Vergallo^{1*}  and for the Alzheimer's Precision Medicine Initiative (APMI)

Abstract

β -Secretase1 (BACE1) protein concentrations and rates of enzyme activity, analyzed in human bodily fluids, are promising candidate biological markers for guidance in clinical trials investigating BACE1 inhibitors to halt or delay the dysregulation of the amyloid- β pathway in Alzheimer's disease (AD). A robust body of evidence demonstrates an association between cerebrospinal fluid/blood BACE1 biomarkers and core pathophysiological mechanisms of AD, such as brain protein misfolding and aggregation, neurodegeneration, and synaptic dysfunction.

In pharmacological trials, BACE1 candidate biomarkers may be applied to a wide set of contexts of use (CoU), including proof of mechanism, dose-finding, response and toxicity dose estimation. For clinical CoU, BACE1 biomarkers show good performance for prognosis and disease prediction.

The roadmap toward validation and qualification of BACE1 biomarkers requires standardized pre-analytical and analytical protocols to reduce inter-site variance that may have contributed to inconsistent results.

BACE1 biomarker-drug co-development programs, including biomarker-guided outcomes and endpoints, may support the identification of sub-populations with a higher probability to benefit from BACE1 inhibitors with a reduced risk of adverse effects, in line with the evolving precision medicine paradigm.

Keywords: Alzheimer's disease, Amyloid- β pathway, Axonal damage, BACE1, Clinical trials, Context of use, Fluid biomarkers, Neurodegeneration

Introduction

β -Site amyloid precursor protein (APP) cleaving enzyme 1 (BACE1) is a type I transmembrane aspartyl protease widely expressed in the brain, particularly in neurons, oligodendrocytes, and astrocytes [1–3]. BACE1 is expressed at the plasma endothelial membrane and in the endosomal compartments and has been detected in

healthy synaptic terminals. BACE1 functions as the β -secretase enzyme by cleaving the transmembrane APP to release the β -stubs and represents the rate-limiting catalytic step for A β production (see Fig. 1) [1–3].

High BACE1 concentrations (probably reflecting gene expression levels) and enzymatic activity were found in human AD brain extracts, consistent with experimental evidence that neurons express higher levels of A β in AD compared to “cognitively healthy aging.” In addition, a relatively large accumulation of BACE1 was found in neuritic dystrophies in close proximity of A β plaques both in AD amyloidogenic transgenic mouse models and

* Correspondence: avergallo@outlook.com

¹Sorbonne University, GRC no 21, Alzheimer Precision Medicine (APM), AP-HP, Pitié-Salpêtrière Hospital, Paris, France

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 1 Schematic representation of amyloidogenic and non-amyloidogenic pathways. Footnote: Three main proteases— α -, β -, and γ -secretases—are involved in APP processing through the amyloidogenic pathway (sequential cleavage by β - and γ -secretases), promoting amyloid- β ($A\beta$) production, and the non-amyloidogenic pathway in which $A\beta$ is cleaved in the middle, either directly by α -secretase (generating soluble APP α) or by the sequential cleavage by β -secretase and α -secretase (generating shorter $A\beta$ species such as $A\beta_{1-15}$ and $A\beta_{1-16}$). The two pathways lead to the production of different by-products with different intrinsic functional properties, putative physiological roles, and pathophysiological potential. In particular, BACE1 serves as the β -secretase enzyme by cleaving the transmembrane APP to release the β -stubs. BACE1 cleavage of APP represents the rate-limiting step for $A\beta$ production. Cleavage of APP by BACE1 liberates the soluble N-terminus of APP, while the C-terminal fragment (CTF- β or C99) remains bound to the membrane. To produce $A\beta$, the fragment CTF- β is cleaved by γ -secretase, an aspartyl-type protease membrane protein complex, which finally releases $A\beta$ into the extracellular space and the APP intracellular domain into the cytoplasm. The γ -secretase consists of different components. The catalytic components of the membrane-embedded tetrameric γ -secretase complex are represented by presenilins 1 and 2, intramembrane-cleaving proteases (I-CLiPs), responsible for generating the $A\beta$ carboxyl terminus from APP. In a parallel competing non-amyloidogenic pathway, APP is cleaved either by α -secretase or η -secretase to release two additional variants of the APP ectodomain, namely sAPP α and sAPP η . In vitro studies have shown that ADAM-10, a disintegrin and metalloprotease belonging to the family proteases, is the major α -secretase responsible for the ectodomain shedding of APP in the mouse brain and likely to be active in humans. APP is a type I transmembrane protein, highly expressed in neurons and abundant at the synapse. Although a full understanding of its function remains elusive, studies have suggested a role in the remodeling of dendritic spines, neurotransmission, synaptic plasticity, and maintenance of excitation-inhibition (E/I) balance. Soluble sAPP α and sAPP β are hypothesized to modulate basal synaptic transmission and short-term synaptic facilitation likely through GABAB receptor subunit 1a-mediated synaptic effect. Note: Adapted from [4]. Reproduced with permission

in AD brains, and this presence may promote cyclic $A\beta$ production [3, 5, 6].

Although BACE1 mutations have not yet been linked to AD risk, genetic variants surrounding the β -secretase site in the APP (including the Swedish mutation *KM/NL*, the Italian variant *A673V*, and the *A673T*) are associated with either higher or lower affinity for BACE1 to initiate APP cleavage, thus exerting a protective or risk effect, respectively [3].

The reported translational results provide robust proof of principle for the pathophysiological and pharmacological model, indicating that reducing the β -cleavage of APP may be a resilience mechanism for AD [3].

With the advent of oral and blood-brain barrier (BBB)-permeable inhibitors, BACE1 has become a central target in several drug AD R&D pipelines. Despite intense pharmacological efforts, all clinical trials so far have been discontinued for futility or signs of cognitive worsening or some systemic toxic effects, thus raising relevant safety and efficacy concerns [7, 8].

One of the most significant issues after a first reflection on discontinued clinical trials is that they did not introduce any direct BACE1 biomarkers for any relevant context of use (CoU), such as proof of mechanism, dose-

finding, and efficacy/safety measures. Implementation of existing BACE1 biomarkers would support the mapping of drug response, optimization of go/no-go decision-making, and mitigation of side effects due to non-specific or too high BACE1 inhibition [3].

Search strategy and selection criteria

The narrative inherent to this review article is based on the authors' knowledge and experience in the field. As such, no systematic literature search was performed.

BACE1 biomarkers in the cerebrospinal fluid (CSF)

BACE1 protein concentrations, probably reflecting levels of gene expression, and rates of enzymatic activity, have been measured in human CSF samples to investigate their diagnostic/predictive values as well as their association with critical pathophysiological alterations of AD, including the amyloid- β pathway, tau pathophysiology, neurodegeneration, and synaptic dysfunction [3].

CSF BACE1 biomarker: diagnostic and predictive performance

The first study analyzed BACE1 CSF concentrations and activity in a pooled cohort of cognitively healthy control

(HC) individuals, clinically diagnosed patients with AD dementia (ADD), and individuals with mild cognitive impairment (MCI) [9].

The authors reported that individuals in which BACE1 both enzymatic activity and protein levels are in the higher ranges showed an increased relative risk of association with the MCI group when compared to HC or ADD individuals. The finding of elevated BACE1 biomarkers in individuals with MCI compared to ADD was discussed in relation to extensive dendritic remodeling and neuronal loss characterizing dementia stages [9].

Zetterberg and colleagues investigated BACE1 activity in a comparable pooled cohort, reporting a statistically significant difference between patients with ADD and HC individuals, but not between ADD patients and a diagnostic group of HC combined with MCI individuals [10]. When further differentiating the group of MCI individuals into ADD converters and non-converters, they found higher mean BACE1 activity in the former subgroup than in the latter one [10]. Such results were corroborated in subsequent studies [11–13].

In a separate study, Pernecky and colleagues assessed BACE1 activity in a population of 342 individuals, including HC, stable MCI, converted MCI individuals, and ADD patients. Contrary to these findings, no significant differences in BACE1 activity were found between the investigated groups. The authors argued that such unexpected results could be partly due to the different used assays, which may have impacted BACE1 activity due to higher intra-assay variability [14].

Likewise, Savage and colleagues found no significant difference regarding BACE1 activity between individuals with HC, MCI, and ADD. In line with the argumentation of Pernecky and colleagues, they hypothesized that the wide inter-subject variability of BACE1 activity along with technical differences of laboratory assays could have limited the analytical standardization and clinical validation of the CSF BACE1 diagnostic candidate biomarker [15].

CSF BACE1 and amyloid- β biomarkers

Studies investigating CSF BACE1 biomarkers and indicators of brain accumulation of A β , including CSF 42-amino acid forms of amyloid- β protein (A β 42) and A β positron emission tomography (A β -AB-PET), do not show consistent results, as some studies did not find any significant association [10, 14, 16, 17], while other studies showed significant correlations [18–20].

A significant association was found in studies that stratified the whole study population according to the clinical diagnosis, reporting positive correlations only in HC individuals and patients with ADDs [18–20], but not in individuals with MCI [18]. There is no clear biological

explanation for the variation of results across different studies but rather a potential methodological issues related to employing different study designs, populations, and assays. The diverging results (i.e., A β 42 is a product of the BACE1 pathway and thus an association was expected) have generated some discussion.

Some authors raised the question of whether A β 42 monomers truly provide comprehensive information on the whole A β pathway that encompasses small aggregation species, oligomers, protofibrils, fibrils, and eventually senile plaques [3, 10]. Effective clearance of A β aggregates will impact the concentrations of A β 42 in CSF as well. Therefore, the whole unfolding of the amyloid- β pathway can account for a non-linear association between BACE1 and A β 42 monomers (see below). Despite the lack of a robust correlation of CSF A β 42 with BACE1 concentrations, a multimodal study showed that CSF BACE1 activity is correlated with a global uptake of the Pittsburgh Compound B PET (PiB-PET) tracer, a radiotracer that binds to the fibrillary component of amyloid plaques [21].

BACE1 CSF parameters correlate, at least in part, with other A β markers. In particular, a strong positive correlation between BACE1 and levels of A β 40, sAPP- α , and sAPP- β has been reported. Interestingly, although sAPP- α is a by-product of the alternative pathways of α -secretase, it negatively correlates with BACE1 [10]. This finding may be explained through collinearity between sAPP- α and sAPP- β that are highly associated with one another or by the fact that both by-product may reflect the rate of APP processing [10].

CSF BACE1 and biomarkers of tau-related pathophysiology

Concerning the association between BACE1 and tau-related pathophysiology, multiple groups—using different methodological approaches and study designs—found a positive correlation between CSF BACE1 biomarkers and CSF tau phosphorylated at threonine181 (t-tau and p-tau, respectively) [22].

Experimental evidence and translational studies can help explain the association between p-tau and the amyloid- β pathway, including the putative upstream role of BACE1. Indeed, injection of A β fibrils into the brains of P301L mutant tau transgenic mice triggers a fivefold increase in NFTs in cell bodies within the amygdala where neurons project to the injection sites [23]. In another study, crossing transgenic mice showing the spread of tau from the entorhinal cortex to other brain regions with APP/PS1 mice showed that cortical amyloid deposition caused a dramatic increase in tau spreading to distal brain regions. Hence, several findings point toward an upstream role of A β , and on an inferring speculative basis BACE1, on tau phosphorylation and neurofibrillary tangle generation by facilitating and promoting the

conversion of tau from a normal to a toxic state, which may enhance A β toxicity via a feedback loop [24, 25]. Such experimental evidence supports the data-driven (biomarker-based) hypothetical model of AD clinical-biological continuum whereby brain accumulation of A β may either facilitate being permissive to spreading of tau pathology that is tightly associated with the clinical evolution of the disease [22].

CSF BACE1 biomarkers and neurodegeneration

CSF BACE1 biomarkers have been investigated in relation to hippocampal volume loss, a biomarker of regional neurodegeneration occurring during early stages of AD.

The only published structural MRI study reported that an increase in CSF BACE1 activity is associated with bilateral decreased hippocampus volume [11]. The interpretation of this finding is unclear; however, it may suggest a BACE1-mediated neurotoxicity. The observed BACE1 activity in CSF inversely correlating with hippocampal volume supports the hypothesis that elevated BACE1 may induce downstream amyloidogenic effects by triggering stepwise neurodegeneration leading to hippocampal atrophy.

Evidence indicating an association between BACE1 and neurodegeneration can be derived from studies reporting positive correlations between CSF BACE1 biomarkers and CSF total tau protein [9–12, 14–20, 26], a surrogate marker of axonal damage and neuronal loss. It is conceivable that BACE1 is released into CSF by degenerating neurons and that the concentrations may correlate with the severity of neurodegeneration and the progression of synaptopathy and neuronal loss. Dysregulation of synaptic BACE1 functions may account in part for a non-amyloidogenic impact on synaptic homeostasis.

Beyond any preliminary data-driven and knowledge-based consideration, it must be outlined that major parts of evidence regarding the association between BACE1 and tau biomarkers in CSF have been studied cross-sectional. Longitudinal observational studies are needed to investigate the spatial-temporal relationship between BACE1 biomarker expression, gene expression levels and activity, and neurodegeneration.

By contrast, the results of the phase 3 trial of verubecestat (12 or 40 mg/day) conducted in mild-to-moderate AD patients (EPOCH, ClinicalTrials.gov NCT01739348) showed significant BACE1 inhibition correlating with decreased hippocampal volumes [3]. Although a univocal interpretation of this trial result is challenging, it has been hypothesized that there is an over inhibition of BACE1 regulation of synaptic substrates and or physiological functions of A β species, essential also for hippocampal homeostasis [3] (see Fig. 1 for more details about the amyloidogenic pathway). Follow-up analysis is needed in

BACE1 inhibitor trials to ascertain whether this hippocampal effect is related to the cognitive worsening reported in some studies and whether it may be reversible.

CSF BACE1 and synaptic biomarkers

Two studies investigated the association between CSF BACE1 and biomarkers of synaptic dysfunction. De Vos and colleagues analyzed CSF BACE1 and neurogranin (NGR)—a dendritic protein proposed as a biomarker of hippocampal synaptic impairment [27]—in a pooled cohort of HC individuals HC and positive A β biomarkers individuals diagnosed with MCI or ADD [18]. Despite no significant inter-group differences, they found that the NGR/BACE1 ratio differentiates both the MCI and ADD diagnostic groups from the HC individuals group with good accuracy [18]. The NGR/BACE1 ratio also showed potential prognostic value since individuals with higher concentrations had a more severe cognitive decline at follow-up [18].

In agreement with De Vos and colleagues, a recent report showed that NGR/BACE1 ratio levels are (i) elevated in both individuals with subjective cognitive decline (SCD) and MCI compared to HC individuals, (ii) associated with smaller hippocampal and amygdala volumes, and (iii) correlate with worse baseline and longitudinal cognitive performance [28].

A recent study investigated a broad set of candidate biomarkers tracking distinct pathophysiological processes in patients with AD and reported a positive cross-sectional correlation between BACE1 CSF concentrations and NGR [20]. Interestingly, another group reported that the CSF NGR/BACE1 ratio, along with core AD biomarkers, displays good accuracy to distinguish between depression with cognitive impairment and AD dementia [29].

Given the established neurobiological overlap among depression, MCI, and AD, the NGR/BACE1 ratio may represent a suitable clinical tool for the AD diagnostic workup [30]. Further prospective longitudinal studies are needed to understand the role of the NGR/BACE1 ratio as a biomarker to improve the classification between non-neurodegenerative forms of MCI, including but not exclusively depression, and early AD.

The understanding of the relationship between BACE1 and synaptic homeostasis remains an unmet objective. There is evolving experimental evidence, i.e., data from conditional deletion of BACE1 in mouse models, that points at BACE1 as a molecular orchestrator of hippocampal synaptic remodeling at the dendritic and axonal level [3, 7]. Arguably, BACE1 overactivation may excessively accelerate synaptic turnover until it triggers downstream detrimental pathways resulting in synaptic damage. The generated hypothesis is further supported by studies investigating the association between BACE1 and other candidate biomarkers of neurodegeneration and synaptic loss.

CSF BACE1 biomarkers and the APOE $\epsilon 4$ allele

Ewers and colleagues reported an association of the apolipoprotein E (*APOE*) $\epsilon 4$ allele with increased BACE1 activity in both patients with ADD and subjects with MCI compared to HC individuals [31]. This finding is in agreement with experimental models of AD, indicating increased activity of BACE1 in individuals carrying the *APOE* $\epsilon 4$ allele [32]. It is unclear whether this correlation is induced by *APOE* $\epsilon 4$ allele in promoting A β deposition, which subsequently can induce increase of BACE1 activity. Consistently, in-human neuropathological studies show higher concentrations of BACE1 in HC individuals or ADD patients, carrying the *APOE* $\epsilon 4$ allele [33]. Two independent studies did not show any association between the *APOE* genotype and BACE1 concentrations or rates of activity. Of note, the two studies differed significantly with each other and compared to other investigations, in terms of experimental design, including the assay utilized [15, 16]; see the section “Potential explanation of controversial results in CSF (and blood-based) BACE1 studies” for a more in-depth argumentation.

CSF-based studies provide evidence that BACE1 biomarkers, both protein concentrations and enzymatic activity, support further analytical and clinical investigations in patients with AD to investigate their potential as candidate biomarkers suitable for clinical practice (i.e., early diagnosis, prediction, and progression) and pharmacological trials targeting BACE1 (i.e., target engagement and efficacy response, among others). Further studies with longer follow-up, standardized pre-analytical procedures, and analytical protocols are required to address the open questions based on conflicting study data.

BACE1 blood-based biomarkers

BACE1 biomarkers in plasma

While BACE1 is primarily expressed in the CNS, the protease is expressed in platelets, leukocytes, and is circulating as a soluble protein in the plasma.

For different CoU, blood-based biomarkers provide unique opportunities and decision-making tools in clinical trial programs. Blood-based biomarkers have numerous advantages, i.e., they are widely accessible and minimally invasive and are more time- and cost-effective for healthcare systems compared with CSF. In particular, they are appropriate tools to inform biomarker-guided medicine applied to individuals with preclinical AD [27].

Both BACE1 activity and protein concentrations in blood (mostly plasma but in some cases serum) are significantly increased in MCI individuals or patients with ADD compared to HC individuals, with a trend across different disease stages reflecting the direction of expression of the reported CSF biomarkers [34–36].

Shen and colleagues reported a study in which both blood BACE activity and protein concentration were measured and explored in parallel with correlations of CSF AD core biomarkers [36]. The population included individuals with ADD, HC, MCI converters to AD, and MCI stable at follow-up. They showed that BACE1 activity was elevated in both individuals with ADD and MCI converters when compared to stable MCI or HC individuals; at the same time, BACE1 protein concentrations were significantly increased in individuals with ADD compared to HC or stable MCI, while BACE1 concentrations in converter MCI individuals were significantly elevated compared to subjects with stable MCI, but not compared to the HC group [36]. BACE1 activity was positively correlated with CSF t-tau protein and negatively correlated with CSF A $\beta 42$, further supporting a link between the plasma biomarker and brain AD pathology [36].

A recent plasma-based study tested associations between plasma BACE1 concentrations and the degree of cerebral accumulation of A β in a cohort of HC with subjective memory complaints (SMC), a condition associated with increased risk for AD. For this objective, brain accumulation of A β was investigated using A β positron emission tomography (PET) imaging (A β -PET) [37] showing, for the first time, that plasma BACE1 concentrations impact the level of brain A β in individuals with SMC [37].

The same study further investigated the question of whether other relevant biological factors, such as sex, besides the *APOE* $\epsilon 4$ allele and age, may affect plasma BACE1 concentrations [37]. They found highly significantly increased baseline and longitudinal mean concentrations of BACE1 in women compared to men, irrespective of age and time [37]. These results indicate a potential sexual dimorphism in plasma BACE1 concentrations, in agreement with experimental evidence about the role of estradiol in the control of BACE1 expression [38].

BACE1 assessed in platelets

Studies conducted in platelets show an increase of BACE1 protein concentrations and higher activity rates in individuals with ADD and or MCI compared to HC [39–41]. One study did not find differences between individuals with HC and MCI, while Decourt and colleagues found decreased BACE1 protein concentrations in individuals with ADD when compared to HC [42].

Another platelet-based study reported that patients with ADD treated with a stable 6-month dose of donepezil, but not cognitively healthy controls, showed downregulation of BACE1 gene expression in platelets [43]. Given the consistency between the study results and AD pathophysiology, including BACE1 overexpression, the authors argued that BACE1 platelet levels should be investigated to ascertain whether they might

represent an additional exploratory outcome measure to employ in BACE1 inhibitor trials.

BACE1 transcriptomic studies in blood

Besides assessing enzymatic biomarkers, molecular biology investigations of BACE1 have used blood samples. In particular, three studies explored the variability of BACE1 gene expression in AD. In 2019, Wongchitrat and colleagues measured the rate of BACE1 mRNA expression in peripheral leukocytes and showed significantly higher mRNA levels in ADD patients compared to HC individuals [44]. In the same year, Vakilian and colleagues obtained similar results, and in their extension study, they assessed BACE1 concentration in using the same blood samples as in the genetic research. Although they found increased concentrations of BACE1 in ADD patients, they did not observe any correlations between BACE1 and mRNA expression [45].

A different approach was chosen by Fotuhi and colleagues, which evaluated circulating long noncoding RNA (lncRNA) related to the BACE1 gene (BACE1-AS) in plasma and plasma-derived exosomes. The lncRNA BACE1-AS is believed to improve BACE synthesis, via mRNA stabilization [46].

They did not observe any difference in exosome lncRNAs; however, they found that lncRNA BACE1-AS plasmatic concentration was significantly lower in mild ADD compared to HC individuals, differentiating HC individuals from mild ADD with good sensitivity and specificity [46].

Potential explanation of controversial results in CSF (and blood-based) BACE1 studies

Inconsistent results complicate clinical validation and qualification of BACE1-related biomarkers and their potential integration into the evolving AD biomarker matrix. Existing discrepancies need to be carefully scrutinized to see whether methodological issues, rather than biological implications, may determine differences.

First, as indicated by human neuropathological and experimental models of aging and AD, BACE1 gene expression and rates of activity may vary throughout AD progression. In this context, Rosen and colleagues found that BACE1 activity was significantly increased in AD patients with mild dementia compared to patients at more severe stages [12]. A recent study showed that BACE1 biomarker candidates are significantly increased in individuals with MCI, but not with ADD, when compared with the HC group [13]. Therefore, it is likely that the disease stage of the individual patient may influence BACE1 concentration and activity, and thus, clinical heterogeneity of included individuals may have neutralized inter-group differences.

Second, since BACE1 had been proposed as a potential AD-specific biomarker, a considerable part of the reported studies, in particular the older publications, were performed in clinically, but not biologically, characterized study populations. Hence, it is not possible to rule out that the enrollment of individuals with non-AD pathophysiology may have biased the data and created conflicting outcomes.

Most of the recent studies used CSF core biomarkers of AD, where BACE1 correlated indeed with A β and tau markers. In particular, Mulder and colleagues found that BACE1 activity was increased in individuals showing characteristic AD biological features compared to individuals with negative AD biomarkers [16], while Alexopoulos and colleagues showed significantly decreased CSF BACE1 activity in individuals with MCI without AD pathophysiology compared to patients with MCI due to AD [13].

Third, some studies did not report any association between CSF BACE1 biomarkers and CSF A β 42 raising questions about the potential of BACE1 biomarkers in AD pharmacological trials. However, given the tendency of A β 42 monomers of aggregating into oligomers and fibrils and preliminary evidence that BACE1 is associated with A β -PET radiotracer uptake, we suggest further studies using multimodal outcome measures such as PET tracers and emerging CSF candidates for the assessment of different A β species, including A β -oligomers and protofibrils [47–49].

Fourth, conflicting data may be partly explained by sexual dimorphism in BACE1 biology. Indeed, in a study conducted in patients suffering from bipolar disorder, a sex-based dimorphism in BACE1 gene expression levels was reported, with men displaying upregulation of BACE1 expression [50].

Furthermore, sexual dimorphism was reported in males exhibiting higher *BACE1* expression compared to females with schizophrenia and HC individuals [51].

However, female sex-biased dimorphism may exist regarding BACE1 concentrations in cognitively healthy individuals at risk for AD [37]. This finding is in line with most of the experimental evidence—mouse models of aging and AD—that indicates that intracellular effects of estrogens induce upregulation of BACE1 gene expression levels [32, 52], confirming the widely accepted notion that women bear higher vulnerability to AD [53, 54].

Apart from the necessity to elucidate in humans whether male or female sex may be a determinant of BACE1 gene expression in AD—where other genetic/biological factors may synergize with sex hormones or act independently to upregulate BACE1—evidence of sexual dimorphism in BACE1 biology may be relevant for clinical BACE1 inhibitor trial outcomes.

If a BACE1 sexual dimorphism was corroborated, sex-related outcome analyses and comparative active treatment dose-finding studies should be taken into account.

Lastly, the abovementioned CSF studies assessed different BACE1 biomarkers and some studies evaluated BACE1 enzymatic activity [10, 14, 15], while others investigated either BACE1 protein levels [18] or both biomarkers.

It should be acknowledged, however, that several *in vitro* and animal data point to a non-linear relationship between the levels of gene expression and rates of enzyme activity that is highly influenced by post-translational modifications [55]. Indeed, experimental studies indicate that BACE1 activity significantly increases over time while its expression levels are less likely to be altered during cognitively healthy aging as well as in the presence of AD-related cognitive decline [3].

Methodological and technological challenges

Inconsistent results in fluid biomarker studies can derive from methodological differences.

Pre-analytical factors such as the sample collection, processing, and storage protocol, as well as analytical factors including sample handling and immunoassays used, are likely the most relevant determinants of the inter-study variability in terms of results (see Tables 1 and 2).

Regarding pre-analytical factors, besides those that concern AD CSF biomarkers in general, a recent study conducted in cell lines and iPSC-derived neurons reported that 7 of the 8 BACE1 inhibitors evaluated show increased BACE1 protein concentrations [66]. A thorough pre-analytical evaluation is required to understand better the effect of BACE1 inhibitors in BACE1 biomarker assays [66].

Regarding the assessment of BACE1 biomarkers, poor specificity (i.e., other enzymatic activities may contribute to the signal) of activity-based assays may represent a plausible explanation for the observed differences [39]. In particular, peptide-based activity assays show questionable reliability for measuring BACE1 activity. To our knowledge, one of the most robust assays used is reported by Sinha and colleagues in 1999, utilizing membrane-bound substrates for measuring BACE1 activity [67]. Specifically, they purified BACE1 activity to homogeneity from human brains using a substrate analog inhibitor of the enzyme activity [67].

Additionally, antibody-based assays used in the described investigations differ in the binding site and epitope recognition. It is plausible to infer that distinct forms of BACE1 may have been explored in previous studies.

Moreover, recent research suggests that the existence of multiple enzyme isoforms could affect the correct estimation of BACE1 concentration. It is worth noting that some spliced forms do not have APP-cleaving activity and that it is not known which (and whether) specific forms vary in AD [55]. The same study shows that other enzymes, detectable in CSF, such as cathepsin B, meprin β , and BACE2, could exert β -secretase activity.

Some studies employed assays based on polyclonal antibodies [34, 68], which do not support the reproducibility of the assays on the longer term (cfr. long-term supply of antibodies with steady characteristics, lot-to-lot consistency of the assay, etc.). ELISA-based platforms, however, may have non-specificity due to weak polyclonal antibodies, and this can cause a discrepancy in data variation between different groups. The stability of BACE1 protein concentrations measured by sandwich ELISA shows limited change under standard storage and freeze/thaw conditions [69].

It will be crucial to develop and standardize the most appropriate methodologies, to understand the corresponding readout, and eventually focusing on CoU qualification to establish the potential role of BACE1 as an AD biomarker. While the activity-based measurements encounter low specificity, analyzing protein concentrations should consider the various known BACE1 protein isoforms also generated by post-translational modifications, membrane association, and truncated fragments [65]. Indeed, post-translational modifications influence the rate of activity of BACE1, thus accounting for a non-linear association with BACE1 gene expression levels [3].

A relatively easy step forward to increase specificity and analytical robustness of BACE1 biomarker assessment is to use immunoassays with well-defined antibodies. In this regard, an immunoassay based on two monoclonal antibodies (mAbs) has been established. It consists of a clone 10B8 that recognizes BACE1 within its extracellular, active domain (aa46-240) (aa numbering of human BACE1) and, as a complementary monoclonal antibody, clone 5G7, recognizing cleaved, non-membrane-bound BACE1 via a conformational epitope (aa299-312, a helical region of BACE1, and a free C-terminal Q386 end). In the first explorative study with this immunoassay, significant differences were observed in the CSF from AD patients compared to control individuals [26].

Although still at an initial stage, innovative molecular imaging for BACE1 assessment is under development. In line with the high-affinity of BACE1 inhibitors, recent efforts to develop brain-penetrant PET radiotracers, such as the highly potent selective aminothiazine inhibitor, PF-06684511 have been made [70].

Table 1 BACE1 CSF-based biomarkers

BACE1 activity measured (method)	Antibody	Substrate	Clinical study	Result
Sandwich ELISA	Anti-BACE1 Ab SECB1&2 [56, 57] Anti-BACE1 Ab B280 and anti-BACE1 monoclonal Ab (R&D Systems Inc) [58]	Synthetic fluorescence substrate—containing the BACE1 cleavage site	Zhong et al. [59]	Significant elevation of BACE1 levels in MCI and AD Strong and significant correlation of BACE1 activity with BACE1 protein and A β peptide level
Solution-based detection of CSF BACE activity	Polyclonal NF neopeptide-specific Ab [60]	Biotinylated BACE1 substrate	Zetterberg et al. [10]	Significant differences in BACE1 activity between MCI, AD, and HC Positive correlation between BACE1 activity, CSF t-tau, and A β 40 levels in the MCI and AD
Sandwich ELISA	Anti-BACE1 Ab SECB1&2 [56, 57] Anti-BACE1 Ab B280 and anti-BACE1 monoclonal Ab (R&D Systems Inc) [58]	Synthetic fluorescence substrate—containing the BACE1 cleavage site [59]	Ewers et al. [11]	BACE1 activity and protein levels were significantly increased in AD compared to healthy HC Increase in CSF BACE1 activity in AD Increased activity associated with increased CSF t-tau but not A β 42 in AD
2 step-solution-based detection of CSF BACE activity	Anti-NF c-terminal neopeptide polyclonal Ab	Biotinylated peptide substrate (bBACE (aa1–460)) and Sapphire-II Enhancer substrate	Pernecky et al. [14] Savage et al. [15]	No significant difference in BACE1 levels between HC, MCI, and AD
ELISA	mAbs ADx401 (clone 5G7) and 10B8 and mAb ADx402 (clone 10B8F1) biotinylated with peroxidase [26]		De Vos et al. [18]	Significant correlation of ratio of CSF neurogranin trunc P75/BACE1 between HC, MCI, and AD
Solution-based assay	BACE1 Activity Assay Kit (Sigma CS1060)	BACE1 Activity Assay Kit (Sigma CS1060)	Mulder et al. [16]	No significant differences in BACE1 levels between MCI, AD, and HC
SignalClimb technology	Time-resolved fluorescence activity [61]	Synthetic TruePoint BACE1 substrate	Tsolakidou et al. [62]	Positive correlation between BACE1 activity and SORL1 in CSF t-tau and sAPP β levels in AD
Sandwich ELISA	Anti-BACE1 monoclonalAbs (mAbs) 5G7 and 10B8 [26]		Timmers et al. [19]	CSF BACE1 correlated positively with age (weak) and with A β 37 (strong), sAPP-total, p-tau181 (strong)
Commercially available assays	Euroimmun, Luebeck, Germany		Schaefferbeke et al. [20]	Correlation of BACE1 activity with brain volumes and A β load in regions typically involved early in AD
SignalClimb technology	Time-resolved fluorescence activity [61]	Synthetic TruePoint BACE1 substrate	Grimmer et al. [21]	Strong association between BACE1 activity and in vivo A β pathology in brain regions close to the ventricles
Solution-based detection of CSF BACE activity	Polyclonal NF neopeptide-specific Ab [59]	Biotinylated BACE1 substrate	Rosén et al. [12]	BACE1 correlated slightly with sAPP α , sAPP β , and A β 40
2 step ELISA	Time-resolved fluorescence activity [61]	Synthetic TruePoint BACE1 substrate	Alexopoulos et al. [63]	No significant difference in BACE1 activity between AD and HC or MCI while BACE1 activity was significantly higher in MCI-AD compared to both HC

Table 1 BACE1 CSF-based biomarkers (Continued)

BACE1 activity measured (method)	Antibody	Substrate	Clinical study	Result
Solution-based detection of CSF BACE activity	Fluorescence-based detection in the presence of inhibitor Calbiochem	Synthetic peptide substrates containing the BACE 1 cleavage site	Ewers et al. [31]	CSF BACE1 activity significantly increased in MCI compared to AD. No significant difference between AD and HC
BACE activity fluorometric assay kit	K360-100, BioVision, Milpitas, CA, USA		Hou et al. [32]	BACE1 activity significantly increased in the hippocampus of ApoE4/3xTg mice especially in females
Sandwich ELISA	Used anti-BACE1 ectodomain MAB9311 (R&D Systems) and detection with rabbit anti-BACE1 N-terminus B0681 (Sigma–Aldrich) Abs [64]		Decourt et al. [42]	BACE1 levels 12% lower in the AD frontal cortex compared to HC 6.5% decrease in the temporal cortex
Solution-based assay	Used modified procaspase-3 as detection enzyme	Caspase substrate AspGluValAsp-p-nitroanilide	Verheijen et al. [65]	Assay detects BACE1 activity in extracts of human brain tissue and CSF

Abbreviations: CSF cerebral spinal fluid, MCI individuals with mild cognitive impairment, AD patients with Alzheimer's disease dementia, HC cognitively healthy individuals, ELISA enzyme-linked immunosorbent assay, Ab antibody, *t-tau* total peptide tau protein, A β amyloid beta, BACE1 beta secretase1

Table 2 BACE1 blood-based biomarkers

BACE1 activity (method)	Antibody	Substrate	Clinical study	Result
ELISA	Anti-BACE1 Ab SECB1&2 [56, 57]	Synthetic peptide substrates containing the β -cleavage site (Calbiochem, EMD, Gibbstown, NJ, USA) [59]	Shen et al. [36]	Plasma BACE1 activity significantly increased by 53.2% in MCI and by 68.9% in AD compared to HC
ELISA	Anti-BACE1 Ab SECB1&2 [56, 57]	Synthetic substrate-C-terminally labeled with the fluorescent, Luciferase Yellow, and N-terminally labeled with the quenching, Dabsyl	Cervellati et al. [35]	Increased BACE1 activity in serum of AD
ELISA	Biotinylated detector mAb, diluted in a buffer adapted for the plasma matrix	Kit-based assay EQ 6541–9601-L; Euroimmun AG, Lübeck, Germany [18]	Vergallo et al. [37]	Plasma BACE1 significantly higher in women than in men in cognitively healthy individuals at clinical risk for AD
Solution-based platelet BACE1 assay		Fluorogenic substrate (Calbiochem, BACE1 substrate I)	Johnston et al. [39]	17% increase in platelet membrane BACE1 activity in AD compared to HC
Solution-based platelet BACE1 assay		Fluorescence-quenching substrate (Calbiochem, Merck, Darmstadt, Germany)	Bermejo-Bescós et al. 2013 [40]	No significant difference in BACE1 activity between MCI and HC
Solution-based platelet BACE1 assay		Fluorogenic substrate (Sigma A1472 or Bachem M2465)	Wongchitrat et al. [44]	Baseline platelet membrane BACE1 activity not significantly different between MCI vs HC
Immunoassay		Immunoassay kit (CUSABIO, USA)	Vakilian et al. [45]	Elevated plasma levels of BACE1 in AD vs HC
RNA expression analysis		Real-time quantitative RT-PCR	Ghafouri-Fard et al. [50]	BACE1 levels significantly high in bipolar disorder
RNA expression analysis		Real-time quantitative RT-PCR	Nafisi-Far et al. [51]	BACE1 levels significantly high in schizophrenia

Abbreviations: CSF cerebral spinal fluid, MCI individuals with mild cognitive impairment, AD patients with Alzheimer's disease dementia, HC cognitively healthy individuals, ELISA enzyme-linked immunosorbent assay, Ab antibody, *t-tau* total peptide tau protein, A β amyloid beta, BACE1 beta secretase1

As of other AD PET biomarkers, PET BACE1 ligands can investigate regional patterns of BACE1 activity and monitor BACE1 inhibitor regional brain effects.

In summary, pre-analytical and analytical protocols for BACE1, as well as other biomarkers for AD, should be harmonized and then standardized at a global scale to drastically reduce inter-study and longitudinal variability and eventually speed up the validation and qualification process of BACE1 biomarker candidates. The validation process will be facilitated not only by internationally accepted general requirements for the competence of testing and calibration laboratories but also by recently proposed standard operating procedures (SOPs) for Alzheimer's biomarkers, including BACE1 (see Table 3 for specific information).

Conclusions: challenges and perspectives

A potential explanation for the observed cognitive worsening in some of the previously reported late-stage BACE1 inhibitor trials may be related to insufficient APP substrate specificity. Other physiologically relevant BACE1 substrates may have been inhibited; some of these are involved in neuroplasticity, repair, and synaptic pathways. It may also be possible that BACE1-mediated APP processing could have been inhibited too strongly impairing physiological APP turnover or alternative APP-processing pathways may have been induced. More extensive research is needed to answer these questions.

Non-clinical, translational studies have shown that BACE1 activity is a relevant facilitator of axonal

sprouting, dendritic remodeling, and synaptic plasticity, through both amyloidogenic and non-amyloidogenic pathways [3, 7]. In this regard, complete suppression of BACE1 enzymatic activity may substantially impair adult hippocampal neurogenesis [72, 73], which is a crucial mechanism for hippocampal synaptic plasticity and essential for memory formation and learning [72, 73].

Experimental studies in adult conditional BACE1 knockout mice indicated that pharmacological BACE1 inhibition might disrupt the organization of axonal pathways in the hippocampus [3]. Regarding peripheral toxicity, most of the BACE1 inhibitors block the activity of BACE2 as well, a close homolog of BACE1, which may cause additional unwanted on-target side effects.

Among several open scientific issues, we highlight the question of whether the negative correlation between CSF BACE1 enzymatic activity and degree of hippocampal atrophy may be primarily induced by the BACE1 downstream amyloidogenic effects or on affected synaptic pathways as well. The interrelation between BACE1 and progressive neurodegeneration deserves further investigation. Neurodegeneration biomarker panels, including tau and NFL proteins, provide partially differential information of related pathophysiological processes [74].

There is evidence of a complex interaction between the amyloidogenic pathway and other pathomechanistic alterations of AD, including neuroinflammation. TNF receptor (TNFR1) knockout mice show decreased A β peptides and cerebral accumulation of amyloid plaques

Table 3 Proposed stepwise validation path for Alzheimer's disease biomarkers

Parameter	Definition
Robustness	The ability of a method to remain unaffected by small variations in method parameters
Precision	The closeness of agreement between independent test results obtained under stipulated conditions
Trueness	The closeness of agreement between the average value obtained from an extensive series of test results and an accepted reference value
Uncertainty	A parameter associated with the result of a measurement that characterizes the dispersion of the values could reasonably be attributed to the measurand
Limits of quantification	Highest and lowest concentrations of analyte that have been demonstrated to be measurable with acceptable levels of precision and accuracy
Dilutional linearity	Dilutional linearity is performed to demonstrate that a sample with a spiked concentration above the ULOQ can be diluted to a concentration within the working range and still give a reliable result
Parallelism	Relative accuracy from recovery tests on the biological matrix or diluted matrix against the calibrators in a substitute matrix
Recovery	The recovery of an analyte in an assay is the detector response obtained from an amount of the analyte added to and extracted from the biological matrix, compared to the detector response obtained for the true concentration of the analyte in the solvent
Selectivity	The ability of the bioanalytical method to measure and differentiate the analytes in the presence of components that may be expected to be present
Sample stability	The chemical stability of an analyte in a given matrix under specific conditions for given time intervals

Note: A committee, within the international research framework BIOMARKAPD, recently convened to propose the ten key requirements to fulfill within a step-by-step validation process. The BIOMARKAPD project aims for the standardization of biomarker measurements for AD and Parkinson's disease (PD), including pre-analytical and analytical procedures, assay validation, and development of reference measurement procedures (RMP) and certified reference materials (CRM) for harmonization of results across assay formats and laboratories. The table captures stepwise standard operating procedures (SOP)

Table adapted from [71]

through regulation of BACE1 gene expression via the nuclear factor κ B (NF- κ B) pathway [75, 76].

Addressing these scientific questions will provide key pathophysiological insights and facilitate the implementation of standardized drug-biomarker co-development programs that are necessary to achieve successful BACE1 targeting strategies for precision medicine.

BACE1, either in CSF or blood and either activity or protein concentration, does not show a remarkable performance as a clinical diagnostic or pathognomonic AD biomarker. However, whether BACE1 biomarkers could increase diagnostic accuracy if combined with AD core biomarkers has been poorly investigated. Given the preliminary evidence about the association between BACE1 and A β biomarkers as well as neurodegeneration biomarkers (namely hippocampal volumes and t-tau) and synaptic biomarkers (neurogranin), we support the investigation of BACE1 parameters in combination with the AD core biomarker panel, across different matrixes such as CFS and blood, to assess diagnostic performance in the AD continuum (preclinical, prodromal, dementia stages). Association studies indicate that BACE1 biomarkers may be useful for COU in a clinical trial setting, including proof of mechanism, treatment response, and safety assessment in clinical trials, as well as COU in a clinical practice setting such as prognostic evaluation in MCI individuals. From a therapeutic perspective, BACE1 inhibitors dosing could be personalized to engage targets based on direct concentrations and activity rate measurements from individual bodily fluids [27, 74]. Furthermore, the reduction of cleavage products, such as sAPP β , or enriched alternatively processed peptides such as A β 5-X, which are correlated to BACE1 inhibition, could be used to monitor target engagement and optimize efficacy [15, 77]. Future investigations using combinatorial strategies and biomarker-guided or personalized dose selection may allow the application of lower doses with an optimized specificity for BACE1 over BACE2 [3].

Developing assays for the analysis of A β species, including bioactive oligomers, provide a more profound understanding of human pathophysiology and the relationship between BACE1 and elements of the downstream amyloid pathway, from A β species supporting synaptic and homeostatic functions to more bioactive and toxic species [3, 78].

While experimental studies foster knowledge generation of BACE1's complex biology, including synaptic homeostasis, BACE1 fluid biomarker development still needs to transition through the validation and standardization process with harmonized pre-analytical and analytical protocols.

From a pharmacological standpoint, BACE1 biomarkers are expected to be essential components of drug-biomarker co-development programs supporting

successful outcome generations and lowering drug attrition rates in pipelines targeting BACE1. From a medical practice standpoint, liquid biopsy with first availability of CSF followed by maturation of blood-based BACE1 biomarkers [79] may expand the current descriptive A/T/N classification system into developing a comprehensive and integrative biological staging model of AD.

Acknowledgements

HH and AV are employees of Eisai Inc.

HZ is a Wallenberg Scholar supported by grants from the Swedish Research Council (#2018-02532), the European Research Council (#681712), Swedish State Support for Clinical Research (#ALFGBG-720931), and the UK Dementia Research Institute at UCL.

KB is supported by the Swedish Research Council (#2017-00915), the Alzheimer Drug Discovery Foundation (ADDF), USA (#RDAPB-201809-2016615), the Swedish Alzheimer Foundation (#AF-742881), Hjärnfonden, Sweden (#FO2017-0243), and the Swedish state under the agreement between the Swedish government and the County Councils, and the ALF-agreement (#ALFGBG-715986).

Contributors to the Alzheimer Precision Medicine Initiative-Working Group (APMI-WG):

Mohammad AFSHAR (France), Lisi Flores AGUILAR (Canada), Leyla AKMAN-ANDERSON (USA), Joaquín ARENAS (Spain), Jesús ÁVILA (Spain), Claudio BABILONI (Italy), Filippo BALDACCIO (Italy), Richard BATRLA (Switzerland), Norbert BENDA (Germany), Keith L. BLACK (USA), Arun L.W. BOKDE (Ireland), Ubaldo BONUCCELLI (Italy), Karl BROICH (Germany), Francesco CACCIOLA (Italy), Filippo CARACI (Italy), Giuseppe CARUSO (Italy), Juan CASTRILLO (Spain), Enrica CAVEDO (France), Roberto CERAVOLO (Italy), Patrizia A. CHIESA (France), Massimo CORBO (Italy), Jean-Christophe CORVOL (France), Augusto Claudio CUELLO (Canada), Jeffrey L. CUMMINGS (USA), Herman DEPYPERE (Belgium), Bruno DUBOIS (France), Andrea DUGGENTO (Italy), Enzo EMANUELE (Italy), Valentina ESCOTT-PRICE (Wales), Howard FEDEROFF (USA), Maria Teresa FERRETTI (Switzerland), Massimo FIANDACA (USA), Richard A. FRANK (USA), Francesco GARACI (Italy), Hugo GEERTS (USA), Ezio GIACOBINI (Switzerland), Filippo S. GIORGI (Italy), Edward J. GOETZL (USA), Manuela GRAZIANI (Italy), Marion HABERKAMP (Germany), Marie-Odile HABERT (France), Britta HÄNISCH (Germany), Harald HAMPEL (France), Karl HERHOLZ (England), Felix HERNANDEZ (Spain), Bruno P. IMBIMBO (Italy), Dimitrios KAPOGIANNIS (USA), Eric KARRAN (USA), Steven J. KIDDLE (USA), Seung H. KIM (South Korea), Yosef KORONYO (USA), Maya KORONYO-HAMAOU (USA), Todd LANGEVIN (USA), Stéphane LEHÉRICY (France), Pablo LEMERCIER (France), Simone LISTA (France), Francisco LLAVERO (Spain), Jean LORENCEAU (France), Alejandro LUCÍA (Spain), Dalila MANGO (Italy), Mark MAPSTONE (USA), Christian NERI (France), Robert NISTICÒ (Italy), Sid E. O'BRYANT (USA), Giovanni PALE RMO (Italy), George PERRY (USA), Craig RITCHIE (Scotland), Simone ROSSI (Italy), Amira SAIDI (Italy), Emiliano SANTARNECCHI (USA), Lon S. SCHNEIDER (USA), Olaf SPORNS (USA), Nicola TOSCHI (Italy), Pedro L. VALENZUELA (Spain), Bruno VELLAS (France), Steven R. VERDOONER (USA), Andrea VERG ALLO (France), Nicolas VILLAIN (USA), Kelly VIRECOULON GIUDICI (France), Mark WATLING (England), Lindsay A. WELIKOVITCH (Canada), Janet WOODCOCK (USA), Erfan YOUNESI (France), José L. ZUGAZA (Spain).

Alzheimer Precision Medicine Initiative (APMI)

<https://www.apmiscience.com/>

Authors' contributions

Author contribution: HH conceptualized the study, wrote the article, and provided a critical review of the literature; SL contributed to the conceptualization of the study and the writing of all sections of the article and supported the review of the literature; EVM provided expert input on fluid biomarkers of BACE1 and the related methodology; HZ provided expert input on existing data-driven studies involving fluid biomarkers of BACE1 and other Alzheimer's disease biomarkers; FSG contributed to the writing of a part of the article and supported the review of the literature; AG contributed to the writing of a part of the article and supported the review of the literature; KB provided expert input on existing data-driven studies involving fluid biomarkers of BACE1 and other Alzheimer's disease; FC contributed to the writing of a part of the article and supported the review of the literature; BD provided the table and contributed to the review of the literature; RY

contributed to the writing of the article and provided expert input on fluid biomarkers of BACE1 and the related methodology; AV conceptualized the study, wrote the article, and provided a critical review of the literature. The author(s) read and approved the final manuscript.

Authors' information

Nothing to declare.

Funding

Not applicable

Availability of data and materials

Not applicable

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Competing interests

HH is an employee of Eisai Inc. This work has been performed during his previous position at Sorbonne University, Paris, France. At Sorbonne University, he was supported by the AXA Research Fund, the "Fondation partenariale Sorbonne Université," and the "Fondation pour la Recherche sur Alzheimer," Paris, France. HH serves as Senior Associate Editor for the journal *Alzheimer's & Dementia* and does not receive any fees or honoraria since May 2019; before May 2019, he had received lecture fees from Servier, Biogen, and Roche; research grants from Pfizer, Avid, and MSD Avenir (paid to the institution); travel funding from Eisai, Functional Neuromodulation, Axovant, Eli Lilly and company, Takeda and Zinfandel, GE Healthcare, and Oryzon Genomics; and consultancy fees from Qynapse, Jung Diagnostics, Cytos Ltd., Axovant, Anavex, Takeda and Zinfandel, GE Healthcare, Oryzon Genomics, and Functional Neuromodulation; and participated in scientific advisory boards of Functional Neuromodulation, Axovant, Eisai, Eli Lilly and company, Cytos Ltd., GE Healthcare, Takeda and Zinfandel, Oryzon Genomics, and Roche Diagnostics.

He is a co-inventor in the following patents as a scientific expert and has received no royalties:

- In Vitro Multiparameter Determination Method for The Diagnosis and Early Diagnosis of Neurodegenerative Disorders Patent Number: 8916388.
- In Vitro Procedure for Diagnosis and Early Diagnosis of Neurodegenerative Diseases Patent Number: 8298784.
- Neurodegenerative Markers for Psychiatric Conditions Publication Number: 20120196300.
- In Vitro Multiparameter Determination Method for The Diagnosis and Early Diagnosis of Neurodegenerative Disorders Publication Number: 20100062463.
- In Vitro Method for The Diagnosis and Early Diagnosis of Neurodegenerative Disorders Publication Number: 20100035286.
- In Vitro Procedure for Diagnosis and Early Diagnosis of Neurodegenerative Diseases Publication Number: 20090263822.
- In Vitro Method for The Diagnosis of Neurodegenerative Diseases Patent Number: 7547553.
- CSF Diagnostic In Vitro Method for Diagnosis of Dementias and Neuroinflammatory Diseases Publication Number: 20080206797.
- In Vitro Method for The Diagnosis of Neurodegenerative Diseases Publication Number: 20080199966.
- Neurodegenerative Markers for Psychiatric Conditions Publication Number: 20080131921.

SL received lecture honoraria from Roche and Servier.

AV is an employee of Eisai Inc. This work has been performed during his previous position at Sorbonne University, Paris, France. He does not receive any fees or honoraria since November 2019. Before November 2019, he had received lecture honoraria from Roche, MagQu LLC, and Servier.

HZ has served at scientific advisory boards for Roche Diagnostics, Denali, Wave, Samumed, and CogRx; has given lectures in symposia sponsored by Alzecure and Biogen; and is a co-founder of Brain Biomarker Solutions in Gothenburg AB, a GU Ventures-based platform company at the University of Gothenburg.

KB has served as a consultant or at advisory boards for Abcam, Axon, Biogen, Lilly, MagQu, Novartis, and Roche Diagnostics and is a co-founder of Brain Biomarker Solutions in Gothenburg AB, a GU Venture-based platform company at the University of Gothenburg, all unrelated to the work presented in this paper.

ADV, FSG, AG, FC, BD, and YR report no biomedical financial interests or potential conflicts of interest.

Author details

¹Sorbonne University, GRC no 21, Alzheimer Precision Medicine (APM), AP-HP, Pitié-Salpêtrière Hospital, Paris, France. ²Brain & Spine Institute (ICM), INSERM U 1127, CNRS UMR 7225, Boulevard de l'hôpital, F-75013 Paris, France. ³Institute of Memory and Alzheimer's Disease (IM2A), Department of Neurology, Pitié-Salpêtrière Hospital, AP-HP, Boulevard de l'hôpital, F-75013 Paris, France. ⁴ADx NeuroSciences NV, Technologiepark 4, 9052 Ghent, Belgium. ⁵Department of Psychiatry and Neurochemistry, Institute of Neuroscience & Physiology, the Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden. ⁶Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden. ⁷Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square, London, UK. ⁸UK Dementia Research Institute at UCL, London, UK. ⁹Human Anatomy, Department of Translational Research and New Technologies in Medicine and Surgery, University of Pisa, Pisa, Italy. ¹⁰Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy. ¹¹Department of Drug Sciences, University of Catania, Catania, Italy. ¹²Oasi Research Institute-IRCCS, Troina, Italy. ¹³Department of Neuroscience, University of Connecticut Health, Farmington, CT, USA.

Received: 8 June 2020 Accepted: 15 September 2020

Published online: 16 October 2020

References

1. Luo Y, Bolon B, Kahn S, Bennett BD, Babu-Khan S, Denis P, et al. Mice deficient in BACE1, the Alzheimer's beta-secretase, have normal phenotype and abolished beta-amyloid generation. *Nat Neurosci*. 2001;4:231–2.
2. Vassar R, Bennett BD, Babu-Khan S, Kahn S, Mendiáz EA, Denis P, et al. Beta-secretase cleavage of Alzheimer's amyloid precursor protein by the transmembrane aspartic protease BACE. *Science*. 1999;286:735–41.
3. Hampel H, Vassar R, De Strooper B, Hardy J, Willem M, Singh N, et al. The β -Secretase BACE1 in Alzheimer's Disease. *Biol Psychiatry*. 2020;S0006-3223(20)30063–9.
4. Zolezzi JM, et al. Alzheimer's disease: relevant molecular and physiopathological events affecting amyloid- β brain balance and the putative role of PPARs. *Front Aging Neurosci*. 2014;6:176. <https://doi.org/10.3389/fnagi.2014.00176>.
5. Kandalepas PC, Sadleir KR, Eimer WA, Zhao J, Nicholson DA, Vassar R. The Alzheimer's beta-secretase BACE1 localizes to normal presynaptic terminals and to dystrophic presynaptic terminals surrounding amyloid plaques. *Acta Neuropathol*. 2013;126:329–52.
6. Sadleir KR, Kandalepas PC, Buggia-Prevot V, Nicholson DA, Thinakaran G, Vassar R. Presynaptic dystrophic neurites surrounding amyloid plaques are sites of microtubule disruption, BACE1 elevation, and increased A β generation in Alzheimer's disease. *Acta Neuropathol*. 2016;132:235–56.
7. Barão S, Moechars D, Lichtenthaler SF, De Strooper B. BACE1 physiological functions may limit its use as therapeutic target for Alzheimer's disease. *Trends Neurosci*. 2016;39:158–69.
8. Yan R, Vassar R. Targeting the beta secretase BACE1 for Alzheimer's disease therapy. *Lancet Neurol*. 2014;13:319–29.
9. Zhong Z, Ewers M, Teipel S, Burger K, Wallin A, Blennow K, et al. Levels of beta-secretase (BACE1) in cerebrospinal fluid as a predictor of risk in mild cognitive impairment. *Arch Gen Psychiatry*. 2007;64:718–26.
10. Zetterberg H, Andreasson U, Hansson O, Wu G, Sankaranarayanan S, Andersson ME, et al. Elevated cerebrospinal fluid BACE1 activity in incipient Alzheimer disease. *Arch Neurol*. 2008;65:1102–7.
11. Ewers M, Cheng X, Zhong Z, Nural HF, Walsh C, Meindl T, et al. Increased CSF-BACE1 activity associated with decreased hippocampus volume in Alzheimer's disease. *J Alzheimers Dis*. 2011;25:373–81.
12. Rosén C, Andreasson U, Mattsson N, Marcusson J, Minthon L, Andreassen N, et al. Cerebrospinal fluid profiles of amyloid β -related biomarkers in Alzheimer's disease. *Neuromolecular Med United States*. 2012;14:65–73.

13. Alexopoulos P, Thierjung N, Grimmer T, Ortner M, Economou P, Assimakopoulos K, et al. Cerebrospinal fluid BACE1 activity and sA β PP β as biomarker candidates of Alzheimer's disease. *Dement Geriatr Cogn Disord Switzerland*. 2018;45:152–61.
14. Pernecky R, Alexopoulos P. Cerebrospinal fluid BACE1 activity and markers of amyloid precursor protein metabolism and axonal degeneration in Alzheimer's disease. *Alzheimers Dement*. 2014;10:5425–9 e1.
15. Savage MJ, Holder DJ, Wu G, Kaplow J, Siuciak JA, Potter WZ. Soluble BACE-1 activity and sA β PP β concentrations in Alzheimer's disease and age-matched healthy control cerebrospinal fluid from the Alzheimer's Disease Neuroimaging Initiative-1 baseline cohort. *J Alzheimers Dis*. 2015;46:431–40.
16. Mulder SD, van der Flier WM, Verheijen JH, Mulder C, Scheltens P, Blankenstein MA, et al. BACE1 activity in cerebrospinal fluid and its relation to markers of AD pathology. *J Alzheimers Dis*. 2010;20:253–60.
17. Tsolakidou A, Alexopoulos P, Guo L-H, Grimmer T, Westerteicher C, Kratzer M, et al. β -Site amyloid precursor protein-cleaving enzyme 1 activity is related to cerebrospinal fluid concentrations of sortilin-related receptor with A-type repeats, soluble amyloid precursor protein, and tau. *Alzheimers Dement United States*. 2013;9:386–91.
18. De Vos A, Struyfs H, Jacobs D, Franssen E, Klewansky T, De Roeck E, et al. The cerebrospinal fluid neurogranin/BACE1 ratio is a potential correlate of cognitive decline in Alzheimer's disease. *J Alzheimers Dis*. 2016;53:1523–38.
19. Timmers M, Barao S, Van Broeck B, Tesseur I, Slemmon J, De Waepenaert K, et al. BACE1 dynamics upon inhibition with a BACE inhibitor and correlation to downstream Alzheimer's disease markers in elderly healthy participants. *J Alzheimers Dis*. 2017;56:1437–49.
20. Schaevebeke J, Gille B, Adamczuk K, Vanderstichele H, Chassaing E, Bruffaerts R, et al. Cerebrospinal fluid levels of synaptic and neuronal integrity correlate with gray matter volume and amyloid load in the precuneus of cognitively intact older adults. *J Neurochem*. 2019;149:139–57.
21. Grimmer T, Alexopoulos P, Tsolakidou A, Guo L-H, Henriksen G, Yousefi BH, et al. Cerebrospinal fluid BACE1 activity and brain amyloid load in Alzheimer's disease. *ScientificWorldJournal*. 2012;2012:712048.
22. Dubois B, Feldman HH, Jacova C, Hampel H, Molinuevo JL, Blennow K, et al. Advancing research diagnostic criteria for Alzheimer's disease: the IWG-2 criteria. *Lancet Neurol*. 2014;13:614–29.
23. Götz J, Chen F, van Dorpe J, Nitsch RM. Formation of neurofibrillary tangles in P301L tau transgenic mice induced by A β 42 fibrils. *Science*. 2001;293:1491–5.
24. Bloom GS. Amyloid- β and tau: the trigger and bullet in Alzheimer disease pathogenesis. *JAMA Neurol United States*. 2014;71:505–8.
25. Pooler AM, Polydoro M, Maury EA, Nicholls SB, Reddy SM, Wegmann S, et al. Amyloid accelerates tau propagation and toxicity in a model of early Alzheimer's disease. *Acta Neuropathol Commun*. 2015;3:14.
26. Barao S, Zhou L, Adamczuk K, Vanhoutvin T, van Leuven F, Demedts D, et al. BACE1 levels correlate with phospho-tau levels in human cerebrospinal fluid. *Curr Alzheimer Res*. 2013;10:671–8.
27. Hampel H, O'Bryant SE, Molinuevo JL, Zetterberg H, Masters CL, Lista S, et al. Blood-based biomarkers for Alzheimer disease: mapping the road to the clinic. *Nat Rev Neurol*. 2018;14:639–52.
28. Kirsebom B-E, Nordengen K, Selnes P, Waterloo K, Torsetnes SB, Gisladdóttir B, et al. Cerebrospinal fluid neurogranin/ β -site APP-cleaving enzyme 1 predicts cognitive decline in preclinical Alzheimer's disease. *Alzheimer's Dement (New York, N Y)*. 2018;4:617–27.
29. Schipke CG, De Vos A, Fuentes M, Jacobs D, Vanmechelen E, Peters O. Neurogranin and BACE1 in CSF as potential biomarkers differentiating depression with cognitive deficits from early Alzheimer's disease: a pilot study. *Dement Geriatr Cogn Dis Extra*. 2018;8:277–89.
30. Caraci F, Spampinato SF, Morgese MG, Tascadda F, Salluzzo MG, Giambitone MC, et al. Neurobiological links between depression and AD: the role of TGF- β 1 signaling as a new pharmacological target. *Pharmacol Res Netherlands*. 2018;130:374–84.
31. Ewers M, Zhong Z, Burger K, Wallin A, Blennow K, Teipel SJ, et al. Increased CSF-BACE 1 activity is associated with ApoE-epsilon 4 genotype in subjects with mild cognitive impairment and Alzheimer's disease. *Brain*. 2008;131:1252–8.
32. Hou X, Adeosun SO, Zhang Q, Barlow B, Brents M, Zheng B, et al. Differential contributions of ApoE4 and female sex to BACE1 activity and expression mediate A β deposition and learning and memory in mouse models of Alzheimer's disease. *Front Aging Neurosci*. 2015;7:207.
33. Decourt B, Gonzales A, Beach TG, Malek-Ahmadi M, Walker A, Sue L, et al. BACE1 levels by APOE genotype in non-demented and Alzheimer's post-mortem brains. *Curr Alzheimer Res*. 2013;10:309–15.
34. Manzine PR, Souza M da S, Cominetti MR. BACE1 levels are increased in plasma of Alzheimer's disease patients compared with matched cognitively healthy controls. *Per Med*. 2016;13:531–40.
35. Cervellati C, Trentini A, Rosta V, Passaro A, Bosi C, Sanz JM, et al. Serum beta-secretase 1 (BACE1) activity as candidate biomarker for late-onset Alzheimer's disease. *GeroScience*. 2020;42:159–67.
36. Shen Y, Wang H, Sun Q, Yao H, Keegan AP, Mullan M, et al. Increased plasma beta-secretase 1 may predict conversion to Alzheimer's disease dementia in individuals with mild cognitive impairment. *Biol Psychiatry*. 2018;83:447–55.
37. Vergallo A, Houot M, Cavedo E, Lemercier P, Vanmechelen E, De Vos A, et al. Brain A β load association and sexual dimorphism of plasma BACE1 concentrations in cognitively normal individuals at risk for AD. *Alzheimers Dement*. 2019;15:1274–85.
38. Zhao L, Morgan TE, Mao Z, Lin S, Cadenas E, Finch CE, et al. Continuous versus cyclic progesterone exposure differentially regulates hippocampal gene expression and functional profiles. *PLoS One*. 2012;7:e31267.
39. Johnston JA, Liu WW, Coulson DTR, Todd S, Murphy S, Brennan S, et al. Platelet beta-secretase activity is increased in Alzheimer's disease. *Neurobiol Aging*. 2008;29:661–8.
40. Bermejo-Bescós P, Martín-Aragón S, Jiménez-Aliaga K, Benedi J, Felici E, Gil P, et al. Processing of the platelet amyloid precursor protein in the mild cognitive impairment (MCI). *Neurochem Res*. 2013;38:1415–23.
41. Marksteiner J, Humpel C. Platelet-derived secreted amyloid-precursor protein-beta as a marker for diagnosing Alzheimer's disease. *Curr Neurovasc Res*. 2013;10:297–303.
42. Decourt B, Walker A, Gonzales A, Malek-Ahmadi M, Liesback C, Davis KJ, et al. Can platelet BACE1 levels be used as a biomarker for Alzheimer's disease? Proof-of-concept study Platelets. 2013;24:235–8.
43. Sarno TA, Talib LL, Joaquim HPG, Bram JM de F, Gattaz WF, Forlenza OV. Protein expression of BACE1 is downregulated by donepezil in Alzheimer's disease platelets. *J Alzheimers Dis*. 2017;55:1445–51.
44. Wongchitrat P, Pakpian N, Kitidee K, Chopin K, Dharmasaroja PA, Govitrapong P. Alterations in the expression of amyloid precursor protein cleaving enzymes mRNA in Alzheimer peripheral blood. *Curr Alzheimer Res United Arab Emirates*. 2019;16:29–38.
45. Vakilian A, Masoumi J, Mirzaee S, Khorramdelazad H. Expression analysis of beta-secretase 1 (BACE1) enzyme in peripheral blood of patients with Alzheimer's disease. *Casp J Intern Med*. 2019;10:276–80.
46. Faghihi MA, Modarresi F, Khalil AM, Wood DE, Sahagan BG, Morgan TE, et al. Expression of a noncoding RNA is elevated in Alzheimer's disease and drives rapid feed-forward regulation of beta-secretase. *Nat Med*. 2008;14:723–30.
47. Yang T, O'Malley TT, Kanmert D, Jerecic J, Zieske LR, Zetterberg H, et al. A highly sensitive novel immunoassay specifically detects low levels of soluble A β oligomers in human cerebrospinal fluid. *Alzheimers Res Ther*. 2015;7:14.
48. Savage MJ, Kalinina J, Wolfe A, Tugusheva K, Korn R, Cash-Mason T, et al. A sensitive A β oligomer assay discriminates Alzheimer's and aged control cerebrospinal fluid. *J Neurosci*. 2014;34:2884–97.
49. Esparza TJ, Zhao H, Cirrito JR, Cairns NJ, Bateman RJ, Holtzman DM, et al. Amyloid- β oligomerization in Alzheimer dementia versus high-pathology controls. *Ann Neurol*. 2013;73:104–19.
50. Ghafouri-Fard S, Taheri M, Arsang-Jang S, Kholghi Oskooei V, Omrani MD. Sex-based dimorphisms in expression of BDNF and BACE1 in bipolar patients. *Compr Psychiatry*. 2019;91:29–33.
51. Nafisi-Far N, Ghafouri-Fard S, Panah AST, Sayad A, Taheri M. A gender dimorphism in up-regulation of BACE1 gene expression in schizophrenia. *Metab Brain Dis*. 2018;33:933–7.
52. Nyarko JNK, Quartey MO, Pennington PR, Heistad RM, Dea D, Poirier J, et al. Profiles of beta-amyloid peptides and key secretases in brain autopsy samples differ with sex and APOE epsilon4 status: impact for risk and progression of Alzheimer disease. *Neuroscience*. 2018;373:20–36.
53. Hampel H, Vergallo A, Giorgi FS, Kim SH, Depypere H, Graziani M, et al. Precision medicine and drug development in Alzheimer's disease: the importance of sexual dimorphism and patient stratification. *Front Neuroendocrinol*. 2018;50:31–51.
54. Ferretti MT, Iulita MF, Cavedo E, Chiesa PA, Schumacher Dimech A, Santuccione Chadha A, et al. Sex differences in Alzheimer disease - the gateway to precision medicine. *Nat Rev Neurol*. 2018;14:457–69.

55. Lopez-Font I, Boix CP, Zetterberg H, Blennow K, Sáez-Valero J. Characterization of Cerebrospinal Fluid BACE1 Species. *Mol Neurobiol*. 2019; 12:8603–16.
56. Li R, Lindholm K, Yang L-B, Yue X, Citron M, Yan R, et al. Amyloid beta peptide load is correlated with increased beta-secretase activity in sporadic Alzheimer's disease patients. *Proc Natl Acad Sci U S A*. 2004;101:3632–7.
57. Yang L-B, Lindholm K, Yan R, Citron M, Xia W, Yang X-L, et al. Elevated beta-secretase expression and enzymatic activity detected in sporadic Alzheimer disease. *Nat. Med. United States*. 2003;3–4.
58. Yan R, Han P, Miao H, Greengard P, Xu H. The transmembrane domain of the Alzheimer's beta-secretase (BACE1) determines its late Golgi localization and access to beta -amyloid precursor protein (APP) substrate. *J Biol Chem United States*. 2001;276:36788–96.
59. Zhong Z, Ewers M, Teipel S, Bürger K, Wallin A, Blennow K, et al. Levels of beta-secretase (BACE1) in cerebrospinal fluid as a predictor of risk in mild cognitive impairment. *Arch Gen Psychiatry United States*. 2007;64:718–26.
60. Shi X-P, Tugusheva K, Bruce JE, Lucka A, Chen-Dodson E, Hu B, et al. Novel mutations introduced at the beta-site of amyloid beta protein precursor enhance the production of amyloid beta peptide by BACE1 in vitro and in cells. *J Alzheimers Dis Netherlands*. 2005;7:139–80.
61. Holsinger RMD, Lee JS, Boyd A, Masters CL, Collins SJ. CSF BACE1 activity is increased in CJD and Alzheimer disease versus [corrected] other dementias. *Neurology United States*. 2006;67:710–2.
62. Tsolakidou A, Alexopoulos P, Guo L-H, Grimmer T, Westerteicher C, Kratzer M, et al. β -Site amyloid precursor protein-cleaving enzyme 1 activity is related to cerebrospinal fluid concentrations of sortilin-related receptor with A-type repeats, soluble amyloid precursor protein, and tau. *Alzheimers Dement*. 2013;9:386–91.
63. Alexopoulos P, Thierjung N, Grimmer T, Ortner M, Economou P, Assimakopoulos K, et al. Cerebrospinal fluid BACE1 activity and sA β PP β as biomarker candidates of Alzheimer's disease. *Dement Geriatr Cogn Disord*. 2018;45:152–61.
64. Gonzales A, Decourt B, Walker A, Condjella R, Nural H, Sabbagh MN. Development of a specific ELISA to measure BACE1 levels in human tissues. *J Neurosci Methods*. 2011;202:70–6.
65. Verheijen JH, Huisman LGM, van Lent N, Neumann U, Paganetti P, Hack CE, et al. Detection of a soluble form of BACE-1 in human cerebrospinal fluid by a sensitive activity assay. *Clin Chem*. 2006;52:1168–74.
66. Liu L, Lauro BM, Ding L, Rovere M, Wolfe MS, Selkoe DJ. Multiple BACE1 inhibitors abnormally increase the BACE1 protein level in neurons by prolonging its half-life. *Alzheimers Dement*. 2019;15:1183–94.
67. Sinha S, Anderson JP, Barbour R, Basi GS, Caccavello R, Davis D, et al. Purification and cloning of amyloid precursor protein beta-secretase from human brain. *Nature*. 1999;402:537–40.
68. Decourt B, Sabbagh MN. BACE1 as a potential biomarker for Alzheimer's disease. *J Alzheimers Dis*. 2011;24(Suppl 2):53–9.
69. Willems EAJ, Vermeiren Y, Garcia-Ayllon M-S, Bridel C, De Deyn PP, Engelborghs S, et al. Pre-analytical stability of novel cerebrospinal fluid biomarkers. *Clin Chim Acta*. 2019;497:204–11.
70. Zhang L, Chen L, Dutra JK, Beck EM, Nag S, Takano A, et al. Identification of a novel positron emission tomography (PET) ligand for imaging β -site amyloid precursor protein cleaving enzyme 1 (BACE-1) in brain. *J Med Chem*. 2018;61:3296–308.
71. Andreasson U, Perret-Liaudet A, van Waalwijk van Doorn LJ, et al. A practical guide to immunoassay method validation. *Front Neurol*. 2015;6:179. Published 2015 Aug 19. doi:<https://doi.org/10.3389/fneur.2015.00179>.
72. Moreno-Jiménez EP, Flor-García M, Terreros-Roncal J, Rábano A, Cafini F, Pallas-Bazarra N, et al. Adult hippocampal neurogenesis is abundant in neurologically healthy subjects and drops sharply in patients with Alzheimer's disease. *Nat Med*. 2019;25:554–60.
73. Lazarov O, Hollands C. Hippocampal neurogenesis: learning to remember. *Prog Neurobiol*. 2016;138–140:1–18.
74. Cavado E, Lista S, Houot M, Vergallo A, Grothe MJ, Teipel S, et al. Plasma tau correlates with basal forebrain atrophy rates in people at risk for Alzheimer disease. *Neurology*. 2020;94:e30–41.
75. Jiang H, Hampel H, Prvulovic D, Wallin A, Blennow K, Li R, et al. Elevated CSF levels of TACE activity and soluble TNF receptors in subjects with mild cognitive impairment and patients with Alzheimer's disease. *Mol Neurodegener*. 2011;6:69.
76. He P, Zhong Z, Lindholm K, Berning L, Lee W, Lemere C, et al. Deletion of tumor necrosis factor death receptor inhibits amyloid beta generation and prevents learning and memory deficits in Alzheimer's mice. *J Cell Biol*. 2007; 178:829–41.
77. Portelius E, Dean RA, Andreasson U, Mattsson N, Westerlund A, Olsson M, et al. Beta-site amyloid precursor protein-cleaving enzyme 1 (BACE1) inhibitor treatment induces Abeta5-X peptides through alternative amyloid precursor protein cleavage. *Alzheimers Res Ther*. 2014;6:75.
78. Giuffrida ML, Tomasello MF, Pandini G, Caraci F, Battaglia G, Busceti C, et al. Monomeric β -amyloid interacts with type-1 insulin-like growth factor receptors to provide energy supply to neurons. *Front Cell Neurosci*. 2015;9: 297.
79. Hampel H, Goetzl EJ, Kapogiannis D, Lista S, Vergallo A. Biomarker-Drug and Liquid Biopsy Co-development for Disease Staging and Targeted Therapy: Cornerstones for Alzheimer's Precision Medicine and Pharmacology. *Front Pharmacol*. 2019;10:310.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.