

Symmetries and special solutions of a parabolic chemotaxis system

M. S. Bruzón¹ | M. L. Gandarias¹  | M. Torrisi²  | R. Tracinà²

¹Departamento de Matemáticas, Universidad de Cádiz, Puerto Real, Cádiz, 11510, Spain

²Dipartimento di Matematica e Informatica, University of Catania, Viale A. Doria, 6, Catania, 95125, Italy

Correspondence

M. Torrisi, Dipartimento di Matematica e Informatica, University of Catania, Viale A. Doria, 6, 95125 Catania, Italy.
Email: torrisi@dmi.unict.it

Communicated by: J. Muñoz Rivera

Funding information

Istituto Nazionale di Alta Matematica "Francesco Severi", Grant/Award Number: G.N.F.M.; Junta de Andalucía FQM-201 group and University of Cádiz; Università di Catania, Grant/Award Number: Piano triennale della Ricerca 2016-2018; Junta de Andalucía FQM-201 group; University of Cádiz; Università di Catania

In this paper, we consider a class of chemotaxis models with two arbitrary constitutive functions $g(u)$ and $f(v)$. After having performed a complete symmetry group classification with respect to them the reduced systems are derived. By considering $g(u)$ of the logistic form, wide classes of exact solutions are found.

KEYWORDS

chemotaxis, nonlinear parabolic equations, symmetries

MSC CLASSIFICATION

92C17; 58J70; 35K55

1 | INTRODUCTION

The phenomena concerned with the motion of bacteria and other unicellular organisms due to chemical stimuli (in a *chemical substrate*) are called *chemotaxis*. These phenomena are quite common in biology and depend on the essential characteristic of living organisms to sense signals in the environment and to adapt their motion consequently. These signals allow to approach the chemically favorable environments and to avoid the unfavorable ones.

In chemotaxis, the chemical signals can come from external sources, or they can be secreted by the organisms themselves. Chemotaxis can be either positive (chemoattraction) when cells move to the chemical or negative (chemorepulsion) when cells move away from the chemical. Keller and Segel in some remarkable papers,¹⁻³ concerned with studies about chemotactical features of the slime mold amoebae *Dictyostelium discoideum* and *Escherichia coli*, considered a phenomenological model from which the existence and properties of migrating bacterial bands can be deduced. Following Keller and Segel,³ the equations of a reduced model for a population of density u , with a chemoattractant of concentration v , could be written as

$$\begin{aligned} u_t &= \nabla \cdot (\mu(v)\nabla u - u\chi(v)\nabla v), \\ v_t &= D_v \nabla^2 v - k(v)u, \end{aligned} \quad (1)$$

where $k(v)$ is the rate of consumption of the substrate per cell, D_v is the diffusion constant of the substrate while χ is called the chemotactic sensitivity; it shows how strong the cells (species) react to the chemical. In general, the sign of χ

depends on the type of chemotaxis: in case of chemoattraction, its sign is positive, whereas in case of chemorepulsion, its sign is negative. The motility parameter μ takes the place of a diffusion coefficient, and in general, it may be considered as a function of v .

Hillen and Painter on 2001⁴ modified system (1) as follows:

$$\begin{aligned} u_t &= \nabla \cdot (\nabla u - V(u, v)\nabla v), \\ v_t &= D_v \nabla^2 v + h(u, v). \end{aligned} \quad (2)$$

In the first equation, they put $\mu = 1$ and rewritten $u\chi$ as a more general function V depending on u and v . While, in the second equation, the function describing production and degradation of the external stimulus has been generalized. In this way, the population density directly modulates its own sensitivity response. In particular in Hillen and Painter,⁴ they substituted the *chemotactic cross-diffusion* $V(u, v)\nabla v$ with the following $u\beta(u)\chi(v)\nabla v$ to show global existence of classical solutions.

The same authors⁵ used a probabilistic approach to arrive at the following equations:

$$\begin{aligned} u_t &= D_u \nabla^2 u - \nabla \cdot (u\chi(u, v)\nabla v) + g(u, v) \\ v_t &= D_v \nabla^2 v + h(u, v) \end{aligned} \quad (3)$$

that are of the same form originally proposed by Patlak⁶ and by Keller and Segel,² to whom have been added cell and chemical kinetics of the form $g(u, v)$ and $h(u, v)$, respectively.

In several biologically relevant processes, when the chemical substance is produced by the individuals of the population, the function h of (3) satisfies the inequality $h_u > 0$. So, as in the classical Keller–Segel system, it is possible to consider a degradation of v and simplify the term $h(u, v)$ by the linear expression $h(u, v) = h_0 u - h_1 v$ with $h_0 > 0$, $h_1 > 0$ that without loss of generality, we could normalize putting both equal to 1.

In their study,⁷ Negreanu and Tello considered a volume filling model with fast diffusion process for the chemical substance with logistic growth term

$$\begin{aligned} u_t &= \nabla^2 u - c\nabla \cdot (u(N - u)\nabla v) + \lambda u(1 - u), \\ \nabla^2 v &= v - u, \end{aligned} \quad (4)$$

where $c > 0$ is a constant referred to the chemotactic sensitivity, while as in Painter and Hillen⁵ a threshold value $N > 0$ is considered. The sign change of the chemotactic cross-diffusion term characterizes the possibility of the system to evolve from positive to negative taxis or vice versa.

Negreanu and Tello in that paper proved the existence of global bounded classical solutions and analyzed the stability of the constant steady state $u = 1$, $v = 1$.

By assuming

$$V(u, v)\nabla v := cu\nabla v,$$

we can get the following simplified Keller–Segel system:

$$\begin{aligned} u_t &= u_{xx} + cuv_{xx} - cu_x v_x, \\ 0 &= v_{xx} - \beta v + u, \end{aligned} \quad (5)$$

whose properties have been studied in several papers.^{8–12}

In particular, in Cherniha and Didovych⁹ and Didovych,¹⁰ Lie symmetries have been found and applied for construction of exact solutions.

Here, taking into account other studies,^{5,9,10} we restrict ourselves to the following set of the constitutive relations:

$$D_u = D_v = 1, \quad V(u, v) = c(N - u), \quad g(u, v) = g(u), \quad h(u, v) = -f(v) + u,$$

where $f = f(v)$ and $g = g(u)$ are non-negative smooth functions. So we are able to rewrite (3) as

$$\begin{aligned}u_t &= \nabla^2 u - c(\nabla \cdot ((N - u)\nabla v)) + g(u) \\v_t &= \nabla^2 v - f(v) + u,\end{aligned}\tag{6}$$

that in one space dimension reads:

$$\begin{aligned}u_t &= u_{xx} - c(N - u)v_{xx} + cu_x v_x + g(u) \\v_t &= v_{xx} - f(v) + u.\end{aligned}\tag{7}$$

When growth effects in chemotaxis systems are considered to study the large time behavior, the growth term g in the first equation is defined by a logistic function and, after normalization, g has the following expression:

$$g(u) = u(1 - u).\tag{8}$$

In this paper, we look for symmetries of system (7) whose knowledge allows to reduce the system of partial differential equations (PDEs) into a system of ordinary differential equations (ODEs) and to find exact solutions.

Lie group methods offer good tools as conservation laws, reductions, exact solutions, and give a systematic way to analyze differential equations.

The structure of the work is as follows. In Section 2, we look for the Lie symmetry classification with respect to functions $f(u)$ and $g(u)$ of system (7). We also derive in Section 3 the reductions obtained from some Lie subalgebras admitted. In Section 4, by assuming the function $g(u)$ of the logistic type¹³⁻¹⁵ together with some suitable specializations of $f(v)$, we find some exact solutions of system (7). For more general polynomial forms of $f(u)$ and $g(u)$, in Section 5, solutions of the kink type and soliton type are carried out. Conclusions are given in Section 6.

2 | LIE SYMMETRIES

The search for solutions via classical Lie symmetries leads to the search for those that are invariant with respect to the same one parameter Lie groups of invariant transformations admitted by system (7). By following a well-established procedure (for details, the interested reader can see well-known monographs¹⁶⁻¹⁸) the first step consists in determining the infinitesimal components $\xi(t, x, u, v)$, $\tau(t, x, u, v)$, and $\eta(t, x, u, v)$ and $\phi(t, x, u, v)$ of transformation generator

$$\mathbf{v} = \xi \partial_x + \tau \partial_t + \eta \partial_u + \phi \partial_v.\tag{9}$$

Following the usual techniques, from the request invariance of system (7), by applying the Lie infinitesimal criterion, we get a system of equations (*determining system*) to determine the infinitesimal components of \mathbf{v} .

After a first simplification of this one, we get that

$$\begin{aligned}\xi &= \frac{1}{2}\tau_t x + \omega(t), \quad \tau = \tau(t), \\ \eta &= \alpha(x, t)u + \beta(x, t, v), \quad \phi = \delta(x, t)\end{aligned}$$

where $\tau(t)$, $\omega(t)$, $\alpha(t, x)$, $\beta(x, t, v)$ and $\delta(x, t)$ must satisfy the following equations:

$$\begin{aligned}\beta_v c + \beta_{vv} &= 0, \\ \alpha_x c u + \beta_x c + 2\beta_{vx} &= 0, \\ \tau_t u + \alpha u - f \tau_t - \delta f_v - \delta_t + \delta_{xx} + \beta &= 0, \\ \frac{\tau_{tt} x}{2} + \omega_t + c \delta_x + 2\alpha_x &= 0, \\ \frac{\tau_{tt} x}{2} + \omega_t &= 0, \\ c(\alpha N + \beta) &= 0, \\ \alpha c u N - \alpha c f N + c \delta_{xx} N - \alpha g_u u - c \delta_{xx} u + \beta c u + \beta_v u - \alpha_{xx} u + \\ \alpha_t u - g \tau_t - \beta g_u + \alpha g - \beta c f - \beta_v f - \beta_{xx} + \beta_t &= 0.\end{aligned}\tag{10}$$

TABLE 1 Symmetry classification for system (7) with $c \neq 0$. Here k, α, β, f_0 are constitutive constants. Of course, in this table appear only the extensions of \mathcal{L}_p

i	f	g	\mathbf{v}_3^i	\mathbf{v}_4^i
1	Arbitrary	Arbitrary		
2	α	Arbitrary	∂_v	
3	$\alpha v + f_0, \alpha \neq 0$	Arbitrary	$\exp(-\alpha t) \partial_v$	
4	α	$k(N - u)^2$	$x\partial_x + 2t\partial_t + 2(N - u)\partial_u + 2(N - \alpha)t\partial_v$	∂_v
5	$N + \alpha \exp(\beta v),$	$k(N - u)^2$	$x\partial_x + 2t\partial_t + 2(N - u)\partial_u - \frac{2}{\beta}\partial_v$	

TABLE 2 Similarity variables and similarity solutions

i	\mathbf{v}^i	z	u	v
1	$\lambda \mathbf{v}_1 + \mu \mathbf{v}_2$	$\mu x - \lambda t$	$H(z)$	$F(z)$
2	$\lambda \mathbf{v}_1 + \mu \mathbf{v}_2 + \mathbf{v}_3^2$	$\mu x - \lambda t$	$H(z)$	$\frac{x}{\lambda} + F(z)$
3	$\lambda \mathbf{v}_1 + \mu \mathbf{v}_2 + \mathbf{v}_3^3$	$\mu x - \lambda t$	$H(z)$	$F(z) - \frac{e^{-\alpha t}}{\alpha \mu}$
4	\mathbf{v}_3^4	$\frac{x}{\sqrt{t}}$	$\frac{H(z)}{t} + N$	$F(z) + (N - \alpha)t$
5	\mathbf{v}_3^5	$\frac{x}{\sqrt{t}}$	$\frac{H(z)}{t} + N$	$F(z) - \frac{\ln t}{\beta}$

A further analysis allows to ascertain that

$$\xi = \frac{k_1 x}{2} + k_3, \quad \tau = k_1 t + k_2, \quad \eta = k_1(N - u), \quad \phi = \gamma(t),$$

with k_2, k_3 arbitrary constants, while the function $\gamma = \gamma(t)$ and the constant k_1 are related to the constitutive functions $f = f(v)$ and $g = g(u)$ by the following conditions:

$$k_1 N - f k_1 - \gamma_t - f_v \gamma = 0, \quad k_1 (g_u N - g_u u + 2g) = 0. \quad (11)$$

If f and g are arbitrary functions, we obtain

$$\xi = k_3, \quad \tau = k_2, \quad \eta = 0, \quad \phi = 0, \quad (12)$$

and the only symmetries admitted by (7) are defined by the group of space and time translations,

$$\mathbf{v}_1 = \partial_x, \quad \mathbf{v}_2 = \partial_t,$$

and they constitute the principal Lie algebra \mathcal{L}_p of system (7). Moreover, by discussing the classifying system (11), we get additional extensions of the principal Lie algebra and a classification with respect to the functions f and g .

The picture of the results is summarized in Table 1.

By considering the invariance generators listed in Table 2 and following the well-known procedures, we find the corresponding similarity variables and the structure of the invariant solutions so that we write the corresponding ODE reduced systems.

3 | REDUCED SYSTEMS

The form of the invariant solutions admitted by system (7) is derived from their invariance conditions

$$\mathbf{v}[u, v]^T = 0, \quad (13)$$

where $[u, v]^T$ denotes the transposed of the vector row $[u, v]$. By specializing (13) to the case of the principal Lie algebra that is for $\mathbf{v} = \lambda \partial_x + \mu \partial_t$, we get the following PDE system:

$$\lambda u_x + \mu u_t = 0 \quad (14)$$

$$\lambda v_x + \mu v_t = 0 \quad (15)$$

TABLE 3 System of ODEs

<i>i</i>	<i>ODE_i</i>
1	$H' \lambda - c \mu^2 F'' N + \mu^2 H'' + c \mu^2 F' H' + c \mu^2 F'' H + g = 0$ $f - F' \lambda - H - \mu^2 F'' = 0.$
2	$H' \lambda + \frac{c \mu H'}{\lambda} - c \mu^2 F'' N + \mu^2 H'' + c \mu^2 F' H' + c \mu^2 F'' H + g = 0,$ $-F' \lambda - H - \mu^2 F'' + \alpha = 0.$
3	$H' \lambda - c \mu^2 F'' N + \mu^2 H'' + c \mu^2 F' H' + c \mu^2 F'' H + g = 0,$ $-F' \lambda - H - \mu^2 F'' + \alpha F + f_0 = 0.$
4	$2H'' + 2cF' H' + zH' + 2kH^2 + 2cF'' H + 2H = 0,$ $2H + 2F'' + zF' = 0.$
5	$2H'' + 2cF' H' + zH' + 2kH^2 + 2cF'' H + 2H = 0,$ $2\beta H + 2\beta F'' + \beta zF' - 2\alpha \beta e^{\beta F} + 2 = 0.$

from where we derive the invariant solutions

$$\begin{aligned} u(t, x) &= H(z), \\ v(t, x) &= F(z), \end{aligned} \tag{16}$$

with the similarity variable $z = \mu x - \lambda t$. Substituting (16) into (7) we obtain the system ODE'_1 of the Table 3 for the search of solution of the traveling wave type

$$\begin{aligned} H' \lambda - c \mu^2 F'' N + \mu^2 H'' + c \mu^2 F' H' + c \mu^2 F'' H + g &= 0, \\ f - F' \lambda - H - \mu^2 F'' &= 0, \end{aligned} \tag{17}$$

where, of course, f and g are arbitrary functions.

In the following sections, we focus our attention only on the search for some special cases.

Solutions and analyses of additional cases will be considered in further researches.

4 | EXACT SOLUTIONS WITH $g(u)$ OF THE LOGISTIC FORM.

4.1 | Case 1

Here we consider the ODE system (17) derived by reducing system (7) and specializing the constitutive functions $g(u)$ and $f(v)$ as

$$g(u) = u(1 - u), \quad f(v) = v, \tag{18}$$

that is by assuming the growth term of the first equation to be of logistic type. On the basis of these specializations, the system (17) reads

$$\begin{aligned} H' \lambda - c \mu^2 F'' N + \mu^2 H'' + c \mu^2 F' H' + c \mu^2 F'' H + H(1 - H) &= 0, \\ F - F' \lambda - H - \mu^2 F'' &= 0. \end{aligned} \tag{19}$$

It is a simple matter to verify that in the case $N < 1$ an exact solution of the system (19) has the form

$$H = 1 - b_0 \frac{(N - 1)}{2} e^{b_1 z}, \tag{20}$$

$$F = 1 + b_0 e^{b_1 z} \tag{21}$$

with $b_0, b_1 \neq 0$ arbitrary constants and


$$z = \pm x \sqrt{\frac{1 - N}{4b_1^2 c}} + \frac{1 - N - 2c(N + 1)}{4b_1 c} t. \tag{22}$$

Then for the system (7), we get

$$u(t, x) = 1 - b_0 \frac{N - 1}{2} e^{b_1 \left(\pm x \sqrt{\frac{1 - N}{4b_1^2 c}} + \frac{1 - N - 2c(N + 1)}{4b_1 c} t \right)}, \tag{23}$$

$$v(t, x) = 1 + b_0 e^{b_1 \left(\pm x \sqrt{\frac{1 - N}{4b_1^2 c}} + \frac{1 - N - 2c(N + 1)}{4b_1 c} t \right)}. \tag{24}$$

FIGURE 1 The solution $u(t, 1)$ given by (23) with $b_0 = 1$, $b_1 = 1$, $c = 1$, $\lambda = 1$, $\mu = 1$ [Colour figure can be viewed at wileyonlinelibrary.com]



In Figure 1, we show the density $u(t, 1)$ given by (23) for different values of N .

Taking into account that with these forms of f and g , we fall in the Case 3 of Table 1, we can use the infinitesimal generator $\mathbf{v} = \lambda \mathbf{v}_1 + \mu \mathbf{v}_2 + \mathbf{v}_3^3$. Moreover, we observe that the corresponding reduced system is the same of system (19); then, by using the same solution (20), (21), going back to the original variables for system (7), we get

$$u(t, x) = 1 - b_0 \frac{N-1}{2} e^{b_1 \left(\pm x \sqrt{\frac{1-N}{4b_1^2 c}} + \frac{1-N-2c(N+1)}{4b_1 c} t \right)}, \quad (25)$$

$$v(t, x) = 1 + b_0 e^{b_1 \left(\pm x \sqrt{\frac{1-N}{4b_1^2 c}} + \frac{1-N-2c(N+1)}{4b_1 c} t \right)} - \frac{e^{-t}}{\sqrt{\frac{1-N}{4b_1^2 c}}}. \quad (26)$$

4.2 | Case 2

Here, taking into account Painter and Hillen⁵ and Galakhova et al,¹⁹ we consider $g(u)$ of the following generalized logistic form:

$$g(u) = g_0 u(1 - u^2), \quad (27)$$

and

$$f(v) = \alpha = \text{const}. \quad (28)$$

We fall in the Case 2 (Table 1), and we use the infinitesimal generator $\mathbf{v} = \lambda \mathbf{v}_1 + \mu \mathbf{v}_2 + \mathbf{v}_3^3$. From where we get the invariant solutions,

$$\begin{aligned} u(x, t) &= H(z), \\ v(x, t) &= F(z) + \frac{x}{\lambda}, \end{aligned} \quad (29)$$

with the similarity variable $z = \mu x - \lambda t$.

The reduced system is written in Table 3 (Case 2) and, by setting $g_0 = \frac{3c}{N}$, it is possible to verify that a solution is

$$H(z) = h_1 e^{\frac{-\lambda z}{2\mu^2}}, \quad (30)$$

$$F(z) = \frac{2h_1^2 \mu^2}{\lambda^2 N} e^{\frac{-\lambda z}{\mu^2}} + \frac{4\mu^2 h_1}{\lambda^2} e^{-\frac{\lambda z}{2\mu^2}} + \frac{(12c\mu^2 - 4cN^2\mu^2 - 2cN\mu - \lambda^2 N)z + 2f_0 \lambda c N \mu^2}{2cN \lambda \mu^2}. \quad (31)$$

that for the system (7) becomes

$$u(t, x) = h_1 e^{\frac{-\lambda(\mu x - \lambda t)}{2\mu^2}} \quad (32)$$

$$\begin{aligned} v(t, x) &= \frac{2h_1^2 \mu^2}{\lambda^2 N} e^{\frac{-\lambda(\mu x - \lambda t)}{\mu^2}} + \frac{4\mu^2 h_1}{\lambda^2} e^{-\frac{\lambda(\mu x - \lambda t)}{2\mu^2}} \\ &+ \frac{(12c\mu^2 - 4cN^2\mu^2 - 2cN\mu - \lambda^2 N)(\mu x - \lambda t) + 2f_0 \lambda c N \mu^2}{2cN \lambda \mu^2} + \frac{x}{\lambda}. \end{aligned} \quad (33)$$

5 | SPECIAL TRAVELING WAVE SOLUTIONS

We consider solutions for system (17) of the following form:

$$\begin{aligned} H(z) &= b_1 \tanh^s z + b_2, \\ F(z) &= b_3 \tanh^r z + b_4, \end{aligned} \quad (34)$$

where $b_i, (i = 1, \dots, 4)$ are constants with $b_1 \neq 0, b_3 \neq 0$, that for system (7) read as

$$u(t, x) = b_1 \tanh^s(\mu x - \lambda t) + b_2, \quad (35)$$

$$v(t, x) = b_3 \tanh^r(\mu x - \lambda t) + b_4. \quad (36)$$

By choosing f and g in some more general polynomial forms, it is a simple matter to verify that the system (17) admits solutions of the form (34).

In the following, we consider special solutions of the kink type and soliton type by assuming, respectively, $r = s = 1$ and $r = s = 2$.

5.1 | Solutions of the kink type

For $r = s = 1$, specializing f and g in the form

$$g(u) = j_0 + j_1 u + j_2 u^2 + j_3 u^3 + j_4 u^4, \quad (37)$$

$$f(v) = k_3 v^3 + k_2 v^2 + k_1 v + k_0, \quad (38)$$

with the following constraints between the constitutive constants

$$\begin{aligned} j_0 &= \frac{(b_2 - b_1)(b_1 + b_2)(b_1^2 \lambda + b_1^2 b_3 \mu^2 c + 2 b_2 b_1 \mu^2 - b_2^2 b_3 \mu^2 c - 2 b_2 c \mu^2 N b_3)}{b_1^3}, \\ j_1 &= 2 \frac{3 b_2^3 b_3 \mu^2 c - 3 b_1^2 c \mu^2 b_3 b_2 - b_1^2 c \mu^2 N b_3 + 3 b_2^2 c \mu^2 N b_3 + b_1^3 \mu^2 - 3 b_2^2 b_1 \mu^2 - b_2 b_1^2 \lambda}{b_1^3}, \\ j_2 &= \frac{6 b_2 b_1 \mu^2 + 4 b_1^2 b_3 \mu^2 c + b_1^2 \lambda - 12 b_2^2 b_3 \mu^2 c - 6 b_2 c \mu^2 N b_3}{b_1^3}, \\ j_3 &= -2 \frac{\mu^2 (b_1 - 5 c b_3 b_2 - c N b_3)}{b_1^3}, \\ j_4 &= -3 \frac{b_3 \mu^2 c}{b_1^3}, \\ k_0 &= -\frac{-b_3^3 \lambda - 2 b_4 \mu^2 b_3^2 - b_3^2 b_2 + 2 b_4^3 \mu^2 + b_4^2 b_3 \lambda + b_4 b_3 b_1}{b_3^2}, \\ k_1 &= \frac{-2 \mu^2 b_3^2 + b_3 b_1 + 6 b_4^2 \mu^2 + 2 b_4 b_3 \lambda}{b_3^2}, \\ k_2 &= -\frac{b_3 \lambda + 6 b_4 \mu^2}{b_3^2}, \\ k_3 &= 2 \frac{\mu^2}{b_3^2}, \end{aligned} \quad (39)$$

we obtain the solution

$$H(z) = b_1 \tanh z + b_2, \quad (40)$$

$$F(z) = b_3 \tanh z + b_4, \quad (41)$$

and for system (7), we have the kink solution

$$u(t, x) = b_1 \tanh(\mu x - \lambda t) + b_2, \quad (42)$$

$$v(t, x) = b_3 \tanh(\mu x - \lambda t) + b_4. \quad (43)$$

In Figure 2, we show the density u given by (42) for different values of x .

FIGURE 2 The kink solution u given by (42) with $b_1 = 1$, $b_2 = 2$, $\lambda = 1$, $\mu = 1$ [Colour figure can be viewed at wileyonlinelibrary.com]





FIGURE 3 The solution $u(t, x)$ given by (49) with $b_1 = -1$, $b_2 = 0$, $\lambda = 1$, $\mu = 1$ [Colour figure can be viewed at wileyonlinelibrary.com]



5.2 | Solutions of the soliton type

For $r = s = 2$, specializing f and g in the form

$$g(u) = g_0 + g_1 u + g_2 u^2 + g_3 u^3 + g_4 \left(\frac{u}{b_1} - \frac{b_2}{b_1} \right)^{\frac{3}{2}} + g_5 \left(\frac{u b_1}{-b_1} - \frac{b_2}{-b_1} \right)^{\frac{1}{2}}, \quad (44)$$

$$f(v) = c_0 + c_1 v + c_2 v^2 + c_3 \left(\frac{v}{b_3} - \frac{b_4}{b_3} \right)^{\frac{3}{2}} + c_4 \left(\frac{v}{b_3} - \frac{b_4}{b_3} \right)^{\frac{1}{2}}, \quad (45)$$

with

$$\begin{aligned} g_0 &= \frac{2(b_2 + b_1)\mu^2(3b_2 b_3 c N + b_1 b_3 c N + 2b_2^2 b_3 c + 2b_1 b_2 b_3 c - 3b_1 b_2 - b_1^2)}{b_1^2}, \\ g_1 &= -\frac{2\mu^2(6b_2 b_3 c N + 4b_1 b_3 c N + 9b_2^2 b_3 c + 12b_1 b_2 b_3 c + 3b_1^2 b_3 c - 6b_1 b_2 - 4b_1^2)}{b_1^2}, \\ g_2 &= \frac{2\mu^2(3b_3 c N + 12b_2 b_3 c + 8b_1 b_3 c - 3b_1)}{b_1^2}, \\ g_3 &= -\frac{10b_3 c \mu^2}{b_1^2}, \\ g_4 &= 2\lambda b_1, \\ g_5 &= -2\lambda b_1, \\ c_0 &= \frac{((6b_3 + 12)b_4^2 + (8b_3^2 + 8b_3)b_4 + 2b_3^3)\mu^2 - b_1 b_4 + b_2 b_3^2}{b_3^2}, \\ c_1 &= -\frac{12b_4 \mu^2}{b_3} - 8\mu^2 + \frac{b_1}{b_3}, \\ c_2 &= \frac{6\mu^2}{b_3}, \\ c_3 &= -2\lambda b_3, \end{aligned} \quad (46)$$

we obtain a solution

$$H(z) = b_1 \tanh^2 z + b_2, \quad (47)$$

$$F(z) = b_3 \tanh^2 z + b_4 \quad (48)$$

and for the system (7), we have the soliton solution

$$u(t, x) = b_1 \tanh^2(\mu x - \lambda t) + b_2, \quad (49)$$

$$v(t, x) = b_3 \tanh^2(\mu x - \lambda t) + b_4. \quad (50)$$

In Figure 3, we show the density u given by (49) for different values of x .

6 | CONCLUSIONS

In this short paper, a wide class of generalized chemotaxis systems, characterized by suitable constitutive relations, has been considered. In the growth term of the first equation, an arbitrary function $g(u)$ appears while in the second equation, assuming function $h(u, v) = -f(v) + u$, we introduce an arbitrary positive function $f(v)$. The complete symmetry classification has been performed with respect to the arbitrary functions $g(u)$ and $f(v)$. Once reduced systems were obtained, we focused our attention in forms of $g(u)$ of the logistic type. Wide classes of solutions have been derived. Finally, by choosing $g(u)$ and $f(v)$ of a more general polynomial form, solutions of the kink type and of the soliton type are shown.

ACKNOWLEDGEMENTS

The support of Junta de Andalucía FQM-201 group and University of Cádiz is gratefully acknowledged by M.L. Gandarias and M.S. Bruzón. M. Torrisi acknowledges support from G.N.F.M. of INdAM. R. Tracina acknowledges Università di Catania for financial support: Ricerca di Ateneo PTR *Metodi gruppali e umbrali per modelli di diffusione e trasporto*.

CONFLICT OF INTEREST

This work does not have any conflicts of interest.

ORCID

M. L. Gandarias  <https://orcid.org/0000-0001-8604-8272>

M. Torrisi  <https://orcid.org/0000-0003-0386-6216>

REFERENCES

1. Keller EF, Segel LA. Initiation of slime mold aggregation viewed as an instability. *J theor Biol.* 1970;399-4(15):26.
2. Keller EF, Segel LA. A model for chemotaxis. *J Theoret Biol.* 1971;30:225–34.
3. Keller EF, Segel LA. Traveling bands of chemotactic bacteria: a theoretical analysis. *J theor Biol.* 1971;30:235–248.
4. Hillen T, Painter K. Global existence for a parabolic chemotaxis model with prevention of overcrowding. *Adv Appl Math.* 2001;26(4):280–301.
5. Painter K, Hillen T. Volume-filling and quorum-sensing in models for chemosensitive movement. *Can Appl Math Q.* 2002;10(4):501–543.
6. Patlak CS. Random walk with persistence and external bias. *Bull Math Biophys.* 1953;15:311–338.
7. Negreanu M, Tello JI. On a parabolic-elliptic chemotactic system with non-constant chemotactic sensitivity. *Nonlin Anal.* 2013;80:1–13.
8. Calvez V, Yasmin DS. Asymptotic behavior of a two-dimensional Keller–Segel model with and without density control. *Math Model Biol Syst.* 2007;2:323–329.
9. Cherniha R, Didovych M. Exact solutions of the simplified Keller–Segel model. *Commun Nonlinear Sci Numer Simulat.* 2013;18:2960–2971.
10. Didovych M. A (1 + 2)-dimensional simplified Keller–Segel model: Lie symmetry and exact solutions. *Symmetry.* 2015;7:1463–1474.
11. Horstmann D. From 1970 until present: the Keller–Segel model in chemotaxis and its consequences, Leipzig: Max-Planck-institute for Mathematics in the Sciences, Jahresber. *Dtsch Math-Ver.* 2003;105:103–165.
12. Nagai T. Blow-up of radially symmetric solutions to a chemotaxis system. *Adv Math Sci Appl.* 1995;5:581–601.
13. Galakov E, Salieva O, Tello JI. On a parabolic-elliptic system with chemotaxis and logistic type growth. *J Differential Equations.* 2016;261:4631–4647.
14. Marras M, Vernier-Piro S, Viglialoro G. Decay in chemotaxis systems with a logistic term. *Discr Contin Dyn Syst Ser S.* 2020;13(2):257–268.
15. Salako RB, Shen W. Existence of traveling wave solutions to parabolic-elliptic-elliptic chemotaxis systems with logistic source. *Discr Contin Dyn Syst - S.* 2020;13(2):293–319.
16. Olver PJ. *Applications of Lie Groups to Differential Equations.* New York: Springer; 1986.
17. Bluman GW, Kumei S. *Symmetries of Differential Equations.* New York: Springer; 1989.
18. Ibragimov NH. *CRC Handbook of Lie Group Analysis of Differential Equations.* Boca Raton, FL, USA: CRC Press; 1996.
19. Galakhova E, Salievab O, Tello JI. On a parabolic–elliptic system with chemotaxis and logistic type growth. *J Differential Equations.* 2016;261:4631–4647.

How to cite this article: Bruzón MS, Gandarias ML, Torrisi M, Tracina R. Symmetries and special solutions of a parabolic chemotaxis system. *Math Meth Appl Sci.* 2021;44:2050–2058. <https://doi.org/10.1002/mma.6914>