

A class of functionals possessing multiple global minima

Biagio Ricceri

To Professor Gheorghe Moroşanu, with friendship, on his 70th birthday.

Abstract. We get a new multiplicity result for gradient systems. Here is a very particular corollary: Let $\Omega \subset \mathbf{R}^n$ ($n \geq 2$) be a smooth bounded domain and let $\Phi : \mathbf{R}^2 \rightarrow \mathbf{R}$ be a C^1 function, with $\Phi(0, 0) = 0$, such that

$$\sup_{(u,v) \in \mathbf{R}^2} \frac{|\Phi_u(u, v)| + |\Phi_v(u, v)|}{1 + |u|^p + |v|^p} < +\infty$$

where $p > 0$, with $p = \frac{2}{n-2}$ when $n > 2$.

Then, for every convex set $S \subseteq L^\infty(\Omega) \times L^\infty(\Omega)$ dense in $L^2(\Omega) \times L^2(\Omega)$, there exists $(\alpha, \beta) \in S$ such that the problem

$$-\Delta u = (\alpha(x) \cos(\Phi(u, v)) - \beta(x) \sin(\Phi(u, v)))\Phi_u(u, v) \quad \text{in } \Omega$$

$$-\Delta v = (\alpha(x) \cos(\Phi(u, v)) - \beta(x) \sin(\Phi(u, v)))\Phi_v(u, v) \quad \text{in } \Omega$$

$$u = v = 0 \quad \text{on } \partial\Omega$$

has at least three weak solutions, two of which are global minima in $H_0^1(\Omega) \times H_0^1(\Omega)$ of the functional

$$(u, v) \rightarrow \frac{1}{2} \left(\int_{\Omega} |\nabla u(x)|^2 dx + \int_{\Omega} |\nabla v(x)|^2 dx \right) - \int_{\Omega} (\alpha(x) \sin(\Phi(u(x), v(x))) + \beta(x) \cos(\Phi(u(x), v(x)))) dx .$$

Mathematics Subject Classification (2010): 35J47, 35J50, 49K35.

Keywords: Minimax, multiple global minima, variational methods, semilinear elliptic systems.

1. Introduction

Let S be a topological space. A function $g : S \rightarrow \mathbb{R}$ is said to be inf-compact if, for each $r \in \mathbb{R}$, the set $g^{-1}(]-\infty, r])$ is compact.

If Y is a real interval and $f : S \times Y \rightarrow \mathbb{R}$ is a function inf-compact and lower semicontinuous in S , and concave in Y , the occurrence of the strict minimax inequality

$$\sup_Y \inf_S f < \inf_S \sup_Y f$$

implies that the interior of the set A of all $y \in Y$ for which $f(\cdot, y)$ has at least two local minima is non-empty. This fact was essentially shown in [4], giving then raise to an enormous number of subsequent applications to the multiplicity of solutions for nonlinear equations of variational nature (see [7] for an account up to 2010).

In [6] (see also [5]), we realized that, under the same assumptions as above, the occurrence of the strict minimax inequality also implies the existence of $\tilde{y} \in Y$ such that the function $f(\cdot, \tilde{y})$ has at least two global minima. It may happen that \tilde{y} is unique and does not belong to the closure of A (see Example 7 of [1]).

In [8] and [12], we extended the result of [6] to the case where Y is an arbitrary convex set in a vector space. We also stress that such an extension is not possible for the result of [4]. We then started to build a network of applications of the results of [8] and [12] which touches several different topics: uniquely remotal sets in normed spaces ([8]); non-expansive operators ([9]); singular points ([10]); Kirchhoff-type problems ([11]); Lagrangian systems of relativistic oscillators ([13]); integral functional of the Calculus of Variations ([14]); non-cooperative gradient systems ([15]); variational inequalities ([16]).

The aim of this paper is to establish a further application within that network.

2. Results

The main abstract result is as follows:

Theorem 2.1. *Let X be a topological space, $(Y, \langle \cdot, \cdot \rangle)$ a real Hilbert space, $T \subseteq Y$ a convex set dense in Y and $I : X \rightarrow \mathbb{R}$, $\varphi : X \rightarrow Y$ two functions such that, for each $y \in T$, the function $x \rightarrow I(x) + \langle \varphi(x), y \rangle$ is lower semicontinuous and inf-compact. Moreover, assume that there exists a point $x_0 \in X$, with $\varphi(x_0) \neq 0$, such that*

- (a) x_0 is a global minimum of both functions I and $\|\varphi(\cdot)\|$;
- (b) $\inf_{x \in X} \langle \varphi(x), \varphi(x_0) \rangle < \|\varphi(x_0)\|^2$.

Then, for each convex set $S \subseteq T$ dense in Y , there exists $y^ \in S$ such that the function $x \rightarrow I(x) + \langle \varphi(x), y^* \rangle$ has at least two global minima in X .*

Proof. In view of (b), we can find $\tilde{x} \in X$ and $r > 0$ such that

$$I(\tilde{x}) + \frac{r}{\|\varphi(x_0)\|} \langle \varphi(\tilde{x}), \varphi(x_0) \rangle < I(x_0) + r \|\varphi(x_0)\| . \quad (2.1)$$

Thanks to (a), we have

$$I(x_0) + r\|\varphi(x_0)\| = \inf_{x \in X} (I(x) + r\|\varphi(x)\|) . \quad (2.2)$$

The function $y \rightarrow \inf_{x \in X} (I(x) + \langle \varphi(x), y \rangle)$ is weakly upper semicontinuous, and so there exists $\tilde{y} \in B_r$ such that

$$\inf_{x \in X} (I(x) + \langle \varphi(x), \tilde{y} \rangle) = \sup_{y \in B_r} \inf_{x \in X} (I(x) + \langle \varphi(x), y \rangle) , \quad (2.3)$$

B_r being the closed ball in X , centered at 0, of radius r . We distinguish two cases. First, assume that $\tilde{y} \neq \frac{r\varphi(x_0)}{\|\varphi(x_0)\|}$. As a consequence, taking into account that $r\|\varphi(x_0)\|$ is the maximum of the restriction to B_r of the continuous linear functional $\langle \varphi(x_0), \cdot \rangle$ (attained at the point $\frac{r\varphi(x_0)}{\|\varphi(x_0)\|}$ only), we have

$$\inf_{x \in X} (I(x) + \langle \varphi(x), \tilde{y} \rangle) \leq I(x_0) + \langle \varphi(x_0), \tilde{y} \rangle < I(x_0) + r\|\varphi(x_0)\| . \quad (2.4)$$

Now, assume that $\tilde{y} = \frac{r\varphi(x_0)}{\|\varphi(x_0)\|}$. In this case, due to (2.1), we have

$$\begin{aligned} \inf_{x \in X} (I(x) + \langle \varphi(x), \tilde{y} \rangle) &\leq I(\tilde{x}) + \langle \varphi(\tilde{x}), \tilde{y} \rangle = I(\tilde{x}) + \frac{r}{\|\varphi(x_0)\|} \langle \varphi(\tilde{x}), \varphi(x_0) \rangle \\ &< I(x_0) + r\|\varphi(x_0)\| . \end{aligned} \quad (2.5)$$

Therefore, from (2.2), (2.3), (2.4) and (2.5), it follows that

$$\sup_{y \in B_r} \inf_{x \in X} (I(x) + \langle \varphi(x), y \rangle) < \inf_{x \in X} \sup_{y \in B_r} (I(x) + \langle \varphi(x), y \rangle) . \quad (2.6)$$

Now, let $S \subseteq T$ be a convex set dense in Y . By continuity, we clearly have

$$\sup_{y \in B_r \cap S} \langle \varphi(x), y \rangle = \sup_{y \in B_r} \langle \varphi(x), y \rangle$$

for all $x \in X$. Therefore, in view of (2.6), we have

$$\begin{aligned} \sup_{y \in B_r \cap S} \inf_{x \in X} (I(x) + \langle \varphi(x), y \rangle) &\leq \sup_{y \in B_r} \inf_{x \in X} (I(x) + \langle \varphi(x), y \rangle) \\ &< \inf_{x \in X} \sup_{y \in B_r} (I(x) + \langle \varphi(x), y \rangle) = \inf_{x \in X} \sup_{y \in B_r \cap S} (I(x) + \langle \varphi(x), y \rangle) . \end{aligned}$$

At this point, the conclusion follows directly applying Theorem 1.1 of [12] to the restriction of the function $(x, y) \rightarrow I(x) + \langle \varphi(x), y \rangle$ to $X \times (B_r \cap S)$. \square

We now present an application of Theorem 2.1 to elliptic systems.

In the sequel, $\Omega \subseteq \mathbf{R}^n$ ($n \geq 2$) is a bounded domain with smooth boundary.

We denote by \mathcal{A} the class of all functions $H : \Omega \times \mathbf{R}^2 \rightarrow \mathbf{R}$ which are measurable in Ω , C^1 in \mathbf{R}^2 and satisfy

$$\sup_{(x, u, v) \in \Omega \times \mathbf{R}^2} \frac{|H_u(x, u, v)| + |H_v(x, u, v)|}{1 + |u|^p + |v|^p} < +\infty$$

where $p > 0$, with $p < \frac{n+2}{n-2}$ when $n > 2$.

Given $H \in \mathcal{A}$, we are interested in the problem

$$\begin{aligned} -\Delta u &= H_u(x, u, v) \text{ in } \Omega \\ -\Delta v &= H_v(x, u, v) \text{ in } \Omega \\ u &= v = 0 \text{ on } \partial\Omega, \end{aligned}$$

H_u (resp. H_v) denoting the derivative of H with respect to u (resp. v).

As usual, a weak solution of this problem is any $(u, v) \in H_0^1(\Omega) \times H_0^1(\Omega)$ such that

$$\begin{aligned} \int_{\Omega} \nabla u(x) \nabla \varphi(x) dx &= \int_{\Omega} H_u(x, u(x), v(x)) \varphi(x) dx, \\ \int_{\Omega} \nabla v(x) \nabla \psi(x) dx &= \int_{\Omega} H_v(x, u(x), v(x)) \psi(x) dx \end{aligned}$$

for all $\varphi, \psi \in H_0^1(\Omega)$.

Define the functional $I_H : H_0^1(\Omega) \times H_0^1(\Omega) \rightarrow \mathbf{R}$ by

$$I_H(u, v) = \frac{1}{2} \left(\int_{\Omega} |\nabla u(x)|^2 dx + \int_{\Omega} |\nabla v(x)|^2 dx \right) - \int_{\Omega} H(x, u(x), v(x)) dx$$

for all $(u, v) \in H_0^1(\Omega) \times H_0^1(\Omega)$.

Since $H \in \mathcal{A}$, the functional I_H is C^1 in $H_0^1(\Omega) \times H_0^1(\Omega)$ and its critical points are precisely the weak solutions of the problem. Moreover, due to the Sobolev embedding theorem, the functional $(u, v) \rightarrow \int_{\Omega} H(x, u(x), v(x)) dx$ has a compact derivative and, as a consequence, it is sequentially weakly continuous in $H_0^1(\Omega) \times H_0^1(\Omega)$.

Also, we denote by λ_1 the first eigenvalue of the Dirichlet problem

$$\begin{aligned} -\Delta u &= \lambda u \text{ in } \Omega \\ u &= 0 \text{ on } \partial\Omega. \end{aligned}$$

Our result is as follows:

Theorem 2.2. *Let $F, G \in \mathcal{A}$, with $p = \frac{2}{n-2}$ when $n > 2$, and let $K \in \mathcal{A}$, with $K(x, 0, 0) = 0$ for all $x \in \Omega$, satisfy the following conditions:*

(a₁) *one has*

$$\lim_{s^2+t^2 \rightarrow +\infty} \frac{\sup_{x \in \Omega} (|F(x, s, t)| + |G(x, s, t)|)}{s^2 + t^2} = 0;$$

(a₂) *there is $\eta \in]0, \frac{\lambda_1}{2}[$ such that*

$$K(x, s, t) \leq \eta(s^2 + t^2)$$

for all $x \in \Omega, s, t \in \mathbf{R}$;

(a₃) *one has*

$$\text{meas}(\{x \in \Omega : 0 < |F(x, 0, 0)|^2 + |G(x, 0, 0)|^2\}) > 0 \quad (2.7)$$

and

$$|F(x, 0, 0)|^2 + |G(x, 0, 0)|^2 \leq |F(x, s, t)|^2 + |G(x, s, t)|^2 \quad (2.8)$$

for all $x \in \Omega, s, t \in \mathbf{R}$;

(a₄) one has

$$\begin{aligned} \text{meas}(\{x \in \Omega : \inf_{(s,t) \in \mathbf{R}^2} (F(x,0,0)F(x,s,t) + G(x,0,0)G(x,s,t)) \\ < |F(x,0,0)|^2 + |G(x,0,0)|^2\}) > 0 . \end{aligned}$$

Then, for every convex set $S \subseteq L^\infty(\Omega) \times L^\infty(\Omega)$ dense in $L^2(\Omega) \times L^2(\Omega)$, there exists $(\alpha, \beta) \in S$ such that the problem

$$\begin{aligned} -\Delta u &= \alpha(x)F_u(x, u, v) + \beta(x)G_u(x, u, v) + K_u(x, u, v) \quad \text{in } \Omega \\ -\Delta v &= \alpha(x)F_v(x, u, v) + \beta(x)G_v(x, u, v) + K_v(x, u, v) \quad \text{in } \Omega \\ u &= v = 0 \quad \text{on } \partial\Omega \end{aligned}$$

has at least three weak solutions, two of which are global minima in $H_0^1(\Omega) \times H_0^1(\Omega)$ of the functional

$$\begin{aligned} (u, v) \rightarrow \frac{1}{2} \left(\int_{\Omega} |\nabla u(x)|^2 dx + \int_{\Omega} |\nabla v(x)|^2 dx \right) \\ - \int_{\Omega} (\alpha(x)F(x, u(x), v(x)) + \beta(x)G(x, u(x), v(x)) + K(x, u(x), v(x))) dx . \end{aligned}$$

Proof. We are going to apply Theorem 2.1, with the following choices: X is the space $H_0^1(\Omega) \times H_0^1(\Omega)$ endowed with the weak topology induced by the scalar product

$$\langle (u, v), (w, \omega) \rangle_X = \int_{\Omega} (\nabla u(x)\nabla w(x) + \nabla v(x)\nabla \omega(x)) dx ;$$

Y is the space $L^2(\Omega) \times L^2(\Omega)$ with the scalar product

$$\langle (f, g), (h, k) \rangle_Y = \int_{\Omega} (f(x)h(x) + g(x)k(x)) dx ;$$

T is $L^\infty(\Omega) \times L^\infty(\Omega)$; I is the function defined by

$$I(u, v) = \frac{1}{2} \left(\int_{\Omega} |\nabla u(x)|^2 dx + \int_{\Omega} |\nabla v(x)|^2 dx \right) - \int_{\Omega} K(x, u(x), v(x)) dx$$

for all $(u, v) \in X$; φ is the function defined by

$$\varphi(u, v) = (F(\cdot, u(\cdot), v(\cdot)), G(\cdot, u(\cdot), v(\cdot)))$$

for all $(u, v) \in X$; x_0 is the zero of X . Let us show that the assumptions of Theorem 2.1 are satisfied. First, from (2.7) and (2.8) it clearly follows, respectively, that

$$\|\varphi(0, 0)\|_Y^2 = \int_{\Omega} (|F(x, 0, 0)|^2 + |G(x, 0, 0)|^2) dx > 0$$

and that

$$\|\varphi(0, 0)\|_Y^2 \leq \|\varphi(u, v)\|_Y^2$$

for all $(u, v) \in X$. Moreover, from (a₂), thanks to the Poincaré inequality, we get

$$\int_{\Omega} K(x, u(x), v(x)) dx \leq \eta \int_{\Omega} (|u(x)|^2 + |v(x)|^2) dx \leq \frac{\eta}{\lambda_1} \int_{\Omega} (|\nabla u(x)|^2 + |\nabla v(x)|^2) dx \quad (2.9)$$

for all $(u, v) \in X$. In particular, since $K(x, 0, 0) = 0$ in Ω and $\frac{\eta}{\lambda_1} < \frac{1}{2}$, from (2.9) we infer that $(0, 0)$ is a global minimum of I in X . So, condition (a) is satisfied. Now, let us verify condition (b). To this end, set

$$P(x, s, t) = F(x, 0, 0)F(x, s, t) + G(x, 0, 0)G(x, s, t) - |F(x, 0, 0)|^2 - |G(x, 0, 0)|^2$$

for all $(x, s, t) \in \Omega \times \mathbf{R}^2$ and

$$D = \left\{ x \in \Omega : \inf_{(s,t) \in \mathbf{R}^2} P(x, s, t) < 0 \right\}.$$

By (a_4) , D has a positive measure. In view of the Scorza-Dragoni theorem, there exists a compact set $C \subset D$, with positive measure, such that the restriction of P to $C \times \mathbf{R}^2$ is continuous. Fix a point $\tilde{x} \in C$ such that the intersection of C and any ball centered at \tilde{x} has a positive measure. Choose $\tilde{s}, \tilde{t} \in \mathbf{R} \setminus \{0\}$ so that $P(\tilde{x}, \tilde{s}, \tilde{t}) < 0$. By continuity, there is $r > 0$ such that

$$P(x, \tilde{s}, \tilde{t}) < 0$$

for all $x \in C \cap B(\tilde{x}, r)$. Set

$$\gamma = \sup_{(x,s,t) \in \Omega \times [-|\tilde{s}|, |\tilde{s}|] \times [-|\tilde{t}|, |\tilde{t}|]} |P(x, s, t)|.$$

Since $F, G \in \mathcal{A}$, γ is finite. Now, choose an open set A such that

$$C \cap B(\tilde{x}, r) \subset A \subset \Omega$$

and

$$\text{meas}(A \setminus (C \cap B(\tilde{x}, r))) < -\frac{\int_{C \cap B(\tilde{x}, r)} P(x, \tilde{s}, \tilde{t}) dx}{\gamma}. \quad (2.10)$$

Finally, choose two functions $\tilde{u}, \tilde{v} \in H_0^1(\Omega)$ such that

$$\tilde{u}(x) = \tilde{s}, \quad \tilde{v}(x) = \tilde{t}$$

for all $x \in C \cap B(\tilde{x}, r)$,

$$\tilde{u}(x) = \tilde{v}(x) = 0$$

for all $x \in \Omega \setminus A$ and

$$|\tilde{u}(x)| \leq |\tilde{s}|, \quad |\tilde{v}(x)| \leq |\tilde{t}|$$

for all $x \in \Omega$. Then, taking (2.10) into account, we have

$$\begin{aligned} \langle \varphi(\tilde{u}, \tilde{v}), \varphi(0, 0) \rangle_Y - \|\varphi(0, 0)\|_Y^2 &= \int_{\Omega} P(x, \tilde{u}(x), \tilde{v}(x)) dx \\ &= \int_{C \cap B(\tilde{x}, r)} P(x, \tilde{s}, \tilde{t}) dx + \int_{A \setminus (C \cap B(\tilde{x}, r))} P(x, \tilde{u}(x), \tilde{v}(x)) dx \\ &< \int_{C \cap B(\tilde{x}, r)} P(x, \tilde{s}, \tilde{t}) dx + \gamma \text{meas}(A \setminus (C \cap B(\tilde{x}, r))) < 0. \end{aligned}$$

This shows that (b) is satisfied. Finally, fix $\alpha, \beta \in L^\infty(\Omega)$. Clearly, the function

$$(x, s, t) \rightarrow \alpha(x)F(x, s, t) + \beta(x)G(x, s, t) + K(x, s, t)$$

belongs to \mathcal{A} , and so the functional

$$(u, v) \rightarrow I(u, v) + \langle \varphi(u, v), (\alpha, \beta) \rangle_Y$$

is sequentially weakly lower semicontinuous in X . Let us show that it is coercive. Set

$$\theta = \max \{ \|\alpha\|_{L^\infty(\Omega)}, \|\beta\|_{L^\infty(\Omega)} \}$$

and fix $\epsilon > 0$ so that

$$\epsilon < \frac{1}{\theta} \left(\frac{\lambda_1}{2} - \eta \right). \quad (2.11)$$

By (a_1) , there is $c_\epsilon > 0$ such that

$$|F(x, s, t)| + |G(x, s, t)| \leq \epsilon(|s|^2 + |t|^2) + c_\epsilon$$

for all $(x, s, t) \in \Omega \times \mathbf{R}^2$. Then, for each $u, v \in H_0^1(\Omega)$, recalling (2.9), we have

$$\begin{aligned} & I(u, v) + \langle \varphi(u, v), (\alpha, \beta) \rangle_Y \\ & \geq \left(\frac{1}{2} - \frac{\eta}{\lambda_1} \right) \int_{\Omega} (|\nabla u(x)|^2 + |\nabla v(x)|^2) dx \\ & \quad - \int_{\Omega} |\alpha(x)F(x, u(x), v(x)) + \beta(x)G(x, u(x), v(x))| dx \\ & \geq \left(\frac{1}{2} - \frac{\eta}{\lambda_1} \right) \int_{\Omega} (|\nabla u(x)|^2 + |\nabla v(x)|^2) dx - \theta\epsilon \int_{\Omega} (|u(x)|^2 + |v(x)|^2) dx - \theta c_\epsilon \text{meas}(\Omega) \\ & \geq \left(\frac{1}{2} - \frac{\eta}{\lambda_1} - \frac{\theta\epsilon}{\lambda_1} \right) \int_{\Omega} (|\nabla u(x)|^2 + |\nabla v(x)|^2) dx - \theta c_\epsilon \text{meas}(\Omega). \end{aligned}$$

Notice that, in view of (2.11), we have $\frac{1}{2} - \frac{\eta}{\lambda_1} - \frac{\theta\epsilon}{\lambda_1} > 0$, and so

$$\lim_{\|(u,v)\|_X \rightarrow +\infty} (I(u, v) + \langle \varphi(u, v), (\alpha, \beta) \rangle_Y) = +\infty,$$

as claimed.

In particular, this also implies that the functional $(u, v) \rightarrow I(u, v) + \langle \varphi(u, v), (\alpha, \beta) \rangle_Y$ is weakly lower semicontinuous, by the Eberlein-Smulyan theorem. Thus, the assumptions of Theorem 2.1 are satisfied. Therefore, for each convex set $S \subseteq L^\infty(\Omega) \times L^\infty(\Omega)$ dense in $H_0^1(\Omega) \times H_0^1(\Omega)$, there exists $(\alpha, \beta) \in S$, such that the functional

$$\begin{aligned} & (u, v) \rightarrow \frac{1}{2} \left(\int_{\Omega} |\nabla u(x)|^2 dx + \int_{\Omega} |\nabla v(x)|^2 dx \right) \\ & \quad - \int_{\Omega} (\alpha(x)F(x, u(x), v(x)) + \beta(x)G(x, u(x), v(x)) + K(x, u(x), v(x))) dx \end{aligned}$$

has at least two global minima in $H_0^1(\Omega) \times H_0^1(\Omega)$. Finally, by Example 38.25 of [17], the same functional satisfies the Palais-Smale condition, and so it admits at least three critical points, in view of Corollary 1 of [3]. The proof is complete. \square

Remark 2.3. We are not aware of known results close enough to Theorem 2.2 in order to do a proper comparison. This sentence also applies to the case of single equations, that is to say when F, G, K depend on x and s only. For an account on elliptic systems, we refer to [2].

Among the various corollaries of Theorem 2.2, we wish to stress the following ones:

Corollary 2.4. *Let $K \in \mathcal{A}$, with $K(x, 0, 0) = 0$ for all $x \in \Omega$, satisfy condition (a_2) . Moreover, let $\Phi : \mathbf{R}^2 \rightarrow \mathbf{R}$ be a non-constant C^1 function, with $\Phi(0, 0) = 0$, belonging to \mathcal{A} , with $p = \frac{2}{n-2}$ when $n > 2$.*

Then, for every convex set $S \subseteq L^\infty(\Omega) \times L^\infty(\Omega)$ dense in $L^2(\Omega) \times L^2(\Omega)$, there exists $(\alpha, \beta) \in S$ such that the problem

$$\begin{aligned} -\Delta u &= (\alpha(x) \cos(\Phi(u, v)) - \beta(x) \sin(\Phi(u, v)))\Phi_u(u, v) + K_u(x, u, v) \text{ in } \Omega \\ -\Delta v &= (\alpha(x) \cos(\Phi(u, v)) - \beta(x) \sin(\Phi(u, v)))\Phi_v(u, v) + K_v(x, u, v) \text{ in } \Omega \\ u &= v = 0 \text{ on } \partial\Omega \end{aligned}$$

has at least three weak solutions, two of which are global minima in $H_0^1(\Omega) \times H_0^1(\Omega)$ of the functional

$$\begin{aligned} (u, v) \rightarrow & \frac{1}{2} \left(\int_{\Omega} |\nabla u(x)|^2 dx + \int_{\Omega} |\nabla v(x)|^2 dx \right) \\ & - \int_{\Omega} (\alpha(x) \sin(\Phi(u(x), v(x))) + \beta(x) \cos(\Phi(u(x), v(x))) + K(x, u(x), v(x))) dx . \end{aligned}$$

Proof. It suffices to apply Theorem 2.2 to the functions $F, G : \mathbf{R}^2 \rightarrow \mathbf{R}$ defined by

$$\begin{aligned} F(s, t) &= \sin(\Phi(s, t)) , \\ G(s, t) &= \cos(\Phi(s, t)) \end{aligned}$$

for all $(s, t) \in \mathbf{R}^2$. □

Corollary 2.5. *Let $F, G : \mathbf{R} \rightarrow \mathbf{R}$ belong to \mathcal{A} , with $p = \frac{2}{n-2}$ when $n > 2$. Moreover, assume that F, G are twice differentiable at 0 and that*

$$\begin{aligned} \lim_{|s| \rightarrow +\infty} \frac{|F(s)| + |G(s)|}{s^2} &= 0 , \\ 0 < |F(0)|^2 + |G(0)|^2 &= \inf_{s \in \mathbf{R}} (|F(s)|^2 + |G(s)|^2) , \\ F''(0)F(0) + G''(0)G(0) &< 0 . \end{aligned} \tag{2.12}$$

Then, for every convex set $S \subseteq L^\infty(\Omega) \times L^\infty(\Omega)$ dense in $L^2(\Omega) \times L^2(\Omega)$, there exists $(\alpha, \beta) \in S$ such that the problem

$$\begin{aligned} -\Delta u &= \alpha(x)F'(u) + \beta(x)G'(u) \text{ in } \Omega \\ u &= 0 \text{ on } \partial\Omega \end{aligned}$$

has at least three weak solutions, two of which are global minima in $H_0^1(\Omega)$ of the functional

$$u \rightarrow \frac{1}{2} \int_{\Omega} |\nabla u(x)|^2 dx - \int_{\Omega} (\alpha(x)F(u(x)) + \beta(x)G(u(x))) dx .$$

Proof. We apply Theorem 2.2 taking $K = 0$. Since 0 is a global minimum of the function $|F(\cdot)|^2 + |G(\cdot)|^2$, we have

$$F'(0)F(0) + G'(0)G(0) = 0$$

and so, in view of (2.12), 0 is a strict local maximum for the function

$$F(\cdot)F(0) + G(\cdot)G(0).$$

Hence, (a_4) is satisfied and Theorem 2.2 gives the conclusion. \square

References

- [1] Cabada, A., Iannizzotto, A., *A note on a question of Ricceri*, Appl. Math. Lett., **25**(2012), 215-219.
- [2] de Figueiredo, D.G., *Semilinear elliptic systems: existence, multiplicity, symmetry of solutions*, Handbook of Differential Equations: Stationary Partial Differential Equations, Vol. V, 1-48, Handb. Differ. Equ., Elsevier/North-Holland, 2008.
- [3] Pucci, P., Serrin, J., *A mountain pass theorem*, J. Differential Equations, **60**(1985), 142-149.
- [4] Ricceri, B., *On a three critical points theorem*, Arch. Math., **75**(2000), 220-226.
- [5] Ricceri, B., *Well-posedness of constrained minimization problems via saddle-points*, J. Global Optim., **40**(2008), 389-397.
- [6] Ricceri, B., *Multiplicity of global minima for parametrized functions*, Rend. Lincei Mat. Appl., **21**(2010), 47-57.
- [7] Ricceri, B., *Nonlinear eigenvalue problems*, in "Handbook of Nonconvex Analysis and Applications" D. Y. Gao and D. Motreanu eds., 543-595, International Press, 2010.
- [8] Ricceri, B., *A strict minimax inequality criterion and some of its consequences*, Positivity, **16**(2012), 455-470.
- [9] Ricceri, B., *A range property related to non-expansive operators*, Mathematika, **60**(2014), 232-236.
- [10] Ricceri, B., *Singular points of non-monotone potential operators*, J. Nonlinear Convex Anal., **16**(2015), 1123-1129.
- [11] Ricceri, B., *Energy functionals of Kirchhoff-type problems having multiple global minima*, Nonlinear Anal., **15**(2015), 130-136.
- [12] Ricceri, B., *On a minimax theorem: an improvement, a new proof and an overview of its applications*, Minimax Theory Appl., **2**(2017), 99-152.
- [13] Ricceri, B., *Another multiplicity result for the periodic solutions of certain systems*, Linear Nonlinear Anal., **5**(2019), 371-378.
- [14] Ricceri, B., *Miscellaneous applications of certain minimax theorems II*, Acta Math. Vietnam., **45**(2020), 515-524.
- [15] Ricceri, B., *An alternative theorem for gradient systems*, Pure Appl. Funct. Anal. (to appear).
- [16] Ricceri, B., *A remark on variational inequalities in small balls*, J. Nonlinear Var. Anal., **4**(2020), 21-26.
- [17] Zeidler, E., *Nonlinear functional analysis and its applications*, vol. III, Springer-Verlag, 1985.

Biagio Ricceri
Department of Mathematics and Informatics,
University of Catania,
Viale A. Doria 6,
95125 Catania, Italy
e-mail: ricceri@dmf.unict.it