

# UNIVERSITÀ DEGLI STUDI DI CATANIA

DOTTORATO DI RICERCA IN SCIENZE DELLE PRODUZIONI ANIMALI

(XXIV Ciclo)

DIPARTIMENTO DI SCIENZE AGRONOMICHE, AGROCHIMICHE  
E DELLE PRODUZIONI ANIMALI

---

Doctoral thesis

PATRICIA LÓPEZ ANDRÉS

## **Bioavailability and antioxidant effect of dietary phenolic compounds in lamb tissues**

Coordinator: Prof. Marcella Avondo

Tutor: Prof. Alessandro Priolo

Years 2008 - 2011



## ***List of Publications***

---

ü López-Andrés P., Luciano G., Vasta V., Gibson TM., Biondi L., Priolo A., Mueller-Harvey I. Bioavailability and antioxidant effect of dietary tannins in lamb tissues. Submitted for publication to *British Journal of Nutrition*.

ü Luciano G., Moloney AP., Priolo A., Röhrle, Vasta V., Biondi L., López Andrés P., Grasso S., Monahan FJ. 2011. Vitamin E and polyunsaturated fatty acids in bovine muscle and the oxidative stability of beef from cattle receiving grass or concentrate-based rations. *Journal of Animal Science* 89, 3759-3768.

ü Luciano G., Vasta V., Monahan FJ., López-Andrés P., Biondi L., Lanza M., Priolo A. 2011. Antioxidant status, colour stability and myoglobin resistance to oxidation of *longissimus dorsi* muscle from lambs fed a tannin-containing diet. *Food Chemistry* 124, 1036-1042.

## ***Foreigner Institutions and Laboratories Visited***

---

ü School of Agriculture, Food Science and Veterinary Medicine, University College of Dublin, Ireland. (February – April 2010). Tutor: Prof. Frank Monahan.

ü Chemistry & Biochemistry Laboratory, Food Production and Quality Research Division, School of Agriculture, Policy and Development, University of Reading, United Kingdom. (January – March 2011). Tutor: Dr. Irene Mueller-Harvey.

## *CONTENTS*

---

➤ Phenolic compounds	6
➤ Effects of phenolic compounds in monogastrics	11
➤ Bioavailability of phenolic compounds in monogastrics	13
➤ Effects of phenolic compounds in polygastrics	18
➤ Bioavailability of phenolic compounds in polygastrics	23
➤ Aim of the experiments	24
➤ Materials and methods	
1. Animal and diets	25
2. Sampling	26
3. Preparation of feed samples and purification of phenolic compounds by SPE	26
4. Preparation of lamb tissues samples and purification of phenolic compounds by SPE	27
5. Feed and tissue extracts LC-MS analysis	28
6. Antioxidant status of liver and plasma samples treated with or without SPE	
6.1. Folin-Ciocalteu assay in RAW-samples	29
6.2. Folin-Ciocalteu assay in SPE-samples	30
6.3. Ferric reducing antioxidant power (FRAP assay) of RAW-samples	30
6.4. Ferric reducing antioxidant power (FRAP assay) of SPE-samples	32
7. Statistical Analysis	32
➤ Results	
1. Analysis of feed samples by LC-MS	33
2. Analysis of animal tissues samples by LC-MS	39
3. Folin-Ciocalteu assay in lamb tissues samples	46
4. Ferric reducing antioxidant power (FRAP) assay in lamb tissues samples	47
➤ Discussions	49
➤ Conclusion	57
➤ References	58




## PHENOLIC COMPOUNDS

---

Phenolic compounds are organic compounds distributed all over the plant kingdom and they are products of the secondary metabolism of plants (Haslam, 1989). Widely distributed in the leaves, seeds, bark and flowers of plants, phenolic compounds afford protection against ultraviolet radiation, pathogens, and herbivores in plants (Harbone and Williams, 2000). Also they contribute to the colour and sensory characteristics for fruits and vegetables. More over 4,000 flavonoids have been identified in plants and phenolic compound have been used in the chemotaxonomic characterisation of plant species or of particular organs of plant tissues (Harborne and Green, 1980).

Polyphenols' chemical structure consists in one or more aromatic rings which can contain hydroxyl (Rice-Evans, 1996) methoxy (Dugas *et al.*, 2000) and/or glycosidic groups (Harborne 1994). Polyphenols also can be associated, with other compounds, such as organic acids (Tapiero *et al.*, 2002), amines (Vitaglione *et al.*, 2004), lipids (Yang *et al.*, 2000) and other phenols (Bravo, 1998). Depending on their chemical structure, polyphenolic compounds can be divided into different groups as shown in Table 1.

Fig.1 Structure of a flavan skeleton




Flavonoids constitute the main and extensively group of phenolic compounds present in plant kingdom (Koes *et al.*, 2005). They present a C6-C3-C6 carbon skeleton (Rice-Evans *et al.*, 1996) and their structures consist in two benzene rings A and B, bonded through three carbons, which form a closed pyran ring (ring C) (Fig. 1).

Table 1. Chemical features of phenolic compounds


<i>Classes of phenolic compounds</i>	<i>Carbon skeletons</i>	<i>Examples</i>
Simple phenolics, benzoquinones	C6	Catechol, pyrogallol
Hydroxybenzoic acids	C6-C1	Gentisic acid
Acetophenones, phenylacetic acids	C6-C2	4-hydroxyphenylacetic acid
Hydroxycinnamic acids	C6-C3	Caffeic acid, cinnamic acid,
Naphthoquinones	C6-C4	Juglone
Xanthones	C6-C1-C6	Mangiferin
Stilbenes, anthraquinones	C6-C2-C6	Resveratrol, picetamol
Flavonoids, isoflavonoids	C6-C3-C6	
anthocyanin		Cyanidin
flavanols		Catechin, gallicocatechin
flavanones		Naringenin
flavones		Luteolin, apigenin
flavonols		Kaempferol, quercetin
isoflavones		Genistein, daidzein
Lignans, neolignans	(C6-C3) <sub>2</sub>	Pinoresinol
Biflavonoids	(C6-C3-C6) <sub>2</sub>	Amentoflavone
Lignins	(C6-C3) <sub>n</sub>	
Condensed tannins	(C6-C3-C6) <sub>n</sub>	Profisetinidin, Procyanidin

Flavonoids are themselves divided into several subclasses, depending on the oxidation state of the central pyran ring and the position of the ring B (Figure 2). In each flavonoids subclass, numerous possible linkages between flavonoids and different sugars (as glucose, galactose, rhamnose, xylose), disaccharides (such as rutose) and methoxy groups can occur (Cook *et al.*, 1996). For example, the flavonol kaempferol can be linked with glucose in the 3 carbon to give astragalol, or be bonded with two rhamnose in the 3 and 7 carbons to give kaempferitrin, or linked with a robinose in 3-C and a rhamnose in 7 to give robinin, etc.


Figure 2. Examples of flavonoids compounds




Cyanidin (Anthocyanin)




Catechin (Flavanol)




Naringenin (Flavanone)



Apigenin (Flavone)



Kaempferol (Flavonol)



Genistein (Isoflavone)



Flavonoids are the monomers of tannins, another group of polyphenolic compounds widespread in plant kingdom (Yoshida *et al.*, 2000). Tannins can be classified into two groups: (i) hydrolysable tannins, which contain gallic acids and their oxidation products (Mueller-Harvey, 2001) and can be classified in ellagitannins and gallotannins (ii) condensed tannins or proanthocyanidins, which are oligomers and polymers of polyhydroxyflavan-3-ols (Porter, 1992). Depending on the chemical structure of the monomer unit, proanthocyanidins can be divided also in different classes (Table 2).

Table 2. Proanthocyanidin nomenclature: types of proanthocyanidin and suggested names for the monomer units

<i>Proanthocyanidin class</i>	<i>Monomer unit</i>	<i>Substitution pattern</i>						
		3	5	7	8	3'	4'	5'
Proapigeninidin	Apigeniflavan	H	OH	OH	H	H	OH	H
Prolyuteolinidin	Luteoliflavan	H	OH	OH	H	OH	OH	H
Protrictetinidin	Trictetiflavan	H	OH	OH	H	OH	OH	OH
Propelargonidin	Afzelechin	OH	OH	OH	H	H	OH	H
Procyanidin	Catechin	OH	OH	OH	H	OH	OH	H
Prodelphinidin	Gallocatechin	OH	OH	OH	H	OH	OH	OH
Proguibourtinidin	Guibourtinidol	OH	H	OH	H	H	OH	H
Profisetinidin	Fisetinidol	OH	H	OH	H	OH	OH	H
Prorobinetinidin	Robinetinidol	OH	H	OH	H	OH	OH	OH
Proteracacinidin	Oritin	OH	H	OH	OH	H	OH	H
Promelacacinidin	Prosopin	OH	H	OH	OH	OH	OH	H

Regardless of their structural diversity, all polyphenolic compounds are reducing agents (Rice-Evans *et al.*, 1996). This antioxidant activity depends on the chemical structure of the polyphenols (Balasundram *et al.*, 2006), first of all, in the degree of hydroxylation and the positions of the –OH groups in the B ring (Rice-Evans *et al.*, 1996) (Fig. 1). The protective effects of flavonoids in biological systems are ascribed

to their capacity to transfer electrons free radicals, chelate metal catalysts, activate antioxidant enzymes, reduce alpha-tocopherol radicals, and inhibit oxidases (Heim *et al.*, 2002). Some studies have reported that the antioxidant activity of polyphenolic compounds decrease when these polyphenols are linked to methyl and/or glucosidic groups (Cao *et al.*, 1997).

Tannins, which are highly polymerized, possess also diverse biological effects, as to be metal ion chelators (McDonald *et al.* 1996) and protein precipitating agents (Santos-Buelga *et al.* 2000). Tannins, like all phenolic compounds are antioxidant agents, but its antioxidant activity has not been studied so greatly than for simple phenols. Hagerman *et al.*, (1998) showed in an *in vitro* study that tannins could be 15-30 times more effective at quenching peroxy radicals than simple phenols. This effect could be due to the fact that tannins are highly polymerized and can contain more –OH groups in their structure, than simple phenols.

Phenolic compounds are present daily in humans and animals diets. Phenolic compounds' effect and their bioavailability have been studied in depth in the last 20 years. The study of bioavailability and effects of phenolic compounds in humans has been focused to relate the consumption of phenols with the decrease of several diseases. In animals, to achieve final products (meat, dairy, eggs) with higher quality, which humans will consume subsequently.

Below, the last studies related with the consumption of phenols, their effects and bioavailability in monogastric and polygastric animals.

## ***EFFECTS OF PHENOLIC COMPOUNDS IN MONOGASTRICS***

---

Phenolic compounds are regular constituents of human food. The main sources of phenolic compounds in human diet are fruits (Cieslik *et al.*, 2006), beverages such as tea (McKay *et al.*, 2002), cocoa (Holt *et al.*, 2002), coffee (Nardini *et al.*, 2002), wine (Wang *et al.*, 2002) and fruit juices (Abad-García *et al.*, 2007), vegetables (Vinson *et al.*, 1998), cereals (Naczka *et al.*, 2006), fruits (Cieślak *et al.*, 2006), chocolate (Sanbongi *et al.*, 1997) and legume seeds (Troszynska *et al.*, 2006). The antioxidant activity of polyphenols present in food is very variable. In table 3 is shown different polyphenol food sources and Trolox equivalent antioxidant activities (TEAC) of each flavonoid. (Heim *et al.*, 2002 and Rice-Evans *et al.*, 1997).

The consumption of phenol-rich foods or beverages is associated with beneficial effects for human health (Hollman *et al.*, 1997). Phenolic compounds are related to the prevention of diseases such as cancer (Ramos, 2008), stroke (Fraser *et al.*, 2007), coronary heart (Ghosh, 2009), osteoporosis (Yamaguchi, 2002), neurodegenerative diseases (Ramassamy, 2006), diabetes (Dembinska-Kiec *et al.*, 2008).

Phenolic compounds have different effects in the organism, they can act as anti-allergenic (Ksouri *et al.*, 2007), anti-atherogenic (Sultana *et al.*, 2008), anti-inflammatory (Cook *et al.*, 1996), anti-microbial (Cowan 1999), antioxidant (Moreno *et al.*, 2006), anti-thrombotic (Stangl *et al.*, 2007), cardioprotective (Zern *et al.*, 2005) and vasodilators compounds (Balasundram *et al.*, 2006). The absorption of polyphenols in humans and animals can increase the vasodilatation (Diebolt *et al.*, 2001), reduce the vasoconstriction of endothelial cells (Aldini *et al.*, 2003) and decrease the platelet aggregation (Freedman *et al.*, 2001). In short, polyphenols can reduce the angiogenesis (Yuasa *et al.*, 2002) and the atherogenesis (Kaliora *et al.*, 2005) of animals fed with a rich phenol diet.

Table 3. Dietary sources of flavonoids and Trolox equivalent antioxidant activities (TEAC). Higher TEAC values reflect greater antioxidant capability.

<i>Classes of phenolic compounds</i>	<i>Dietary sources</i>	<i>Antioxidant activity TEAC (mM)</i>
<i>Hydroxycinnamic acids</i>		
Caffeic acid	White grapes, olive, asparagus	1.3 ± 0.01
Chlorogenic acid	Apple, pear, cherry, tomato, peach	1.3 ± 0.02
Ferulic acid	Grains, tomato, cabbage, asparagus	1.9 ± 0.02
<i>Anthocyanidin</i>		
Cyanidin	Grapes, raspberry, strawberry	4.4 ± 0.12
Apigenidin	Colored fruits	2.35 ± 0.01
<i>Flavanols</i>		
(Epi)catechin	Black grapes, red wine, tea	2.4 ± 0.01
Epigallocatechin gallate	Teas	4.8 ± 0.06
<i>Flavanones</i>		
Naringenin	Citrus, grapefruit	1.5 ± 0.05
Hesperidin	Orange juice	1.0 ± 0.03
Taxifolin	Citrus fruit	1.9 ± 0.03
<i>Flavones</i>		
Apigenin	Parsley, celery	1.5 ± 0.08
Rutin	Onion, apple skin, broccoli, berries	2.4 ± 0.02
Luteolin	Lemon, red pepper, olive, celery	2.1 ± 0.05
<i>Flavonols</i>		
Kaempferol	Endivies, leek, broccoli, grapefruit	1.3 ± 0.08
Quercetin	Tomato, lettuce, broccoli, olive oil	4.7 ± 0.10
<i>Isoflavones</i>		
Genistein	Soybean	2.9 ± 0.01
Daidzein	Soybean	1.25 ± 0.01

## **BIOAVAILABILITY OF PHENOLIC COMPOUNDS IN MONOGASTRIC**

---


It has been suggested that the different chemical structures of the phenolic compounds, which are ingested by animals as part of their diets, are responsible for generating different effects in terms of their absorption and metabolism (Mueller-Harvey, 2006). For example, phenols differ in their basic structures (Urquiaga *et al.*, 2000), degree and pattern of glycosylation (D'Archivio *et al.*, 2007), acylation (Stevenson *et al.*, 2006), polymerisation or molecular size (Wollgast *et al.*, 2000) and solubility (Manach *et al.*, 2004). All of these factors are likely to affect the absorption of each phenolic compound. Due to the different chemical structures of phenolic compounds, it is difficult to describe single metabolic pathways in the organism. The bioavailability of phenols in mammals has been focused in humans (Scalbert *et al.*, 2000, Manach *et al.*, 2005) and in rats (Carbonaro *et al.*, 2001, Silberberg *et al.*, 2006). Polyphenols, depending on their chemical structure, can go by different metabolic pathways in human organism as explained above (see also Figure 3):

1. For polyphenols glycosylated, that it is the main form in which polyphenols are present in foods and beverages, the first step for their absorption is to remove the sugar by glycosidase enzymes. These enzymes can be present in food (enzymes endogenous) and in the gastrointestinal mucosa. It is known that after polyphenols deglycosylation, conjugation reactions such as glucuronidation and methylation, can occur in the jejunal and ileal sections of the small intestine (Spencer *et al.*, 1999).
2. Flavanols as (-)-epicatechin can pass directly through biological membranes and be absorbed without deconjugation or hydrolysis.
3. Phenolic acid esters, i.e. chlorogenic acid, are not absorbed in the small intestine. These compounds reach the colon where they are metabolised by the esterases enzymes. These enzymes are able to break the ester bond between phenol compound and carboxylic acid, in the case of chlorogenic acid (Kroon *et al.*, 1996).
4. Hydroxycinnamic acids as free aglycones are rapidly absorbed directly from small intestine.

5. With regard to proanthocyanidins, contradictory results studies about their bioavailability have been provided. It was thought that they were not absorbed in the small intestine due to their high molecular weight and in this way they were directly eliminated. However, Déprez *et al.*, 2000 showed that although proanthocyanidins were not absorbed through the gut barrier, they could be degraded by the colonic microflora into low molecular weight aromatic acids, which could be absorbed. Furthermore, there are some studies that showed that proanthocyanidins dimmers can be absorbed and identified in human and rat plasma (Holt *et al.*, 2002 and Prasain *et al.* 2009 respectively).

Phenolic compounds that have been absorbed through gut barrier are widely conjugated by enzymes present in different tissues. The conjugation by glucuronidation occurs mainly in liver, although it can also occur in intestine, kidney and human colon (Tapiero *et al.*, 2002). Methylation is undertaken by enzymes found in a wide range of tissues (Lambert *et al.*, 2003) and sulphatation is carried out by the enzyme phenol sulfotransferases found in many tissues as liver and also in the colon, Stahl *et al.*, 2002, Cappiello *et al.*, 1990). These new conjugated phenolic compounds can (i) be excreted in the bile and return to the small intestine, (ii) reach the kidneys and be eliminated in urine (Shahrzad *et al.*, 1998) or (iii) be transported by plasma till other animal tissues. It has to be noted that phenols circulate in plasma as glucuronidated or sulphated compounds and rarely as free aglycones (Manach *et al.*, 1998; Carbonaro *et al.*, 2001).

Figure 3. Possible routes for consumed polyphenols in humans (Scalbert and Williamson 2000).



Polyphenols that have not been absorbed through gut barrier or polyphenols re-excreted in the bile can reach the colon where they are metabolised by colonic microflora. In this way, flavonoid glycosides or glucuronides can be hydrolysed and the aglycone free form can be absorbed. In the case of free aglycones that reach the colon, these can be hydrolysed to low molecular weight and absorbed through the colonic barrier.

It should be noticed that the concentration of polyphenols is usually very low in plasma after their dietary administration. Most dietary polyphenols are quickly eliminated in both urine and bile after ingestion. The elimination depends also on the chemical structure of polyphenols and some studies have showed that the quantities of polyphenols found intact in urine vary from one phenolic compound to another (Scalbert *et al.* 2000). In human plasma, a post-prandial peak is observed 1–2 h after ingestion of various flavonols and flavanols but is longer for isoflavones and other polyphenols only absorbed after degradation by the colon microflora (Scalbert *et al.*, 2000). To keep a high concentration in plasma, it has to repeat ingestion of the polyphenols, as it has been observed with volunteers consuming tea every 2 h (Van Het Hof *et al.*, 1999).

It is well known that some simple flavonoids or their metabolites can be detected in different mammalian tissues not involved in phenol metabolism and excretion. Kalt *et al.*, (2008) demonstrated that when pigs are fed with an enriched blueberry fed, anthocyanins from the diet can be found also in eyes and brain region (cortex and cerebellum) of the pigs. Also, it is known that anthocyanins have been associated with improvements in vision (Matsumoto *et al.*, 2006) and with cancer chemoprevention (Hou *et al.*, 2004). In another study, Bieger *et al.*, (2008) gave to pigs quercetin aglycone as single dose the same day of slaughtering (expt. 1) or mixed into the regular meals of pigs for 4 weeks (expt. 2). In both experiments quercetin was found in muscle *longissimus dorsi* in very low quantity. Bieger also demonstrated that pigs did not accumulate quercetin in most tissues after a long-term of dietary flavonol intake (expt. 2), compared with a single treatment (expt. 1) and that only organs involved in flavonol metabolism and excretion, as small intestine, liver, and kidneys, contained significantly higher flavonol concentrations than plasma in both experiments.

De Boer *et al.*, (2005) identified for the first time quercetin and quercetin metabolites in diverse rat tissues as lung, testes, heart, fat, bones and brain, besides the organs involved directly in phenol metabolism. More recently, Juan *et al.*, (2010) studied the bioavailability of *trans*-resveratrol after an intravenous administration in rats. The results reported that *trans*-resveratrol aglycone was identified in kidney, lungs, liver, testis and brain.

In other studies diverse polyphenols have been identified in different animal tissues. Ha Park *et al.*, (2009) fed tilapia fish with an enriched quercetin diet for 15 weeks. After trial term, quercetin aglycone was identified and quantified in liver and whole body mixture sample. Saitoh *et al.*, 2004 report that when laying hens were fed with an experimental diet containing soy isoflavone-glycosides, most of the isoflavones (daidzein, glycitein, and genistein) and a metabolite, equol, were present in blood and egg yolk in conjugated form. Also Nimalaratne *et al.*, 2011 have shown that when hens are fed with an enriched wheat and corn diet, gallic acid and trace of ferulic acid were found in egg yolk.

In humans, the identification of phenols has been focused on plasma samples. Paganga and Rice-Evans (1997) described the evidence for the absorption of flavonoids and their presence in human plasma in the glycosylated form. The phenol compounds identified were rutin, other quercetin glycosides and phloridzin. Day *et al.*, (2001) showed that after consumption of onions (a rich source of flavonoid glucosides) the main circulating compounds identified in plasma were quercetin-3-glucuronide, 3'-methyl-quercetin-3-glucuronide and quercetin-3'-sulfate.

On the other hand, only few studies have reported the identification of proanthocyanidins in animal tissues. Baba *et al.*, (2002) showed the bioavailability of procyanidin B2 [epicatechin-(4 $\beta$ -8)-epicatechin] in rats. After procyanidin B2 administration, it was absorbed and excreted in urine, and a portion of the PB2 was degraded to (-)-epicatechin and to the metabolized conjugated and/or methylated (-)-epicatechin internally in the rat. Prasain *et al.*, 2009 reported that rats fed with grape seed extracts orally, monomeric catechins, their methylated metabolites, and proanthocyanidins up to trimmers were detected in blood samples. Prasain also



reported that (+)-catechin and (-)-epicatechin were identified in the brain conclusively.

In humans Holt *et al.*, (2002) showed that after cocoa consumption, procyanidin dimer, as well as the flavonol monomers epicatechin and catechin, can be absorbed into the circulation.

More studies are required to understand better the bioavailability of proanthocyanidins in animals.

## ***EFFECTS OF PHENOLIC COMPOUNDS IN POLYGASTRICS***

---

Polygastric animals, also known as ruminants, have a different digestive system to monogastric animals. Ruminants have four different stomachs: rumen, omasum, reticulum and abomasum. In the rumen it is possible to find high quantities of microflora and microfauna like bacteria (Tajima *et al.*, 1999), protozoa (Dehority, 1993) and fungi (Barr, 1988). These microorganisms have a symbiotic relationship with the animals (Russell and Wilson, 1998), because the ruminants give an optimal habitat for the growth of the microorganisms and the microorganisms can provide protein, vitamins and short-chain organic acids for ruminants (Russell and Rychlik, 2001).

Polyphenols consumption by ruminants is highly dependent on the type of the diet. There are different typologies of ruminant feeding systems, i.e. grassland, herbage, silage, concentrate or a mixture between them. It is well known that fresh herbage contain high quantity of vitamins, polyunsaturated fatty acids, antioxidants compounds like tocopherols, carotenoids, ascorbic acid and phenolic compounds (Wood & Enser, 1997). Also it has been widely studied that the consumption of grass or fresh herbage by ruminants, improve qualitative aspects in final products as meat and dairy compare to animals fed a concentrate or silage diet (Nozière *et al.*, 2006). The consumption of grass or concentrate feeds can affect animals performance (Steen *et al.*, 2003), meat color and flavor (Priolo *et al.*, 2001), fatty acid composition in meat and intramuscular fat (Realini *et al.*, 2004, French *et al.*, 2000 respectively), vitamin E concentration in muscle (Turner *et al.*, 2002) and lipid oxidation in meat (Luciano *et al.*, 2001b). The feeding system can affect also to the composition and quality of milk (Morand-Fehr *et al.*, 2007), sensory quality of dairy products (Martin *et al.*, 2005) etc.

It is known that some antioxidant compounds present in the pasture can be transferred to animal tissues. In sheep and goats, the main carotenoid in plasma and adipose tissue is lutein, whereas cattle can also store  $\beta$ -carotene (Yang *et al.*, 1992).

Yang *et al.*, (2002) demonstrated higher concentrations and significant differences of  $\alpha$ -tocopherol and  $\beta$ -carotene compounds in liver from pasture-fed cattle compared to liver from grain-fed cattle. Prache *et al.*, (2003) showed higher plasma carotenoid content for grass-fed lambs, than for stall-fed, long-stall finished or short-stall finished lambs. La Terra *et al.*, (2010) showed that when cows are fed with increasing proportion of fresh forages in their diet, a progressive increase of  $\alpha$ -tocopherol and  $\beta$ -carotene are found in plasma. Vitamin E has also been found in higher concentrations in muscle from cattle fed with a grass diet compared to muscle from cattle fed concentrate feeds (Reailini *et al.*, 2004; Luciano *et al.*, 2011a). However, there is a lack of clear information about the bioavailability of dietary phenolic compounds in animals. Moreover, the possible antioxidant mechanisms of dietary phenolic compounds in animal tissues, whether direct or indirect is still unclear. Given the large variability in the concentration of phenolic compounds in different feedstuffs used for livestock feeding, it could be of great interest to study their potential biological effects when ingested by ruminants.


Among the different phenolic compounds, tannins have been particularly studied in ruminant animals for their effects on animal digestion and performance (Makkar, 2003), for their high antioxidant activity and for their potential effects on products' quality such as meat, milk and dairy products.

It is known that many hydrolysable tannins can be metabolised by microorganisms in the rumen (Brooker *et al.*, 1994, Nelson *et al.*, 1995, Skene *et al.*, 1995, Sly *et al.*, 1997) to gallic acid, pyrogallol and other products that are potentially toxic to ruminants (Goel *et al.*, 2005).

However, it is unclear to what extent the microbial organisms degrade or metabolise condensed tannins. Several publications reported that condensed tannins cannot be degraded (Makkar *et al.* 1995a, Getachew *et al.* 2008) by ruminal microorganisms. In contrast, Perez-Maldonado and Norton (1996) suggested that condensed tannins could be absorbed or degraded during metabolism in the gastrointestinal tract in sheep and goats. Nevertheless it is unclear whether these studies can be applied to all or only to certain types of condensed tannins. It has to be noticed that not all condensed tannins have the same chemical structure and in this way, their chemical structure could interfere in their bioavailability. The main

sources of condensed tannins in animal studies are: grape seed extracts (rich in procyanidins), green tea leaves (prodelphinidin) chestnut, oak, mimosa and quebracho Colorado trees (rich in profisetinidin). The main difference between these kinds of proanthocyanidins is the presence or absence of – OH group in the carbon 5 (see table 2 and Figure 4). Profisetinidin compounds do not contain this group in carbon 5. It is known, that the absence of -OH group increase the stability of the interflavonol linkages in condensed tannin (Mueller-Harvey, 1999).

Fig. 4 Chemical structure of different proanthocyanidin compounds



Tannins are usually added in animal diets in rather low amounts, due to their antinutritional effects when present in high quantities.

When tannins are present in the diet in a proportion higher than 5% of feed dry matter, they have undesirable effects on animal digestion. For example condensed tannins can form complexes with proteins, metal ions, amino acids and polysaccharides, limiting the availability of these nutrients to animals (Goel *et al.*, 2005) and reducing the activity of ruminal microorganisms (Priolo *et al.*, 2000), resulting in an impairment of feed digestion in the rumen. To deactivate the negative effects of tannins it is possible to add the polymers PEG (polyethylene glycol) or PVP (polyvinyl pyrrolidone) in animals' diet: these polymers, in fact, deactivate effects of tannins because tannins have more affinity to these compounds than for other nutrients

(Makkar *et al.*, 1995b). In this way the bioavailability of nutrients to ruminants is higher. Furthermore, it is known that microbial population can adapt to tannin, protecting the animals from their antinutritional effects (Smith *et al.*, 2005).

Nevertheless, it is known that when tannins are present in the diet up to the 4-5% of feed dry matter, they can offer advantages for ruminants which result in increased milk production, wool growth, ovulation rate, and lambing percentage (Min *et al.*, 2003). Also tannins can protect proteins from degradation by ruminal microorganisms thus increasing the amount of proteins available for absorption in the intestine (McSweeney *et al.*, 2001).

There are lots of plants rich in condensed tannins like sulla, sainfoin, birdsfoot trefoil and others which are commonly used for ruminant feeding. Also there are several hardwood tree species rich in condensed tannins like chestnut, oak, mimosa and quebracho trees. Focusing the attention on quebracho trees, the plants commonly used for quebracho extracts production are *Schinopsis lorentzii* and *Schinopsis molle*, also named as quebracho Colorado.

Some positive effects of condensed tannins from quebracho on the quality of ruminants' meat are reported below:

Vasta *et al.*, (2009b) showed that when lambs are fed herbage supplemented with quebracho tannins, the level of  $\Delta^9$  desaturase protein expression in the muscle *longissimus dorsi* was significantly higher compared to lambs receiving the herbage without tannins. The function of the  $\Delta^9$  desaturase protein is to synthesize oleic acid, a monounsaturated  $\omega$ -9 fatty acid, by desaturating the stearic acid, a saturated fatty acid. This study could mean that tannins or their derivatives could act directly in the muscle's tissues.

In another study, Vasta *et al.*, (2009a) demonstrated *in vivo* that the supplementation of quebracho tannins to lambs given fresh herbage or a concentrate diet reduces ruminal biohydrogenation. During the biohydrogenation, the polyunsaturated fatty acids (PUFA) ingested through the diet are gradually hydrogenated by ruminal microorganisms to form stearic acid, a saturated fatty acid (Kepler and Tove 1967). An intermediate of this process is the fatty acid 9*cis* 11*trans* C18:2, known as conjugated linoleic acid (CLA) or rumenic acid. Vasta *et al.*, (2009a)

showed that when ruminants are fed with an enriched diet, the concentration of PUFA and CLA in meat is higher than in animals fed a control diet. The consumption of CLA it has been associated with a reduction of the incidence of cancer (Ip *et al.*, 1991), diabetes (Houseknect *et al.*, 1998), and atherosclerosis (Lee *et al.*, 2005).

It has been showed also that dietary tannins can affect ruminal biohydrogenation through changes in the ruminal microbial community. Vasta *et al.*, 2010 showed for the first time *in vivo* study that the conversion of vaccenic acid (unsaturated fatty acid) to stearic acid (saturated fatty acid) was reduced by tannin supplementation, probably because of a lower proportion of *Butyrivibrio proteoclasticus* bacteria, which are responsible for the last step of biohydrogenation.

Recent studies have shown the effect of condensed tannins on meat oxidative stability. It should be considered that oxidative reactions are the major causes of quality deterioration of raw meat and meat products. Oxidation of lipids and proteins in meat can affect its flavour (giving a rancid taste and smell), colour (giving discolouration, which result in an unpleasant brownish meat colour (Mancini and Hunt, 2005) and its nutritive value. Considering that condensed tannins, as all polyphenolic compounds, exert antioxidant activity, recent studies have investigated the effect of dietary tannins on meat oxidative stability. Luciano *et al.*, (2011b) reported an improvement of the antioxidant status of meat from lambs fed a concentrate-based diet supplement with tannins from a quebracho extract compared to a control tannin-free diet. The effects of phenolic compounds on meat oxidation had been studied previously by Moñino *et al.*, (2008) but in this case the effects of rosemary polyphenols were investigated. Moñino *et al.*, (2008) reported that the muscle of lambs receiving the milk of ewes fed with a rosemary-rich concentrate showed a greater antioxidant status than the control group. These authors reported that when lambs are fed with a polyphenol-enriched diet (tannins or flavonoids); lamb meat displayed a higher scavenging activity compared to the meat of lambs from the control group. Regarding discolouration process in meat, in another study Luciano *et al.*, (2009a) reported that including quebracho tannins in lamb diet improved meat colour stability by delaying myoglobin oxidation during refrigerated storage.

## **BIOAVAILABILITY OF PHENOLIC COMPOUNDS IN POLYGASTRICS**

---

The results obtained in the studies mentioned above imply that dietary quebracho tannins can affect the post-digestive metabolism in animals. Two possible mechanisms could explain these observations: i) ingested quebracho tannins (or their metabolites) might be degraded and absorbed from the ruminant digestive tract before being transferred to tissues or ii) dietary quebracho tannins (and their metabolites) are not absorbed in the digestive tract but act instead as antioxidants in the gastrointestinal tract.

Regarding to the bioavailability of polyphenolic compounds in ruminants, only few studies have showed the bioavailability of simple phenolic. Gladine *et al.*, (2007) found monomeric phenol compounds in plasma of sheep receiving polyphenol-rich plant extracts by ruminal infusion (i.e. grape or rosemary extracts, which contain galloylated condensed tannins or simple phenolics, respectively). Moñino *et al.*, (2008) reported that the muscle of lambs receiving the milk of ewes fed with a rosemary-rich concentrate contained several of the phenols that were present in the diet of the ewe (i.e. rosmarinic acid, carnosol and carnosic acid); while two other flavonoids, genkawanin and hesperetin, which are also present in rosemary extract, were not detected in lamb tissues. These studies suggested that only some phenolic compounds are bioavailable.

Little information is available about the presence of polyphenols in milk or dairy products. Lopez and Lindsay (1993) demonstrated the presence of phenolic compounds in bovine, caprine and ovine milk. It is known that the presence of these compounds takes a very important part in the sensory properties of milk and dairy products (for a review, see O'Connell *et al.*,2001).

## ***AIM OF THE EXPERIMENTS***

---

The aims of the following studies were to investigate whether different polyphenolic compounds or their metabolites could be detected in the tissues of lambs that had been fed previously with a specific diet. For the Experiment 1, lambs were fed with a diet supplemented with quebracho extract (characterised to be a rich source of proflisetinidin compounds, a group of condensed tannins). In the other experiment (Experiment 2), lambs were fed on pasture, specifically with *Lolium perenne* (this plant belongs to Poaceae family plant, which is not rich in polyphenolic compounds but contains some simple polyphenolic compounds). In both studies there was a control group, where lambs were fed with a commercial concentrate diet.

Furthermore, in both experiments the antioxidant status was measured in lamb tissue extracts that had either been treated or not treated with a Solid Phase Extraction (SPE) step. This step was used to purify samples and to isolate any phenolic compounds in the final extract. These experiments sought to investigate whether phenolic compounds have a direct or indirect antioxidant effect in lamb tissues.



## ***MATERIALS AND METHODS***

---

### **1. ANIMAL AND DIETS**

#### **EXPERIMENT 1**

Eighteen Comisana lambs were weaned at 45 days of age (mean weight 14.48 kg  $\pm$  standard deviation, SD 2.41kg). Lambs were blocked in groups of 2 on a descending body weight basis and, within block, were assigned to one of two dietary treatments of 9 animals each (C, control and C+T, control + tannins) and kept in individual pens for the duration of the trial. The C group received a concentrate containing the following ingredients: barley (55.1%), alfalfa hay (30.0%), soybean meal (13.0%), and vitamin and mineral premix (1.9%). The C+T group received the concentrate plus supplementary quebracho tannins (from *Schinopsis lorentzii*; Figli di Guido Lapi S.p.A., Castelfranco di Sotto, Pisa, Italy). For each 1000 g of DM of concentrate plus tannins, 95.7 g consisted of quebracho powder and the remaining 904.3 g of concentrate. The quebracho-supplemented diet was formulated to contain 6.4% (DM basis) tannins. The lambs were adapted to the experimental diets for 7 days before the commencement of the experiment. After 70 days of experiment the lambs were slaughtered,

#### **EXPERIMENT 2**

Sixteen Merinizzata Italian lambs were blocked in groups of 2 on a descending body weight basis after being weaned at 70 days of age. Lambs were assigned to one of two dietary treatments of 8 animals each (S, stall group and G, grazing group). The G group was allowed to graze from 9 am to 5 pm on a 1 ha ryegrass (*Lolium perenne*) at the end of the day the lambs were penned indoor in a multiple box and had *ad libitum* access to water. Before the commencement of the experiment, the animals were adapted to the experimental conditions over a 20 days period in which they were conducted to pasture and in stall received an amount of hay which was gradually

reduced till elimination from the diet. The S lambs were gradually adapted to the experimental feed over a 20 days period during which the weaning concentrate was gradually replaced with the experimental concentrate.

The experimental trial started when the animals were 90 days old and the experimental feeding trial had a duration of 72 days. All the animals were weighed weekly.

## **2. SAMPLING**

In both experiments individual blood samples were taken from the jugular vein of each lamb 12 hours after the last feeding trial and collected in heparin tubes. Blood samples were centrifuged at 3000 x g for 20 min at 4 °C and stored at -80 °C. Liver, taken at slaughtering, was immediately frozen in liquid nitrogen, vacuum packed and stored at -20 °C. Subsequently, a uniform and representative sample of each diet was prepared, vacuum packed and stored at -20 °C until analysis.

## **3. PREPARATION OF FEED SAMPLES AND PURIFICATION OF PHENOLIC COMPOUNDS BY SPE**

Feeds (2.5 g) given to the groups of each experiment (C and C+T for Experiment 1 and S and G for Experiment 2) were placed into 50 ml centrifuge tubes. Samples were homogenised with 15 ml acetone / water (70/30, v/v) for 60 seconds at 4000 rpm using a Heidolph Diax 900 tissue homogenizer (Heidolph Elektro GmbH & Co. KG, Kelheim, Germany). Samples were then sonicated for 6 min (with a break of 2 min after the first 3 min of sonication) using a Bandelin Sonoplus HD2070 sonicator (cycle: 4 × 10%, power: 0.31%). Samples were kept in a water/ice bath during both homogenization and sonication procedures. The sonicated homogenates were centrifuged at 3000 × g for 15 min at 4°C using a Thermo Scientific centrifuge (model IEC CL31R). Then the supernatants were filtered through Whatman 541 filter paper before SPE purification.

Phenolic compounds were isolated from feed samples on a reversed-phase C18 Sep-Pak Vac 6cc (500 mg) cartridge (WAT043395, WATERS SpA, Milan). The method was based on that described by Perez Magariño *et al.* (2008) but was adapted as follows. Prior to use, cartridges were conditioned with 3 ml methanol followed by 3 ml of distilled water. The filtered supernatant (10 ml) was acidified to pH 2.5 with 0.5 M-H<sub>2</sub>SO<sub>4</sub> prior to loading onto the cartridge. Phenolic compounds were eluted with 2 ml of methanol and The final fraction was divided into two 1.5 ml glass vials and kept in a freezer at -30°C.

#### **4. PREPARATION OF LAMB TISSUES SAMPLES AND PURIFICATION OF PHENOLIC COMPOUNDS BY SPE**

Liver (5 g) was placed into 50 ml centrifuge tubes. Preparation steps for liver samples were carried out in the same way as for feed samples. SPE method was based on that described by Perez Magariño *et al.* (2008) and carried out for feed samples. For liver samples phenolic compounds were eluted with 3 ml of ethyl acetate. The collected fraction was evaporated to dryness under nitrogen and then dissolved in 3 ml of methanol. The final fraction was also divided into two 1.5 ml glass vials and kept at -30°C.

The method for plasma samples was based on that described by Juan *et al.*, (2010) but was adapted as follows. A 500 µl plasma aliquot was acidified with 15 µl of glacial acetic acid. Phenolic compounds were isolated from plasma samples on a reversed-phase C18 Sep-Pak 1cc (100 mg) Cartridge (WAT023590 WATERS SpA, Milan). Prior to use, the cartridges were conditioned with 3 ml of methanol followed with 3 ml of distilled water. The acidified plasma sample was completely loaded onto the cartridge followed by 1 ml of distilled water. Phenolic compounds were eluted with 2 ml of methanol. Ascorbic acid (10 µl) was added to the final eluate to avoid any possible oxidation in plasma samples. This fraction was evaporated to dryness under nitrogen and then dissolved in 3 ml of methanol. The final fraction was divided into two 1.5 ml subsamples and stored at -30°C.

To confirm that the SPE method carried out in Experiment 1 and Experiment 2 did not reduce the initial concentration of polyphenolic compounds in a sample during the treatment, a recovery test was performed on 3 solutions of known concentration of gallic acid. Standard solutions were analysed for the concentration of total phenols using the Folin-Ciocalteu assay (Luciano *et al.*, 2011b). Subsequently, the same solution underwent the SPE treatment performed as explained above for plasma samples, but without the addition of ascorbic acid to the final eluate. The elute obtained after SPE step was evaporated to dryness under nitrogen and then dissolved in 1.5 ml of methanol: distilled water (1:1, v/v). The Folin-Ciocalteu assay was then performed on the SPE-treated extract. A 86.35% recovery was found, which shows that during the SPE treatment, exist only minimal loss (< 15%) of polyphenolic compounds in the samples.

## **5. FEED AND TISSUE EXTRACTS LC-MS ANALYSIS**

Feed and lamb tissue extracts were analysed by HPLC-MS using an ACE 5 2.1 × 150 mm C18 column (Hichrom Ltd, Theale, Berkshire, UK) fitted to an Agilent 1100 liquid chromatography with diode array detector. A binary mobile phase system was used where solvent A was HPLC-grade water + 0.1% formic acid and solvent B was HPLC S-grade acetonitrile + 0.1% formic acid (Rathburn Chemicals Ltd, Walkerburn, UK). Metabolites were eluted from the column using a simple gradient program. Initial conditions being 95% A and 5% B held for 1 min changing to 5% A and 95% B over 9 min and then held for 5 min before returning to the initial gradient conditions over 1 min and then held for 9 min to re-equilibrate the column. The pump flow rate was 0.2 ml/min and the column oven temperature was 25°C.

For LC-MS analysis, 5 µl of each sample was injected and the eluted peaks were analysed using an electrospray ionisation (ESI) micrOTOF QII quadrupole time of flight mass spectrometer (Bruker Daltonics, Coventry, UK) operated in the negative ion mode. Using a capillary voltage of 3200 V, nebuliser gas (N<sub>2</sub>) pressure of 1 bar, dry gas (N<sub>2</sub>) flow of 8 l/min and a drying temperature of 180°C. The TOF flight tube was set at +8600 V and the detector at 2010 V. The mass range, of 100 to 1700 daltons, was calibrated using Agilent low concentration Tunemix (G1969-85000). 5 µl of a

standard catechin solution (10 ng/ $\mu$ l) was first injected with each batch of samples to check system integrity and performance. Peak areas of lipophilic compounds were normalized based on catechin areas.

## **6. ANTIOXIDANT STATUS OF LIVER AND PLASMA SAMPLES TREATED WITH OR WITHOUT SPE**

Liver and plasma antioxidant status was determined by means of the FRAP (Ferric Reducing Antioxidant Power) and the Folin-Ciocalteu assays. Both assays were applied to samples either treated (SPE-samples) or not treated (RAW-samples) with SPE.

### **6.1. Folin-Ciocalteu assay in RAW-samples**

For the preparation of RAW-liver for the Folin-Ciocalteu assay, 2 g of liver from lambs of Experiment 1 were placed into 50 ml centrifuge tubes and homogenised with 10 ml of distilled water. For Experiment 2, liver (2 g) were placed into 50 ml centrifuge tubes and homogenised with 15 ml acetone / water (70/30, v/v). For the samples from the two experiments, homogenisation, sonication, centrifugation and filtration steps were performed as above described for SPE-liver. A 1:4 dilution of the extract (3 ml of distilled water added to 1 ml of liver extract) was chosen. The assay was performed as described by Luciano *et al.* (2011b). Briefly, 100  $\mu$ l of the diluted RAW-liver extract was transferred into 15 ml centrifuge tubes and 900  $\mu$ l of distilled water were added. The Folin-Ciocalteu reagent was diluted to 1 N and 500  $\mu$ l were added to the tubes followed by 2.5 ml aqueous solution of sodium carbonate (20% w/v). The mixture was vortex mixed for 30 s and incubated for 40 min in the dark at room temperature. The samples were centrifuged at  $2700 \times g$  for 10 min at 4°C in order to remove any sodium carbonate precipitates. A Shimadzu double-beam spectrophotometer (model UV-1601) was used to measure the absorbance of the samples. The wavelength used was 725 nm and a tube containing all the reagents except tissue extract was used as blank. Aqueous solutions of gallic acid were used for the calibration curve. The concentration range for the calibration curve covered 0  $\mu$ g to

80 µg/µl of gallic acid. The results were expressed as mg of gallic acid equivalents (GAE) / g of liver.

For RAW-plasma samples, 100 µl of plasma diluted 1:10 with distilled water were placed into 15 ml centrifuge tubes and 900 µl of distilled water were added. The Folin-Ciocalteu assay was carried out as described for RAW-liver samples. The results were expressed as mg of gallic acid equivalents (GAE) / ml of plasma.

### **6.2. Folin-Ciocalteu assay in SPE-samples**

Liver and plasma samples, from Experiment 1 and Experiment 2, treated with SPE were subjected to the Folin-Ciocalteu assay as follows. The content of one of the two 1.5 ml glass vials obtained after the SPE step was evaporated to dryness under nitrogen and then dissolved in 1.5 ml of methanol:distilled water (1:1, v/v). In the case of SPE-liver samples, 500 µl of this extract was transferred into a 15 ml centrifuge tube and 500 µl of methanol:distilled water (1:1, v/v) were added. The Folin-Ciocalteu assay was performed as described above for RAW-liver. Solutions of gallic acid in 1:1 (v/v) methanol:distilled water were used to calibrate the assay.

For measuring the total phenolic content and the antioxidant status in plasma, SPE-plasma samples were obtained as described above, with the only difference that ascorbic acid was not added to the final sample; in fact, Georgé *et al.* (2005) showed that ascorbic acid interferes in the Folin-Ciocalteu assay. The final SPE-plasma extracts were evaporated to dryness under nitrogen and then dissolved in 3 ml of methanol:distilled water (1:1, v/v); 500 µl of this sample was placed into a 15 ml centrifuge tube and 500 µl of methanol:distilled water (1:1, v/v) was added. The Folin-Ciocalteu assay was performed as described for RAW-liver. The results were expressed as mg of gallic acid equivalents (GAE) / ml of plasma.

### **6.3. Ferric reducing antioxidant power (FRAP assay) of RAW-samples**

The method described by Luciano *et al.* (2011b) was followed to measure the ferric reducing antioxidant power. The FRAP reagent was prepared by mixing 10

volumes of acetate buffer (300 mM, pH 3.6) with 1 volume of 10 mmol TPTZ solution (2,4,6-tripyridyl-s-triazine in 40 mM HCl) and with 1 volume of 20 mM aqueous ferric chloride. A blank reading at 593 nm was taken immediately after mixing 400  $\mu$ l of distilled water with 3.0 ml of FRAP reagent.

Experiment 1: For RAW-liver samples, 0.5 g of liver was placed into a 50 ml centrifuge tube and 10 ml of distilled water was added. Homogenisation, sonication, centrifugation and filtration steps were performed as above described for SPE-liver samples. Then, in a glass test tube, 300  $\mu$ l of distilled water were mixed with 100  $\mu$ l of liver extract and 3.0 ml of warm FRAP reagent (37°C) were added. The content of the tube was mixed and incubated in a water bath set at 37°C for 4 min, after which the absorbance was recorded at 593 nm.

Experiment 2: For RAW-liver samples, 2 g of liver was placed into a 50 ml centrifuge tube and 15 ml acetone / water (70/30, v/v) of distilled water was added. Homogenisation, sonication, centrifugation and filtration steps were performed as above described for SPE-liver. 74  $\mu$ l of liver extract, 220  $\mu$ l of distilled water and 3.0 ml of warm FRAP reagent (37°C) were added in a glass test tube, 300 were mixed with 100. The content of the tube was mixed and incubated in a water bath set at 37°C for 4 min, after which the absorbance was recorded at 593 nm.

Then for both experiments: The change in absorbance ( $\Delta A_{593 \text{ nm}}$ ) between the final reading and the blank reading was related to that obtained with solutions of  $\text{Fe}^{2+}$  of known concentrations (aqueous  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$  ranging from 0  $\mu\text{M}$  to 1000  $\mu\text{M}$ ). Results of the FRAP assay were therefore expressed as  $\mu\text{moles of Fe}^{+2}$  equivalents /g of liver.

For RAW-plasma sample in both experiments, 74  $\mu$ l of plasma was added in a glass test tube to 220  $\mu$ l of distilled water and 2.2 ml of FRAP reagent (the final dilution of the sample in the mixture was always 1:34). Tubes were mixed and incubated for 4 min in a water bath at 37°C. The absorbance was immediately recorded at 593 nm. Results of the FRAP assay were therefore expressed as  $\mu\text{moles of Fe}^{+2}$  equivalents /ml of plasma.

#### **6.4. Ferric reducing antioxidant power (FRAP assay) of SPE-samples**

The same method described above was applied to the SPE-samples. Since SPE-samples were prepared in a methanol:distilled water solution, the same solution was used instead of distilled water in the assay, as well as for preparing standard  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$  solutions. The assay was performed by adding in a glass test tube 200  $\mu\text{l}$  of the SPE-liver samples or SPE-plasma samples (without ascorbic acid) to 200  $\mu\text{l}$  of methanol:distilled water and 3 ml of FRAP reagent. Incubation time, absorbance measurement and calculations were performed as described above.

#### **7. STATISTICAL ANALYSIS**

For each experiment, the data from the Folin-Ciocalteu assay and FRAP values were analysed by one-way ANOVA including the model treatment effects (C vs C+T or S vs G) and experimental error. Each lamb was considered as an individual experimental unit.



## RESULTS

---

### 1. ANALYSIS OF FEED SAMPLES BY LC-MS

#### EXPERIMENT 1

Representative chromatograms of quebracho enriched and control diets are shown in Fig. 5. The quebracho chromatogram differed from the control by the presence of one peak at 12.1 min. The extracted ion chromatogram revealed that this peak contained several different masses (Fig. 6) with  $m/z$  (H<sup>+</sup>) of 561.150, 833.223 and 1105.293. These compounds were identified as a combination of one or more fisetinidol units plus one catechin unit (Fig. 8; Table 4). Another single peak was observed at 12.8 min with an  $m/z$  (H<sup>+</sup>) of 285.046 and was identified as fisetin (Fig. 5 and 7; Table 4). LC-MS analysis of the control diet revealed the absence of the compounds identified in the quebracho enriched diet.

Table 4. Main phenolic compounds identified in the quebracho-enriched diet by LC-MS

Compound	$m/z$ (H <sup>+</sup> )	Formula	Assignments*
1	285.046	C <sub>15</sub> H <sub>10</sub> O <sub>6</sub>	Fisetin
2	561.150	C <sub>30</sub> H <sub>26</sub> O <sub>11</sub>	One fisetinidol unit plus one catechin unit
3	833.223	C <sub>45</sub> H <sub>38</sub> O <sub>16</sub>	Two fisetinidol units plus one catechin unit
4	1105.293	C <sub>60</sub> H <sub>50</sub> O <sub>21</sub>	Three fisetinidol units plus one catechin unit

\* See Fig. 8 for chemical structures.

Fig. 5. LC chromatograms (wavelength: 214 nm) from quebracho-supplemented diet (a) and control diet (b)




Fig. 6. LC-chromatogram (a) and extracted ion chromatograms for  $m/z$   $H^+$  = 561.15 (b), 833.20 (c) and 1105.29 (d) from quebracho-enriched diet (wavelength: 214 nm).




Fig. 7. LC-chromatogram (a) and extracted ion chromatogram for  $m/z$   $H^+$  = 285.046 (b) from quebracho enriched diet (wavelength: 214nm)







Fig. 8: Possible assignments of profisetinidin ions that were detected in the LC-MS chromatograms of extracts from the quebracho-enriched diet




Chemical Formula :  $C_{15}H_{10}O_6$   
Exact Mass: 285.046 (fisetin)



Chemical Formula :  $C_{30}H_{26}O_{11}$   
Exact Mass: 562.150 (profisetinidin dimer)



Chemical Formula :  $C_{45}H_{38}O_{16}$   
Exact Mass: 834.223 (profisetinidin trimer)



$n = 2$  Chemical Formula :  $C_{60}H_{50}O_{21}$   
Exact Mass: 1106.293


## EXPERIMENT 2

Representative chromatograms of grass and concentrate diets are shown in Fig. 9. Under the LC-MS conditions used in the present study, phenolic compounds were expected to elute between 5 and 13min. In grass chromatogram, only few peaks and with a very low intensity were observed in the range of phenolic compounds (Fig. 9). The phenolic compounds identified in the chromatogram of grass samples belong to the flavonol family and most of the compounds are present in their glycoside form (Table 5).

Table 5. Identification of possible phenolic compounds found in grass diet by LC-MS

RT (min)	<i>m/z</i> (H <sup>-</sup> )	Formula	Phenol compound
10.8	755.2110	C <sub>33</sub> H <sub>45</sub> O <sub>20</sub>	Kaempferol-3-O-glucosyl-rhammosyl-galactoside
11.2	563.1438	C <sub>26</sub> H <sub>28</sub> O <sub>14</sub>	Kaempferol-xylosyl-rhamnoside
11.7	609.1506	C <sub>27</sub> H <sub>30</sub> O <sub>16</sub>	Rutin
12.0	593.1549	C <sub>27</sub> H <sub>30</sub> O <sub>15</sub>	Kaempferol-3-O-rutinoside
12.5	489.1054	C <sub>23</sub> H <sub>22</sub> O <sub>12</sub>	Kaempferol-3-O-acetyl-glucoside
12.7	285.0430	C <sub>15</sub> H <sub>10</sub> O <sub>6</sub>	Kaempferol (tetrahydroxyflavone)

Fig. 9. LC-chromatograms (wavelength: 214 nm) from grass diet (a) and control diet (b). The first chromatogram is a complete chromatogram of the diets; the chromatogram below is an enlargement of the corresponding range of phenolic compounds.




Some of the peaks observed between 10 and 22 min in the chromatograms from grass and control diet are due to lipophilic compounds (Figure 9). Among the lipophilic compounds identified in the diets, the compound observed at 18.2 min with a mass  $m/z$  ( $H^+$ ) of 277.2206 (probably, linolenic acid) was found in a proportion of 7.5 times higher in the grass diet than in the concentrate diet. The peak detected at 20 min with a mass  $m/z$  ( $H^+$ ) of 279.2358 (plausibly linoleic acid) which was 4.5 times higher in the concentrate diet than in the grass diet.

## 2. ANALYSIS OF ANIMAL TISSUES SAMPLES BY LC-MS

### EXPERIMENT 1

Plasma and liver extracts obtained after the SPE step were analysed for quebracho profisetinidins and other phenolic compounds or possible metabolites arising from their degradation. However, no signals from any of these compounds could be detected in liver (Fig. 10) or plasma (Fig. 11) samples from lambs fed the C+T diet. Fig. 10 shows that liver chromatograms from lambs fed the C or C+T diet are exactly the same.

Fig 10. LC-chromatogram (wavelength: 214 nm) of a liver sample from one lamb from the C+T group (a) and of a liver sample from one lamb from the C group (b).



In plasma chromatograms (Fig. 11) no differences could be found between the lambs fed with the different diets. In plasma chromatograms the peak at 2.5 min is ascorbic acid, which had been added during the SPE step. The peaks observed between 12 and 25 min in the chromatograms from plasma and liver samples are due to lipophilic compounds (Table 6); the plasma of the C+T lambs presented greater ( $P \leq 0.011$ ) amounts of  $C_{18}H_{30}O_2$ ,  $C_{18}H_{32}O_2$ ,  $C_{18}H_{34}O_2$  and  $C_{20}H_{30}O_2$  compared to the C animals. No significant differences ( $P > 0.05$ ) were found in the amounts of the lipophilic compounds detected in liver samples of the lambs from the two groups.

Fig 11. LC-chromatogram (wavelength: 214 nm) of a plasma sample from one lamb from the C+T group (a) and of a plasma sample from one lamb from the C group (b).






Table 6. Lipophilic compounds\* found in liver and plasma of lambs fed concentrate (C) or concentrate plus quebracho tannins (C+T).

<i>m/z</i> (H <sup>-</sup> )	<i>Formula</i>	<i>Plasma</i>							
		<i>Liver</i>			<i>Plasma</i>				
		DIET		<i>P value</i>	DIET		<i>P value</i>		
C	C+T	C	C+T		SEM				
255·235	C <sub>16</sub> H <sub>32</sub> O <sub>2</sub>	807,930	2,110,674	0·090	384,486	108,944	154,186	0·200	547,031
265·151	C <sub>15</sub> H <sub>22</sub> O <sub>4</sub>	16,472,511	24,140,824	0·080	2,195,887	2,594,968	3,475,423	0·174	318,125
277·221	C <sub>18</sub> H <sub>30</sub> O <sub>2</sub>	9,246,287	11,522,243	0·222	909,524	39,008	1,811,618	0·006	28,156
279·236	C <sub>18</sub> H <sub>32</sub> O <sub>2</sub>	15,870,152	17,634,140	0·153	608,223	67,688	315,776	0·004	47,583
281·251	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	17,133,219	20,462,115	0·702	4,146,446	312,905	936,492	0·011	130,618
299·263	C <sub>18</sub> H <sub>36</sub> O <sub>3</sub>	1,675,988	1,962,691	0·402	164,443	-	-	-	-
301·221	C <sub>20</sub> H <sub>30</sub> O <sub>2</sub>	1,389,920	13,174,605	0·914	504,255	364,171	478,965	0·008	23,333

\* Data expressed as peak area

## EXPERIMENT 2

Liver and plasma extracts obtained after the SPE step were analysed for the same phenolic compounds found in grass diet or possible metabolites arising from their degradation. However, no signals from any phenolic compounds could be detected in liver (Fig. 12) or plasma (Fig. 13) samples from the G lambs.

Fig. 12. LC-chromatogram (wavelength: 214 nm), from 4 to 16 min (range of phenolic compounds) of a liver sample from one lamb from the G group (a) and of a liver sample from one lamb from the S group (b).





Fig. 13. LC-chromatogram (wavelength: 214 nm), from 4 to 16 min (range of phenolic compounds) of a plasma sample from one lamb from the G group (a) and of a plasma sample from one lamb from the S group (b).



In this experiment, also some peaks observed at the end of the chromatograms were due to lipophilic compounds. In liver chromatograms some plausible fatty acids were identified (Fig. 14). In the case of plasma chromatograms, Extracted Ions Chromatograms with the masses of the compounds identified previously in liver chromatograms were necessary to reveal the presence of these compounds and for their quantification.

The identification of the possible fatty acids and their concentration in liver and plasma can be observed in Table 7. Liver samples from the G lambs showed greater amount ( $P < 0.001$ ) for the compounds identified as  $C_{16}H_{32}O_2$ ,  $C_{18}H_{30}O_2$  and  $C_{20}H_{30}O_2$  compared to liver from the S lambs.

No significant differences were found for the same compounds in plasma samples. In liver and plasma samples from lambs fed the concentrate, higher concentrations ( $P < 0.05$ ) were found for the compounds identified as  $C_{18}H_{32}O_2$ , and  $C_{18}H_{36}O_3$ , compared to liver and plasma from lambs fed the grass diet.

Fig. 14. LC-chromatogram (wavelength: 214 nm), from 15 to 22 min (range of lipophilic compounds) of a liver sample from one lamb from the G group (a) and of a liver sample from one lamb from the S group (b). For identification of lipophilic compounds see Table 7.




Table 7. Lipophilic compounds\* found in liver and plasma of lambs fed in stall (S) or grazing (G).

Compound	<i>m/z</i> (H <sup>+</sup> )	Formula	Liver						Plasma					
			DIET			DIET			DIET			DIET		
			S	G	<i>P value</i>	SEM	S	G	<i>P value</i>	SEM	S	G	<i>P value</i>	SEM
1	281·251	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	2,916,083	3,877,909	0·213	376,738	427,320	179,944	0·004	47,583				
2	255·235	C <sub>16</sub> H <sub>32</sub> O <sub>2</sub>	971,947	1,794,396	<0·001	132,619	174,716	135,092	0·200	547,031				
3	277·221	C <sub>18</sub> H <sub>30</sub> O <sub>2</sub>	1,237,630	4,099,981	<0·001	407,898	32,717	53,346	0·239	8,535				
4	279·236	C <sub>18</sub> H <sub>32</sub> O <sub>2</sub>	7,112,070	4,564,985	0·001	428,736	310,078	56,700	0·009	51,957				
5	299·263	C <sub>18</sub> H <sub>36</sub> O <sub>3</sub>	809,313	56,284	0·001	126,743	143,136	16,149	0·025	29,491				
6	301·221	C <sub>20</sub> H <sub>30</sub> O <sub>2</sub>	2,007,747	6,663,365	<0·001	639,850	10,366	19,515	0·226	3,683				

\* Data expressed as peak area

### 3. FOLIN-CIOCALTEU ASSAY IN LAMB TISSUES SAMPLES

#### EXPERIMENT 1

The Folin-Ciocalteu assay, performed on the RAW samples, gave significantly higher values in both liver and plasma samples from lambs fed the quebracho-enriched diet as compared to tissues from animals fed the control diet (+9.87%,  $P = 0.036$  for liver samples and +5.91%,  $P = 0.006$  for plasma samples; Table 8).

However, when the liver and the plasma samples had been treated with SPE, the Folin-Ciocalteu assay did not show any differences ( $P > 0.05$ ) between the experimental treatments (Table 8). It is also noteworthy that the Folin-Ciocalteu values of the SPE-samples were 100-fold lower compared to the RAW-samples.

#### EXPERIMENT 2

The Folin-Ciocalteu assay, on Experiment 2, gave significantly higher values in liver and plasma samples from lambs fed the grass diet as compared to tissues from animals fed the concentrate diet (+13.15%,  $P = 0.055$  for liver samples and +4.27%,  $P = 0.093$  for plasma samples; Table 8). At the same way, when the liver samples were treated with SPE, the Folin-Ciocalteu assay did not show any differences ( $P > 0.05$ ). The treatment with SPE resulted in a 100-fold reduction of the Folin-Ciocalteu values compared to the RAW-samples (Table 8). Moreover, after SPE purification, plasma samples from S lambs showed higher Folin-Ciocalteu values compared to G lambs (+39.23%,  $P = 0.012$ ).

#### **4. FERRIC REDUCING ANTIOXIDANT POWER (FRAP) ASSAY IN LAMB**

##### **TISSUES SAMPLES**

##### **EXPERIMENT 1**

The FRAP values were also significantly higher (+9.6%,  $P = 0.038$  for liver samples and +28.45%,  $P < 0.0005$  for plasma samples; Table 9) in RAW liver and plasma samples from lambs fed the C + T diet than tissues from lambs fed the C diet. However, for SPE liver and plasma samples, the FRAP values were similar ( $P > 0.05$ ) in all tissues from lambs fed the two diets (Table 9).

##### **EXPERIMENT 2**

The RAW-liver and RAW-plasma from lambs fed the grass diet showed significantly higher FRAP values compared to liver and plasma from lambs fed the concentrate diet (+20.03%,  $P = 0.001$  for liver samples and +14.28%,  $P = 0.003$  for plasma samples, Table 9). However, when the liver and plasma samples were treated with the SPE, differences in FRAP values were not detected ( $P > 0.05$ ; Table 9).

Table 8. Effect of dietary treatments on Folin Ciocalteu values of RAW-samples and SPE-samples in Experiment 1 and Experiment 2

	EXPERIMENT 1				EXPERIMENT 2			
	DIET		<i>P value</i>	SEM	DIET		<i>P value</i>	SEM
	C	C+T			S	G		
RAW-LIVER*	4.071	4.517	0.036	0.1090	1.409	1.6224	0.055	0.056
RAW-PLASMA†	1.926	2.047	0.006	0.0226	1.614	1.686	0.093	0.021
SPE-LIVER*	0.0104	0.0095	0.390	0.0005	0.0121	0.0123	0.932	0.00098
SPE-PLASMA†	0.0163	0.0186	0.180	0.0017	0.0785	0.0477	0.012	0.00649

\* Expressed as mg of gallic acid equivalents (GAE) / g of liver  
† Expressed as mg of gallic acid equivalents (GAE) / ml of plasma

Table 9. Effect of dietary treatments on FRAP values of RAW-samples and SPE-samples in Experiment 1 and Experiment 2

	EXPERIMENT 1				EXPERIMENT 2			
	DIET		<i>P value</i>	SEM	DIET		<i>P value</i>	SEM
	C	C+T			S	G		
RAW-LIVER‡	18.730	20.720	0.038	0.7050	5.309	6.639	0.001	0.226
RAW-PLASMA§	0.410	0.573	<0.0005	0.0255	0.768	0.896	0.003	0.021
SPE-LIVER‡	0.0710	0.0740	0.790	0.0043	0.0622	0.0654	0.694	0.00385
SPE-PLASMA§	0.1620	0.1600	0.948	0.0170	0.1191	0.1195	0.985	0.01070

‡ Expressed as  $\mu\text{moles of Fe}^{+2}$  equivalents /g of liver  
§ Expressed as  $\mu\text{moles of Fe}^{+2}$  equivalents /ml of plasma



## *DISCUSSIONS*

---

### **EXPERIMENT 1**

To the best of our knowledge, this is the first study aiming to determine the metabolic fate of quebracho tannins after being ingested by ruminants. It is well known that tannins complex proteins and interfere with ruminal digestion in general. Protein complexation affects but does not eliminate the antioxidant activities of tannins (Riedl *et al.*, 2001; Arts *et al.*, 2002). Therefore, it is possible that tannins may indirectly affect muscle biochemistry via some other components. In the light of these speculations, the present study investigated whether or not dietary quebracho tannins or their metabolites were present in lamb plasma and liver.

The main types of tannins found in the quebracho-enriched diet were profisetinidins (Table 1) and have been described previously in quebracho extracts (Roux, 1992; Mueller-Harvey, 2006). It should be noted that, in contrast to most other condensed tannins, profisetinidins do not contain 5-OH groups in close proximity to the interflavanol bond. The absence of this 5-OH group increases the stability of the interflavanol linkages in condensed tannins (Mueller-Harvey, 1999), and therefore quebracho tannins are particularly difficult to degrade.

It has been explained previously that the bioavailability of phenolic compounds in ruminants has been only marginally investigated. Gladine *et al.* (2007) found monomeric phenol compounds in plasma when sheep had received polyphenol-rich plant extracts by ruminal infusion (i.e. grape or rosemary extracts, which contain galloylated condensed tannins or simple phenolics, respectively). Moñino *et al.* (2008) reported that the muscle of lambs receiving the milk of ewes fed with a rosemary-rich concentrate contained several of the phenols that were present in the diet of the ewe (i.e. rosmarinic acid, carnosol and carnosic acid); while two other flavonoids, genkawanin and hesperetin, which are also present in rosemary extract, were not detected in lamb tissues. These studies suggested that only some phenolic compounds

are bioavailable. Clearly, the chemical structures of proanthocyanidins and phenolic acids studied by Gladine *et al.*, (2007) and Moñino *et al.*, (2008) differ from the profisetinidins investigated in the present study (Fig. 3).

This study demonstrated that no phenolic compounds could be detected in lamb tissues, which indicates that the profisetinidin tannins from quebracho are not degraded or absorbed in the rumen or in the digestive tract. This agrees with Makkar *et al.*, (1995a) who reported that quebracho tannins, as measured by the butanol-HCl-Fe<sup>3+</sup> reagent (Porter *et al.*, 1986), are not degraded by ruminal microorganisms in an *in vitro* study. Therefore, it is likely that these types of tannins are directly eliminated through the faeces. Nevertheless, other authors reported that ruminal microflora were able to degrade quebracho tannins into smaller phenolics (Bhat *et al.*, 1998).

In the present study the tissue extracts which had been passed through SPE cartridges from lambs fed with the control diet or with the quebracho-enriched diet gave a similar response to the Folin-Ciocalteu and FRAP assays. However, the RAW liver and plasma of the C+T lambs displayed higher FRAP and Folin-Ciocalteu values than the samples from the control lambs. While the SPE method used is highly selective for the isolation and concentration of phenolic compounds (Pérez-Magarino *et al.*, 2008), the Folin-Ciocalteu reagent and the FRAP assay are not specific to phenolic compounds and react to a wide spectrum of reducing compounds (Georgè *et al.*, 2005). Therefore, it is likely that the increased antioxidant capacity of the RAW liver and plasma of the quebracho-fed lambs is due to the presence of reducing compounds other than phenolics and that these were removed by the SPE purification step. This would be in agreement with the results from the LC-MS analysis, which showed that the SPE-treated tissues of the C + T lambs did not possess any phenolic compounds. The results of the current study are consistent with the report by Luciano *et al.*, (2011b) who showed that tannins from quebracho increased the antioxidant status in lamb muscle and the total phenolic content, as measured by the Folin-Ciocalteu reagent. However, it is not possible to make a direct comparison between the results reported by Luciano *et al.*, (2011b) and the present study, as in the former study both FRAP and Folin-Ciocalteu assays were performed only in muscle *longissimus dorsi*.

Although the mechanism of the antioxidant effect of dietary quebracho tannins could not be deduced from the present study, it can be assumed that the greater antioxidant status of the RAW-tissues from the C+T lambs is likely due to an indirect antioxidant effect of dietary tannins. This effect could be mediated by a direct antioxidant activity of the tannins in the gastrointestinal tract, such as a removal or chelation of pro-oxidant compounds and a reduction of lipid peroxidation, which would result in an overall improvement of the animal's antioxidant status (Kerem *et al.*, 2006, Halliwell *et al.*, 2005). Furthermore, it is known that dietary condensed tannins strongly modify lipid metabolism in ruminants (Vasta *et al.*, 2009a) and interfere also with gene (Kresty *et al.*, 2011) and protein expression (Vasta *et al.*, 2009b). In particular, Sgorlon *et al.* (2006) found that supplementing sheep with grape skin extract, which is rich in polyphenols and condensed tannins, increased the expression in plasma of the superoxide dismutase enzyme, which is involved in the endogenous antioxidant defence system.

The lipophilic compounds detected in liver and plasma are likely to be fatty acids and, in particular, the formulae  $C_{18}H_{30}O_2$ ,  $C_{18}H_{32}O_2$  and  $C_{20}H_{30}O_2$  could correspond to some  $C_{18}$  and  $C_{20}$  poly-unsaturated fatty acids (PUFA) and  $C_{18}H_{34}O_2$  could correspond to a mono-unsaturated fatty acid. The plasma of the C+T lambs contained more of these compounds than the plasma of C lambs (Table 2). It is known that feeding tannins results in an increased PUFA accumulation in tissues; this is due to a reduced biohydrogenation of PUFA in the rumen, as tannins depress ruminal microorganism activity and proliferation (Vasta *et al.*, 2009a).

It can be concluded that when lambs are fed with a quebracho supplemented diet, no tannins or other phenolic compounds were found in their liver or plasma. This is in contrast with studies in which plant extracts rich in other types of polyphenols were given to lambs. The present results may be due both to the very low amount of fisetin (a low MW phenol) and to the structural stability of profisetinidins in quebracho. Although no phenolic compounds were detected in the liver and plasma of quebracho-fed lambs, these tissues showed a greater antioxidant capacity compared to liver and plasma of unsupplemented lambs. Therefore, it can be concluded that supplementing quebracho tannins improves the antioxidant capacity of tissues via an indirect effect,

possibly by enhancing the tissue endogenous antioxidant system or by participating in the regeneration of other antioxidant compounds.

## **EXPERIMENT 2**

*Lolium perenne*, commonly named as Perinneal Ryegrass, is a member of the Poaceae (Gramineae) family and it is one of the most important pasture grass species in animals feeding around the world (Cai *et al.*, 2011). Like all plants, *Lolium perenne*, contains polyphenolic compounds. Information about the specific compounds present in this plant is scarce, however, it is known that plants belonging to the family Poaceae, are not as rich in polyphenolic compounds such as plants belonging to the family Fabaceae (Leguminosae) (Reynaud *et al.*, 2010).

In this experiment the grass diet sample was analysed by LC-MS after a SPE treatment to identify the possible phenolic compounds present in the sample. The chromatogram showed a few peaks and with a very low intensity compared to peaks observed at the end of the chromatogram, which belong to lipophilic compounds.

Most of the peaks present in the range of phenolic compounds were identified as different kaempferol glycosides (Table 5) with the masses  $m/z$  (H-) of: 755·211 ( $C_{33}H_{45}O_{20}$ ), 563·1438 ( $C_{26}H_{28}O_{14}$ ), 593·1549 ( $C_{27}H_{30}O_{15}$ ) and 489·1054 ( $C_{23}H_{22}O_{12}$ ). The compound with the mass  $m/z$  (H-): 285·043 could belong to kaempferol compound in its aglycone form. The other compound with  $m/z$  (H-): 609·1506 could be the flavonoid rutin, which is a glycoside of the flavonoid quercetin. All these compounds are part of the group of flavonoids, belonging specifically to the subclass of flavonols. It is well known that the flavonoid kaempferol and its glycosides are very common and they occur in many plants. Chopin and Dellamonica (1988) reported the synthesis and identification of glycosylflavonoids in numerous plant species. In a more recent and specific study, Tu *et al.*, (2010) identified several flavonols compounds in *Lolium perenne* samples. Some of the compounds identified in this study are the same compounds identified previously by Tu *et al.* (2010) i.e. the compounds with a  $m/z$  (H) of: 593·1549, 609·1506 and 755·211. The other compounds identified in the present study were not identified by Tu *et al.*, (2010). As well as Tu *et al.*, (2010) identified other phenolic compounds that in the present study have not been revealed in the chromatogram.

Regarding the chromatograms of the animal tissues, no peaks in the phenolic compounds range were detected in liver or plasma from lambs fed the grass diet. These results could be explained at least in three different ways:

i.- Phenolic compounds from the grass diet are not absorbed in the digestive tract of lambs, and they are directly excreted in the urine. The bioavailability of polyphenols in ruminants is yet unknown, but this supposition could agree with Scalbert *et al.*, 2002, who confirmed that in monogastric animals, most dietary polyphenols are quickly eliminated in both urine and bile after ingestion.

ii.- Flavonoids from grass were absorbed in lambs but were already excreted when sampling was carried out. In the present study blood and liver sampling was taken at least 12 hours after the last grass feeding. There is not information about polyphenols turnover in polygastric, but this hypothesis could be confirmed by Scalbert *et al.*, (2002) that demonstrated that in human plasma, a post-prandial peak is observed 1–2 h after ingestion of various flavonols and flavanols.

iii.- The concentration of phenolic compounds in *Lolium perenne* is not high enough for their transfer into animal tissues. The results obtained in the present study cannot be compared with the results obtained by other authors, which confirm the transfer of phenolic compounds from the diet to animal tissues. For example, Gladine *et al.*, (2007) administered a single acute dose of rosemary, grape, citrus or marigold directly in the rumen of sheep. After the administration, epicatechin was found in plasma from animals dosed with grape, while naringenin was detected in plasma from lambs dosed with citrus. However, some phenolic compounds commonly found in rosemary or marigold were not detected in the plasma of sheep. In another study (Moñino *et al.*, 2008) lambs were fed with distilled rosemary leaves and phenolic compounds were detected in lamb muscles. However, it has to be considered that among the phenolic compounds present in the rosemary leaves, only three phenols were found in meat lamb (rosmarinic acid, carnosol and carnosic acid). Furthermore, rosemary is a rich source of phenolic compounds (Zheng *et al.*, 2001), while *Lolium perenne* is not.

Regarding the results obtained on the antioxidant status of lamb tissues, FRAP values showed significant difference ( $P < 0.005$ ) in both RAW-samples from lambs fed the grass diet compared to lambs the concentrate diet. These results are similar to the results obtained by Descalzo *et al.*, (2007). That demonstrated that fresh meat from cattle raised at pasture had higher FRAP values than meat from grain-fed animals Folin-Ciocalteu assay showed a higher antioxidant potential of RAW-liver from lambs fed the grass diet compare to animals fed the concentrate diet, while a similar trend was found in tendency in RAW-plasma samples. However, SPE-liver and SPE-plasma did not show differences in the Folin-Ciocalteu and in the FRAP values between lambs fed the grass diet and the lambs from stall group. It should be appreciated that Folin-Ciocalteu assay gave significant difference in RAW-plasma from S lambs compared to G lambs. This result was unexpected and the reason for this is unclear.

As speculated in Experiment 1, it is likely that the increased antioxidant capacity of the RAW liver and plasma grazing lambs is due to the presence of reducing compounds other than phenolics and that these were removed by the SPE purification step. This hypothesis is supported by the fact that both the Folin-Ciocalteu and the FRAP assays are not specific for phenolic compounds; rather several reducing compounds can account for the results of both tests (Georgè *et al.*, 2005). In the same way, this would be in agreement with the results from the LC-MS analysis, which showed that after a SPE treatment, tissues from lambs fed the grass diet did not possess any phenolic compounds.

The peaks observed at the end of the chromatograms of the grass and concentrate diets, as well as of the liver and plasma samples are lipophilic compounds, specifically are likely to be fatty acids. It is known that a grass diet is characterised to contain a high quantity of linolenic fatty acid, precursor of the  $n - 3$  series of fatty acids, and, on the other hand, a concentrate diet is characterised to have a high level of linoleic acid, precursor of the  $n - 6$  fatty acid series (Diaz *et al.*, 2002). The results obtained in the present study agree with previous results (Diaz *et al.*, 2002).

Liver and plasma samples from lambs fed the concentrate diet compared to liver from lambs fed the grass, has showed higher concentrations of the compounds identified as  $C_{18}H_{32}O_2$ , and  $C_{18}H_{36}O_3$ , which can be probably identified as linoleic acid and hydroxy stearic acid respectively. Conversely, in liver from grass-lambs the compounds  $C_{16}H_{32}O_2$ ,  $C_{18}H_{30}O_2$  and  $C_{20}H_{30}O_2$  - which could be identified respectively as palmitic acid, linolenic acid and eicosapentaenoic acid (EPA) – were higher compared to the concentrate-based diet. On the other hand, plasma samples did not show significant differences for these compounds.

In data not published yet, fatty acids have been identified and quantified in the muscle *longissimus dorsi* from the same animals of the present study. In muscle from G-lambs group compared to S-lambs group was found higher concentrations of C18:3w3, linolenic acid, (2.77% vs 0.32% of total fatty acids  $P < 0.001$ ) and of C20:5w3, EPA, (1.66% vs 0.30%  $P < 0.001$ ). Also significant difference ( $P < 0.001$ ) and greater concentrations for C18:2c9c12 (linoleic acid) was found for S-lambs group compared to G-lambs group (16.58% vs 8.68% of total fatty acids). These results and the results obtained in the present study agree with the results obtained by Nuernberg *et al.*, (2008) and Scerra *et al.*, (2011) on the fatty acid composition and concentration of the muscle *longissimus dorsi* of lambs fed a grass or a concentrate diet.

It can be concluded that *Lolium perenne* is not a rich source of phenolic compounds, which could explain why, in liver and plasma from lambs fed at pasture with this plant, no phenolic compounds found previously in grass or their metabolites were found. The antioxidant status of liver and plasma from grass-fed lambs was higher compared to tissues from animals fed concentrates and this agrees with previous reports showing a positive effect of grass feeding in improving the antioxidant status of animal tissues compared to concentrate-based diets. However, when samples were purified by SPE in order to isolate phenolic compounds, differences in the antioxidant status of tissues from animals fed grass or concentrates disappeared.

Therefore, in this experiment, it has been shown for the first time, that the higher antioxidant status of tissues from ruminants fed a grass-bases diet, cannot be attributed to a direct transfer of phenolic compounds from plant to tissues.



## *CONCLUSION*

---

These studies have been the first studies to study the bioavailability of profisetinidins compounds from quebracho (Experiment 1) and the bioavailability of flavonoids present naturally on pasture (Experiment 2) on animal tissues.

In both experiments, no signals of the compounds present in feed samples or their corresponding metabolites were found in liver or plasma samples from lambs fed the respective diets. But with the existing knowledge in literature about the antioxidant capacity of polyphenolic compounds, in these studies can be supposed that although polyphenols do not reach the tissues, they improve the antioxidant status of animal tissues via an indirect effect.

Further research is required to study the bioavailability of polyphenols compounds in ruminants and to understand the possible mechanism of action of polyphenolic compounds in improving the antioxidant status in ruminants.

## REFERENCES

---

- Abad-García B., Berrueta LA., López-Márquez DM., Crespo-Ferrer I., Gallo B., Vicente F. 2007. Optimization and validation of a methodology based on solvent extraction and liquid chromatography for the simultaneous determination of several polyphenolic families in fruit juices. *Journal of Chromatography A* 1154, 87-96.
- Aldini G., Carini M., Piccoli A., Rossoni G., Facino RM. 2003. Procyanidins from grape seeds protect endothelial cells from peroxynitrite damage and enhance endothelium-dependent realization in human artery: new evidences for cardio-protection. *Life Sciences* 73, 2883-2898.
- Arts MJTJ., Haenen GRMM., Wilms LC., Beetstra SAJN., Heijnen CGM., Voss HP., Bast A. 2002. Interactions between flavonoids and proteins: effect on the total antioxidant capacity. *Journal of Agriculture and Food Chemistry* 50, 1184-1187.
- Baba S., Osakab N., Natsume M., Terao J. 2002. Absorption and urinary excretion of procyanidin B2 [epicatechin-(4 $\beta$ -8)-epicatechin] in rats. *Free Radical Biology and Medicine* 33, 142-148.
- Balasundram N., Sundram K., Samman S. 2006. Phenolic compounds in plants and agri-industrial by products: Antioxidant activity, occurrence and potential uses. *Food Chemistry* 99, 191-203.
- Barr DJS. 1988. How modern systematic relates to the rumen fungi. *Biosystems* 21, 351-356.
- Bhat TK, Singh B, Sharma OP (1998) Microbial degradation of tannins – A current perspective. *Biodegradation* 9, 343-357.
- Bieger J., Cermak R., Blank R., De Boer VCJ., Hollman PCH., Kamphues J., Wolfram S. 2008. Tissue distribution of quercetin in pigs after long term dietary supplementation. *Journal of Nutrition* 138, 1417-1420.
- Bravo L. 1998. Polyphenols: Chemistry, dietary sources, metabolism and nutritional significance. *Nutrition Reviews* 56, 317-333.
- Brooker JD., O'Donovan LA., Skene I., Clarke K., Blackall L., Muslera P. 1994. *Streptococcus caprinus* sp.nov., a tannin-resistant ruminal bacterium from feral goats. *Letters in Applied Microbiology* 18, 313-318.

- Cai H., Stewart A., Inoue M., Yuyama N., Hirata M. 2011. *Lolium*. Wild crop relatives: genomic and breeding resources, pp 165-173. [C Kole editor]. Springer-Verlag Berlin Heidelberg
- Cao G., Sofic E., Prior RL. 1997. Antioxidant and prooxidant behavior of flavonoids: structure-activity relationship. *Free Radical Biology and Medicine* 22, 749-760.
- Cappiello M., Giuliani L., Pacific G.M. 1990. Differential distribution of phenol and catechol sulphotransferases in human liver and intestinal mucosa. *Pharmacology* 40, 69-76.
- Carbonaro M., Grant G., Pusztai A. 2001. Evaluation of polyphenol bioavailability in rat small intestine. *European Journal of Nutrition*, 40, 84-90.
- Chopin J., Dellamonica G. 1988. C-glycosylflavonoids. The flavonoids; advances in research since 1980, pp 63-97. [JB Harborne editor]. Chapman & Hall, London.
- Cieslik E., Greda A., Adamus W. 2006. Contents of polyphenols in fruit and vegetables. *Food Chemistry* 94, 135-142.
- Cook NC., Samman S. 1996. Flavonoids-Chemistry, metabolism, cardioprotective effects, and dietary sources. *Journal of Nutritional Biochemistry* 7, 66-76.
- Cowan MM. 1999. Plant products as antimicrobial agents. *Clinical Microbiology Reviews* 12, 564-582.
- D'Archivio M., Filesi C., Di Benedetto R., Gargiulo R., Giovannini C., Masella R. 2007. Polyphenols, dietary sources and bioavailability. *Annali dell'Istituto Superiore di Sanità* 43, 4, 348-361.
- Day AJ., Mellon F., Barron D., Sarrazin G., Morgan MRA., Williamson G. 2001. Human metabolism of dietary flavonoids: Identification of plasma metabolites of quercetin. *Free Radical Research* 35, 941-952.
- De Boer VCJ, Dihal AA, Woude vander H., Arts ICW., Wolffram S., Alink GM., Rietjens IMCM., Keijer J., Hollman PCH. 2005. Tissue distribution of quercetin in rats and pigs. *Journal of Nutrition* 135, 1617-1618.
- Dehority BA. 2003. *Rumen microbiology*. University Press, Nottingham, United Kingdom.

- Dembinska-Kiec A., Mykkänen O., Kiec-Wilk B., Mykkänen H. 2008. Antioxidant phytochemicals against type 2 diabetes. *British Journal of Nutrition*, 99, 109–117.
- Déprez S., Brezillon C., Rabot S., Philippe C., Mila I., Lapierre C., Scalbert A. 2000. Polymeric proanthocyanidins are catabolized by human colonic microflora into low-molecular-weight phenolic acids. *Journal of Nutrition* 130, 2733-2738.
- Descalzo AM., Rossetti L., Grigioni G., Irurueta M., Sancho AM., Carrete J., Pensel NA. 2007. Antioxidant status and odour profile in fresh beef from pasture or grain-fed cattle. *Meat Science* 75, 299-307.
- Díaz MT., Velasco S., Cañeque V., Lauzurica S., Ruiz de Huidobro F., Pérez C., González J., Manzanares C. 2002. Use of concentrate or pasture for fattening lambs and its effect on carcass and meat quality. *Small Ruminant Research* 43, 257-268.
- Diebolt M., Bucher B., Andriantsitohaina R. 2001. Wine polyphenols decrease blood pressure, improve NO vasodilatation, and induce gene expression. *Journal of the American Heart Association* 38, 159-165.
- Dugas A.J., Castañeda-Acosta J., Bonin G.C., Price K.L., Fischer N.H., Winston G.W. 2000. Evaluation of the total peroxyl radical-scavenging capacity of flavonoids: structure-activity relationships. *Journal of Natural Products* 63, 327-331.
- Fraser ML., Mok GS., Lee AH. 2007. Green tea and stroke prevention: emerging evidence. *Complementary Therapies in Medicine* 15, 46-53.
- Frecnch P., Stanton C., Lawless F., O'Riordan EG., Monahan FJ., Caffrey PJ., Moloney A. 2000. Fatty acid composition, including conjugated linoleic acid, of intramuscular fat from steers offered grazed grass, grass silage, or concentrate-based diets. *Journal of Animal Science* 78, 2849-2855.
- Freedman JE, Parker CI, Li L., Perlman JA., Frei B., Ivanox V., Deak LR., Iafrati MD., Folts JD. 2001. Select flavonoids and whole juice from purple grapes inhibit platelet function and enhance nitric oxide release. *Circulation*, 103, 2792-2798.
- Georgé S., Brat P., Alter P., Amiot MJ. 2005. Rapid determination of polyphenols and vitamin C in plant-derived products. *Journal of Agriculture and Food Chemistry* 53, 1370-1373.

- Getachew G., Pittroff W., Putnam D.H., Dandekar A., Goyal S., DePeters E.J. 2008. The influence of addition of gallic acid, tannic acid, or quebracho tannins to alfalfa hay on *in vitro* rumen fermentation and microbial protein synthesis. *Animal Feed Science and Technology* 140, 444-461.
- Ghosh D. 2009. Vascular action of polyphenols. *Molecular Nutrition & Food Research* 53, 322-331.
- Gladine C., Rock E., Morand C., Bauchart D., Durand D. 2007. Bioavailability and antioxidant capacity of plant extracts rich in polyphenols, given as a single acute dose, in sheep made highly susceptible to lipoperoxidation. *British Journal of Nutrition* 98, 691-701.
- Goel G., Puniya AK., Singh K. 2005. Tannic acid resistance in ruminal streptococcal isolates. *Journal of Basic Microbiology* 45, 243-245.
- Ha Park K., Rodriguez-Montes de Oca GA., Bonello P., Lee KJ., Dabrowski K. 2009. Determination of quercetin concentrations in fish tissues after feeding quercetin-containing diets. *Aquaculture International* 17, 537-544.
- Hagerman AE., Riedl KM., Jones A., Sovic KN., Ritchard NT., Hartzfeld PW. 1998. High molecular weight plant polyphenolics (tannins) as biological antioxidants. *Journal of Agricultural and Food Chemistry* 46, 1887-1892.
- Halliwell B., Rafter J., Jenner A. 2005. Health promotion by flavonoids, tocopherols, tocotrienols, and other phenols: direct or indirect effects? Antioxidant or not? *American Journal of Clinical Nutrition* 81, 268-276.
- Harborne JB., Green PS. 1980. A chemotaxonomic survey of flavonoids in leaves of the Oleaceae. *Botanical Journal of the Linnean Society* 81, 155-167
- Harborne JB. 1994. *The flavonoids: advances in research since 1986*. [JB Harborne editor]. Chapman & Hall, London, U.K.
- Harborne JB., Williams CA. 2000. Advances in flavonoid research since 1992, *Phytochemistry* 55, 481-504.
- Haslam E., 1989. *Plant Polyphenols. Vegetable Tannins Revisited*, ed.: Cambridge University Press: Cambridge, U.K. 1989.
- Heim KE., Tagliaferro AR., Bobilya DJ. 2002. Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. *The Journal of Nutritional Biochemistry* 13, 572-584.

- Hollman PCH., Katan MB. 1997. Absorption, metabolism and health effects of dietary flavonoids in man. *Biomedicine & Pharmacotherapy* 51, 305-310.
- Holt RR., Lazarus SA., Cameron Sullards M., Yan Zhu Q., Schramm DD., Hammerstone JF., Fraga CG., Schmitz HH., Keen CL. 2002. Procyanidin dimer B2 [epicatechin-(4-8)-epicatechin] in human plasma after the consumption of a flavanol-rich cocoa. *American Journal of Clinical Nutrition* 76,798–804.
- Hou DX., Fujii M., Terahara N., Yoshimoto M. 2004. Molecular mechanisms behind the chemopreventive effects of anthocyanidins. *Journal of Biomedicine and Biotechnology* 5, 321-325.
- Houseknecht KL., Heuvel JPV., Moya-Camarena SY., Portocarrero CP., Peck LW., Nickel KP., Belury MA. 1998. Dietary conjugated linoleic acid normalizes impaired glucose tolerance in the Zucker diabetic fatty *fa/fa* rat. *Biochemical and Biophysical Research Communications* 244, 678-682.
- Ip C., Chin SF., Scimeca JA., Pariza MW. 1991. Mammary cancer prevention by conjugated dienoic derivative of linoleic acid. *Cancer Research* 51, 6118-6124.
- Juan ME., Maijó M., Planas JM. 2010. Quantification of *trans*-resveratrol and its metabolites in rat plasma and tissues by HPLC. *Journal of Pharmaceutical Biomedicine* 51, 391-398.
- Kaliora AC., Dedoussis GVZ., Schmidt H. 2005. Dietary antioxidants in preventing atherogenesis. *Atherosclerosis* 187, 1-17.
- Kalt W., Blumberg JB., McDonald JE., Vinqvist-Tymchuk MR., Fillmore SAE., Graf BA., O'Leary JM., Milbury PE. 2008. Identification of anthocyanins in the liver, eye and brain of blueberry-fed pigs. *Journal of Agricultural and Food Chemistry* 56, 705-712.
- Kepler CR., Tove SB. 1967. Biohydrogenation of unsaturated fatty acids. *The Journal of Biological Chemistry* 242, 5686-5692.
- Kerem Z., Chertrit D., Shoseyov O., Regev-Shoshani G. 2006. Protection of lipids from oxidation by epicatechin, *trans*-resveratrol, and gallic and caffeic acids in intestinal model systems. *Journal of Agricultural and Food Chemistry* 54, 10288-10293.
- Koes RE., Quattrochio F., Mol JNM. 2005. The flavonoid biosynthetic pathway in plants: function and evolution. *Bioessays* 16, 123-132.

- Kresty LA., Howell AB., Maureen B. 2011. Cranberry proanthocyanidins mediate growth arrest of lung cancer cells through modulation of gene expression and rapid induction of apoptosis. *Molecules* 16, 2375-2390.
- Kroon PA., Faulds CB., Ryden P., Williamson G. 1996. Solubilisation of ferulic acid from plant cell wall materials in a model human gut system. *Biochemical Society Transactions* 24, 384-384.
- Ksouri R., Megdiche W., Debez A., Falleh H., Grignon C., Abdelly C. 2007. Salinity effects on polyphenol content and antioxidant activities in leaves of the halophyte *Cakile maritime*. *Plant Physiology and Biochemistry* 45, 244-249.
- La Terra S., Marino VM., Manenti M., Licitra G., Carpino S. 2010. Increasing pasture intakes enhances polyunsaturated fatty acids and lipophilic antioxidants in plasma and milk of dairy cows fed total mix ration. *Dairy Science and Technology* 90, 687-698.
- Lambert JD., Yang CS. 2003. Cancer chemopreventive activity and bioavailability of tea and tea polyphenols. *Mutation Research* 523-524, 201-208.
- Lee JH., Cho KH., Lee KT., Kim MR. 2005. Antiatherogenic effects of structured lipid containing conjugated linoleic acid in C57BL/6J mice. *Journal of Agricultural and Food Chemistry* 53, 7295-7301.
- Liu ZQ., Ma LP., Zhou B., Yang L., Liu, ZL. 2000. Antioxidative affects of green tea polyphenols on free radical initiated and photosensitized peroxidation of human low density lipoprotein. *Chemistry and Physics of Lipids* 106, 53-63.
- Lopez V., Lindsay RC. 1993. Metabolic conjugates as precursors for characterizing flavor compounds in ruminant milks. *Journal of Agriculture and Food Chemistry* 41, 446-454.
- Luciano G., Monahan FJ., Vasta V., Biondi L., Lanza M., Priolo A. 2009a. Dietary tannins improve lamb meat colour stability. *Meat Science* 81, 120-125.
- Luciano G., Monahan FJ., Vasta V., Pennisi P., Bella M., Priolo A. 2009b. Lipid and colour stability of meat from lambs fed fresh herbage or concentrate. *Meat Science* 82, 193-199.
- Luciano G., Moloney AP., Priolo A., Röhrle, Vasta V., Biondi L., López Andrés P., Grasso S., Monahan FJ. 2011a. Vitamin E and polyunsaturated fatty acids in bovine muscle and the oxidative stability of beef from cattle receiving grass or concentrate-based rations. *Journal of Animal Science* 89, 3759-3768.

- Luciano G., Vasta V., Monahan FJ., López-Andrés P., Biondi L., Lanza M., Priolo A. 2011b. Antioxidant status, colour stability and myoglobin resistance to oxidation of *longissimus dorsi* muscle from lambs fed a tannin-containing diet. *Food Chemistry* 124, 1036-1042.
- Makkar HPS., Becker K., Abel H., Szegletti C. 1995a. Degradation of condensed tannins by rumen microbes exposed to quebracho tannins (QT) in rumen simulation technique (RUSITEC) and effects of QT on fermentative processes in the RUSITEC. *Journal of the Science of Food and Agricultural* 69, 495-500.
- Makkar HPS., Blümmel M., Becker K. 1995b. Formation of complexes between polyvinyl pyrrolidones or polyethylene glycols and tannins, and their implication in gas production and true digestibility in vitro techniques. *British Journal of Nutrition* 73, 897-913.
- Makkar HPS. 2003. Effects and fate of tannins in ruminant animals to tannins, and strategies to overcome detrimental effects of feeding tannin-rich feeds. *Small Ruminant Research* 49, 241-256.
- Manach C., Morand C., Crespy V., Demigné C., Texier O., Régéat F., Rémésy C. 1998. Quercetin is recovered in human plasma as conjugated derivatives which retain antioxidant properties. *Federation of European Biochemical Societies* 426, 331-336.
- Manach C., Scalbert A., Morand C., Rémésy C., Jiménez L. 2004. Polyphenols: food sources and bioavailability. *The American Journal of Clinical Nutrition* 79, 727-747.
- Manach C., Williamson G., Morand C., Scalbert A., Rémésy C. 2005. Bioavailability and bioefficacy of polyphenols in humans. I. Review of 97 bioavailability studies. *American Journal of Clinical Nutrition* 81, 230-242.
- Mancini RA., Hunt MC. 2005. Current research in meat color. *Meat Science* 71, 100-121.
- Martin B., Verdier-Metz I., Buchin S., Hurtaud C, Coulon JB. 2005. How do the nature of forages and pasture diversity influence the sensory quality of dairy livestock products? *Animal Science* 81, 205-212.
- Matsumoto H., Nakamura Y., Iida H., Ito K., Ohguro H. 2006. Comparative assessment of blackcurrant anthocyanins in rabbit and rat ocular tissues. *Experimental Eye Research* 83, 348-356.



- McDonald M., Mila I., Scalbert A. 1996. Precipitation of metal ions by plant polyphenols: optimal conditions and origin of precipitation. *Journal of Agricultural and Food Chemistry* 44, 599-606.
- McKay DL., Blumberg JB. 2002. The role of tea in human health: an update. *Journal of the American College of Nutrition* 21, 1-13.
- McSweeney CS., Palmer B., McNeill DM., Krause DO. 2001. Microbial interactions with tannins: nutritional consequences for ruminants. *Animal Feed Science and Technology* 91, 83-93.
- Melanson EL., Astrup A., Donahoo WT. 2009. The relationship between dietary fat and fatty acid intake and body weight, diabetes and the metabolic syndrome. *Nutrition & Metabolism* 55, 229-243.
- Min BR., Barry TN., Attwood GT., McNabb WC. 2003. The effect of condensed tannins on the nutrition and health of ruminants fed fresh temperate forages: a review. *Animal Feed Science and Technology* 106, 3-19.
- Moñino I., Martínez C., Sotomayor JA., Lafuente A., Jordán M.J. 2008. Polyphenolic transmission to Segureño lamb meat from ewes' diet supplemented with the distillate from rosemary (*rosmarinus officinalis*) leaves. *Journal of Agricultural and Food Chemistry* 56, 3363-3367.
- Morand-Fehr P., Fedele V., Decandia M., Le Frileux Y. 2007. Influence of farming and feeding systems on composition and quality of goat and sheep milk. *Small Ruminant Research* 68, 20-34.
- Moreno S., Scheyer T., Romano SC., Vojnov A.A. 2006. Antioxidant and antimicrobial activities of rosemary extracts linked to their polyphenols composition. *Free Radical Research* 40:2, 223-231.
- Mueller-Harvey I. 1999. Tannins: Their nature and biological significance. *Secondary plant products: antinutritional and beneficial actions in animal feeding* pp. 17-39 [JC Caygill and I Mueller-Harvey, editors]. Nottingham, Nottingham University Press.
- Mueller-Harvey I. 2001. Analysis of hydrolysable tannins. *Animal Feed Science and Technology* 91, 3-20.
- Mueller-Harvey I. 2006. Unravelling the conundrum of tannins in animal nutrition and health. *Journal of the Science of Food and Agriculture* 86, 2010-2037.

- Naczek M., Shahidi F. 2006. Phenolics in cereals, fruits and vegetables: Occurrence, extraction and analysis. *Journal of Pharmaceutical and Biomedical Analysis* 41, 1523-1542.
- Nardini M., Cirillo E., Natella F., Scaccini C. 2002. Absorption of phenolic acids in humans after coffee consumption. *Journal of Agriculture and Food Chemistry* 50, 5735-5741.
- Nelson KE., Pell AN., Schofield P., Zinder S. 1995. Isolation and characterization of an anaerobic ruminal bacterium capable of degrading hydrolyzable tannins. *Applied and Environmental Microbiology* 61, 3293-3298.
- Nimalaratne C., Lopes-Lutz D., Schieber A., Wu J. 2011. Free aromatic amino acids in egg yolk show antioxidant properties. *Food Chemistry* 29, 155-161.
- Nozière P., Graulet B., Lucas A., Martin B., Grolier P., Doreau M. 2006. Carotenoids for ruminants: from forages to dairy products (Review). *Animal Feed Science and Technology* 131, 418-450.
- Nuernberg K., Fischer A., Nuernberg G., Ender K., Dannenberger D. 2008. Meat quality and fatty acid composition of lipids in muscle and fatty tissue of Skudde lambs fed grass *versus* concentrate. *Small Ruminant Research* 74, 279-283.
- O'Connell JE., Fox PF. 2001. Significance and applications of phenolic compounds in the production and quality of milk and dairy products: a review. *International Dairy Journal* 11, 103-120.
- Paganga G., Rice-Evans CA. 1997. The identification of flavonoids as glycosides in human plasma. *FEBS Letters* 401, 78-82.
- Pérez-Magariño S., Ortega-Heras M., Cano-Mozo E. 2008. Optimization of a Solid-Phase Extraction method using copolymer sorbents for isolation of phenolic compounds in red wines and quantification by HPLC. *Journal of Agriculture and Food Chemistry* 56, 11560-11570.
- Perez-Maldonado RA., Norton BW. 1996. Digestion of <sup>14</sup>C-labelled condensed tannins from *Desmodium intortum* in sheep and goats. *British Journal of Nutrition* 76, 501-513.
- Porter LJ., Hrstich LN., Chan B. 1986. The conversion of procyanidins and rodolphinidins to cyanidin and delphinidin. *Phytochemistry* 25, 223-230.

- Porter LJ. 1992. Structure and chemical properties of the condensed tannins. *Plant polyphenols, synthesis, properties and significance* pp. 245-258 [Hemingway RW and Laks PE, editors]. Plenum, New York.
- Prache S., Priolo A., Grolier P. 2003. Persistence of carotenoid pigments in the blood of concentrate-finished grazing sheep: Its significance for the traceability of grass-feeding. *Journal of Animal Science* 81, 360-367.
- Prasain JK., Peng N., Dai Y., Moore R., Arabshahi A., Wilson L., Barnes S., Wyss JM., Kim H., Watts R. 2009. Liquid chromatography tandem mass spectrometry identification of proanthocyanidins in rat plasma after oral administration of grape seed extract. *Phytomedicine* 16, 233-243.
- Priolo A., Waghorn GC., Lanza M., Biondi L., Pennisi P. 2000. Polyethylene glycol as a means for reducing the impact of condensed tannins in carob pulp: Effects on lamb growth performance and meat quality. *Journal Animal Science* 78, 810-816.
- Priolo A., Micol D., Agabriel J. 2001. Effects of grass feeding systems on ruminant meat colour and flavour. A review. *Animal Research* 50, 185-200.
- Ramassamy C. 2006. Emerging role of polyphenolic compounds in the treatment of neurodegenerative diseases: A review of their intracellular targets. *European Journal of Pharmacology* 545, 51-64.
- Ramos S. 2008. Cancer chemoprevention and chemotherapy: Dietary polyphenols and signaling pathways. *Molecular Nutrition & Food Research* 52, 507-526.
- Realini CE., Duckett SK., Brito GW, Dalla Rizza M., De Mattos D. 2004. Effect of pasture vs. concentrate feeding with or without antioxidants on carcass characteristics, fatty acid composition, and quality of Uruguayan beef. *Meat Science* 66, 567-577.
- Reynaud A., Fraisse D., Cornu A., Farruggia A., Pujos-Guillot E., Besle JM., Martin B., Lamaison JL., Paquet D., Doreau M., Graulet B. 2010. Variation in content and composition of phenolic compounds in permanent pastures according to botanical variation. *Journal of Agriculture and Food Chemistry* 58, 5485-5494.
- Rice-Evans CA., Miller N.J., Paganga G. 1996. Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radical Biology & Medicine* 20, 7, 933-956.

- Rice-Evans CA., Miller N., Paganga G. 1997. Antioxidant properties of phenolic compounds. *Trend in Plant Science* 2, 152-159.
- Riedl KM., Hagerman E. 2001. Tannin-protein complexes as radical scavengers and radical skins. *Journal of Agriculture and Food Chemistry* 49, 4917-4923.
- Roux DG. 1992. Reflections on the chemistry and affinities of the major commercial condensed tannins in the context of their industrial use. *Plant polyphenols: synthesis, properties, significance*, pp. 7-39 [RW Hemingway and IPE Laks, editors]. Plenum, New York, NY.
- Russell JB., Wilson DB. 1998. Potential opportunities and problems for genetically altered rumen microorganisms. *Journal of Nutrition* 118, 271-279.
- Russell JB., Rychlik JL. 2001. Factors that alter rumen microbial ecology. *Science* 292, 1119.
- Saitoh S., Sato T., Harada H., Matsuda T. 2004. Biotransformation of soy isoflavone-glycosides in laying hens: intestinal absorption and preferential accumulation into egg yolk of equol, a more estrogenic metabolite of daidzein. *Biochimica et Biophysica Acta* 1674, 122-130.
- Sanbongi C., Suzuki N., Sakane T. 1997. Polyphenols in chocolate, which have antioxidant activity modulate immune functions in humans in vitro. *Cellular Immunology* 177, 129-136.
- Santos-Buelga C., Scalbert A. 2000. Proanthocyanidins and tannin-like compounds- nature, occurrence, dietary intake and effects on nutrition and health. *Journal of the Science of Food and Agriculture* 80, 1094-1117.
- Scalbert A., Williamson G. 2000. Dietary intake and bioavailability of polyphenols. *The Journal of Nutrition* 130, 2073-2085.
- Scalbert A., Morand C., Manach C., Rémésy C. 2002. Absorption and metabolism of polyphenols in the gut and impact on health. *Biomedicine and Pharmacotherapy* 56, 276-282.
- Scerra M., Luciano G., Caparra P., Foti F., Cilione C., Giorgi A., Scerra V. 2011. Influence of stall finishing duration of Italian Merino lambs raised on pasture on intramuscular fatty acid composition. *Meat Science* 89, 238-242.

- Sgorlon S., Stradaoli G., Zanin D., Stefanon B. 2006. Biochemical and molecular responses to antioxidant supplementation in sheep. *Small Ruminant Research* 74, 17-33.
- Shahrzad S., Bitsch I. 1998. Determination of gallic acid and its metabolites in human plasma and urine by high-performance liquid chromatography. *Journal of Chromatography B* 705, 87-95.
- Silberberg M., Morand C., Mathevon T., Besson C., Manach C., Scalbert A., Remesy C. 2006. The bioavailability of polyphenols is highly governed by the capacity of the intestine and of the liver to secrete conjugated metabolites. *European Journal of Nutrition* 45, 88-96.
- Skene IK., Brooker JD. 1995. Characterization of tannin acylhydrolase activity in the ruminal bacterium *Selenomonas Ruminantium*. *Anaerobe* 1, 321-327.
- Sly LI., Cahill MM., Osawa R., Fujisawa T. 1997. The tannin-degrading species *Streptococcus gallolyticus* and *Streptococcus caprinus* are subjective synonymous. *International Journal of Systematic Bacteriology* 47, 893-894.
- Smit LA., Baylin A., Campos H. 2010. Conjugated linoleic acid in adipose tissue and risk of myocardial infarction. *The American Journal of Clinical Nutrition* 92, 34-40.
- Smith AH., Zoetendal E., Mackie RI. 2005. Bacterial mechanisms to overcome inhibitory effects of dietary tannins. *Microbial Ecology* 50, 197-205.
- Spencer JPE., Chowrimootoo G., Choudhury R., Debnam ES., Srai SK., Rice-Evans C. 1999. The small intestine can both absorb and glucuronidate luminal flavonoids. *FEBS Letters* 458, 224-230.
- Stahl W., Berg van den H., Arthur J., Bast A., Dainty J., Faulks R.M., Gärtner C., Haenen G., Hollman P., Holst B., Kelly FJ., Polidoei MC., Rice-Evans C., Southon S., Vliet van T., Ribes JV., Williamson G., Asley SB. 2002. Bioavailability and metabolism. *Molecular Aspects of Medicine* 23, 39-100.
- Stangl V., Dreger H., Stangl K., Lorenz M. 2007. molecular targets of tea polyphenols in the cardiovascular system. *Cardiovascular Research* 73, 348-358.
- Steen RWJ., Lavery D., Kilpatrick J., Porter MG. 2003. Effects of pasture and high concentrate diets on the performance of beef cattle, carcass composition at equal growth rates, and the fatty acid composition of beef. *New Zealand Journal of Agricultural Research* 46, 69-81.

- Stevenson DE., Wibisono R., Jensen DJ., Stanley RA., Cooney JM. 2006. Direct acylation of flavonoid glycosides with phenolic acids catalysed by *Candida Antarctica* lipase B (Novozym 435). *Enzyme and Microbial Technology* 39, 1236-1241.
- Sultana B., Anwar F. 2008. Flavonols (kaempferol, quercetin, myricetin) contents of selected fruits, vegetables and medicinal plants. *Food Chemistry* 108, 879-884.
- Tajima K., Aminox RI., Nagamine T., Ogata K., Nakamura M., Matsui H., Benno Y. 1999. Rumen bacterial diversity as determined by sequence analysis of 16S rDNA libraries. *FEMS Microbiology Ecology* 29, 159-169.
- Tapiero H., Tew KD., Nguyen Ba G., Mathé G. 2002. Polyphenols: do they play a role in the prevention of human pathologies? *Biomedicine & Pharmacotherapy* 56, 200-207.
- Troszynska A., Bednarska A., Latosza A., Kozłowska H. 1997. Polyphenolic compounds in the seed coat of legume seeds. *Polish Journal of Food and Nutrition Sciences* 6, 37-45.
- Tu Y., Rochfort S., Liu Z., Ran Y., Griffith M., Badenhorst P., Louie GV., Bowman ME., Smith KF., Noel JP., Mouradov A., Spangenberg G. 2010. Functional analyses of *caffeic acid O-Methyltransferase* and *cinnamoyl-CoA-Reductase* genes from Perennial Ryegrass (*Lolium perenne*). *The Plant Cell* 22, 3357-3373.
- Turner KE., McClure KE., Weiss WP., Borton RJ., Foster JG. 2002. Alpha-tocopherol concentrations and case life of lamb muscle as influenced by concentrate or pasture finishing. *Journal of Animal Science* 80, 2513-2521.
- Urquiaga I., Leighton F. 2000. Plant polyphenol antioxidants and oxidative stress. *Biological Research* 33, 55-64.
- Van het Hof KH., Wiseman SA., Yang CS., Tjiburg LB. 1999. Plasma and lipoprotein levels of tea catechins following repeated tea consumption. *Proceedings of the Society for Experimental Biology and Medicine* 220, 203-209.
- Vasta V., Mele M., Scerra M., Luciano G., Lanza M., Priolo A. 2009a. Metabolic fate of fatty acids involved in ruminal biohydrogenation in sheep fed concentrate or herbage with or without tannins. *Journal of Animal Science* 87, 2674-2684.

- Vasta V., Priolo A., Scerra M., Hallett K.G., Wood J.D., Doran O. 2009b.  $\Delta^9$  desaturase protein expression and fatty acid composition of *longissimus dorsi* muscle in lambs fed green herbage or concentrate with or without added tannins. *Meat Science* 82, 357-364.
- Vasta V., Yañez-Ruiz DR., Mele M., Serra A., Luciano G., Lanza M., Biondi L., Priolo P. 2010. Bacterial and protozoal communities and fatty acid profile in the rumen of sheep fed a diet containing added tannins. *Applied and Environmental Microbiology* 76, 2549-2555.
- Vinson JA., Hao Y., Su X., Zubik L. 1998. Phenol antioxidant quantity and quality in foods: vegetables. *Journal of Agricultural and Food Chemistry*, 46 3630-3634.
- Vitaglione P., Fogliano V. 2004. Use of antioxidants to minimize the human health risk associated to mutagenic/carcinogenic heterocyclic amines in food. *Journal of Chromatography B* 82, 189-199.
- Wang Z., Huang Y., Zou J., Cao K., Xu Y., Wu J.M. 2002. Effects of red wine and wine polyphenol resveratrol on platelet aggregation *in vivo* and *in vitro*. *International Journal of Molecular Medicine* 9, 77-79.
- Wollgast J., Anklam E. 2000. Polyphenols in chocolate: is there a contribution to human health? *Food Research International* 33, 449-459.
- Wood JD., Enser M. 1997. Factors influencing fatty acids in meat and the role of antioxidants in improving meat quality. *British Journal of Nutrition* 78, 49-60.
- Wu LY., Zheng XQ., Lu JL. Liang YR. 2009. Protective effect of green tea polyphenols against ultraviolet B-induced damage to HaCaT cells. *Human Cell* 22, 18-24.
- Yamaguchi M. 2002. Isoflavone and bone metabolism: Its cellular mechanism and preventive role in bone loss. *Journal of Health Science* 48, 209-222.
- Yang A., Larsen TW., Tume RK. 1992. Carotenoid and retinol concentrations in serum, adipose tissue and liver and carotenoid transport in sheep, goats and cattle. *Australian Journal of Agricultural Research* 43, 1807-1809.
- Yang A., Brewster MJ., Lanari MC., Tume RK. 2002. Effect of vitamin E supplementation on  $\alpha$ -tocopherol and  $\beta$ -carotene concentrations in tissues from pasture- and grain-fed cattle. *Meat Science* 60, 35-40.
- Yang CS., Landau JM. 2000. Effects of tea consumption on nutrition and

health. *Journal of nutrition* 130, 2409-2412.

- Yoshida T., Hatano T., Ito H. 2000. Chemistry and function of vegetable polyphenols with high molecular weights. *Journal BioFactors* 13, 121-125.
- Yuasa AK., Hua JJ. Kennedy DO., Yuasa IM. 2002. Green tea extract inhibits angiogenesis of human umbilical vein endothelial cells through reduction of expression of VEGF receptors. *Life Sciences* 73, 1299-1313.
- Zern TL., Fernandez ML. 2005. Cardioprotective effects of dietary polyphenols. *Journal of Nutrition* 135, 2291-2294.
- Zheng W., Wang SY. 2001. Antioxidant activity and phenolic compounds in selected herbs. *Journal of Agricultural and Food Chemistry* 49, 5165-5170.